Preliminary Assessment of Nonfuel Minerals on the Texas Continental Shelf

by

Jeffrey G. Paine, Robert A. Morton, and William A. White

Prepared for

Louisiana Geological Survey Louisiana State University Minerals Management Service

Under Contract No. 14-12-0001-30404

Bureau of Economic Geology
The University of Texas at Austin
Austin, Texas 78713

W. L. Fisher, Director

September 1988

Table of Contents

Introduction	
Purpose of Study	1
Geological Framework	3
Late Wisconsinan Lowstand	
Late Wisconsinan and Holocene Sea-Level Rise	5
Existing Information	
Surface Samples	
Shallow Cores	7
Foundation Borings	9
Geophysical Data	11
Nonfuel Mineral Prospects	14
Shore-Parallel Sands	14
Shelf-Margin Deltas	18
Rio Grande Delta	
Delta 'A'	20
Delta 'B'	21
Delta 'C'	22
사이트 이 마음에게 되는 사람들은 아이가 있었다면 하는 그리는 그리는 아이들 바람이 하는데 모든 그리는 그렇다.	24
Brazos Area Gravels	28
Mustang Island Area Gravels	28
Heavy Minerals	

Potential Markets for Nonfuel Minerals	3	2
Beach Nourishment (Sand)	3	2
Demand		
Sources and Cost	3	3
Construction and Industry (Sand and Gravel)	3	4
Demand	3	4
Sources and Cost	3	5
Conclusions	3	5
Recommendations	3	6
References and Bibliography		
Appendix: Foundation Borings		

Figures

Figure 1.	Bathymetry and lease areas of the Texas continental shelf	2
Figure 2.	Late Pleistocene sea-level curve	4
Figure 3.	Location of short cores	8
Figure 4.	Location of foundation borings	10
Figure 5.	Location of high-resolution seismic surveys	13
Figure 6.	Distribution of shore-parallel sands	16
Figure 7.	Sabine and Heald Banks, eastern Texas shelf	17
Figure 8.	Location of Wisconsinan shelf-margin deltaic complexes	19
Figure 9.	Distribution of borings containing shallow sand	23
Figure 10.	Location of Wisconsinan streamcourses	25
Figure 11.	Distribution of foundation borings containing gravel	27
Figure 12.	Sand and gravel prospect offshore of Mustang Island, Texas	29
Figure 13.	Heavy mineral concentrations	31

DRAFT

Tables


Table 1.	Attributes of prospect types on the Texas continental shelf15
Table 2.	Heavy mineral suite, South Padre Island Area

Introduction

As part of a passive continental margin with a long depositional history, the Texas continental shelf has been richly endowed with mineral resources. Exploitation of oil and gas resources on the shelf extends back decades and the economic value of these deposits has long been recognized. The depositional setting that made the Texas offshore so rich in hydrocarbons has left it barren of exotic nonfuel minerals (polymetallic sulfides and ferromanganese cobalt crusts) such as those found in active tectonic settings near Hawaii and at the Juan de Fuca Ridge along the Pacific Northwest. Nonetheless, there are significant accumulations of potentially economic nonfuel minerals in the Texas Exclusive Economic Zone (EEZ). The most promising of these are sand and gravel deposited on the continental shelf during the sea-level fluctuations of the late Pleistocene and Holocene. Requirements for sand and gravel created by the burgeoning Texas coastal population and the need for compatible sands for beach nourishment projects such as those contemplated for rapidly eroding beaches at south Padre Island, the Brazos delta, and Galveston Island combined with the depletion of nearby onshore sand and gravel resources could make shelf mining operations economically feasible in the future. Before economic feasibility can be determined, however, the location, size, and character of potentially economic shelf deposits must be assessed.

Purpose of study

The primary objective of this study was to prepare a preliminary assessment of nonfuel mineral resources of the EEZ (fig. 1) from the gulf shoreline to near the shelf edge (approximately 200 m water depth). Several steps are required to satisfy this objective, including (1) inventory available geological information, both published (a bibliography) and unpublished (high-resolution seismic surveys, piston cores, vibracores, seafloor samples, and foundation borings); (2) locate potentially economic offshore deposits (prospects) using available data; (3) characterize these prospects as accurately as possible



Bathymetry and lease areas of the Texas continental shelf. HI: High Island; HIE: High Island East Addition; HIS: High Island South Addition; HIES: High Island East Addition South Extension; G: Galveston; GS: Galveston South Addition; B: Brazos; BS: Brazos South Addition; Mal: Matagorda Island; Mul: Mustang Island; MulE: Mustang Island East Addition; NPI: North Padre Island; NPIE: South Padre Island; Sou Padre Island East Addition.

Figure 1.

with existing data, including areal extent, thickness, and sediment composition and texture; and (4) determine whether exploitation of the prospect is economically feasible if sufficient data exist; if data are insufficient, recommend a research program that would help determine whether poorly-known prospects are economic.

Geological Framework

Knowledge of the geology of the Texas continental shelf is helpful in predicting and understanding the distribution of nonfuel minerals. Currently, very little deposition of economically important minerals takes place on the shelf beyond the nearshore zone. However, there are significant concentrations of sand and gravel that occur far offshore. These relict deposits owe their placement to large-scale fluctuations in sea level during the Quaternary (fig. 2); most important for this study are the sea-level lowstand during last glaciation (late Wisconsinan) and the subsequent sea-level rise as the glaciation ended (late Wisconsinan and early to middle Holocene).

Late Wisconsinan Lowstand

As sea level was falling during the onset of the late Wisconsinan glaciation, shelf-phase deltas prograded across the Texas continental shelf leaving relatively thin and discontinuous deltaic deposits. At the shelf edge, however, these deltas encountered steeper gradients that allowed them to reach thicknesses of 90 m or more (Suter and Berryhill 1985). These shelf-margin deltas are largely composed of sand and mud, with sand more abundant in proximal, shallow-water areas such as near distributary channels and at distributary mouth bars. Four major shelf-margin deltaic complexes have been located offshore from Texas (Suter and Berryhill 1985); though they now occur in water that is too deep (more than 90 m) for the deposits to be economic, they represent substantial sand resources that may someday be exploited.




Figure 2. Late Pleistocene sea-level curve. Adapted from Moore (1982).

Also during the late Wisconsinan lowstand, streams associated with the shelf-margin deltas coursed across the Texas continental shelf. As they do onshore today, these streams transported and deposited large quantities of sand and gravel on the shelf. Similar deposits in the Texas coastal zone are commonly exploited for use in the construction industry.

Late Wisconsinan and Holocene Sea-Level Rise

As sea level rose at the end of the last glaciation, thin transgressive deposits composed of reworked deltaic sands capped the shelf margin deltas, and deltaic, estuarine, and finally marine deposits filled the late Wisconsinan streamcourses. These deposits covered fluvial sands and gravels that became the basal postglacial deposits. During minor stillstands and reversals of sea-level rise, waves and longshore currents redistributed relict sediment and sediment brought to the coast by rivers to produce elongate, shore-parallel sandy deposits that were subsequently submerged and partly reworked when sea level began rising again. These shore-parallel sands represent a potentially compatible material for nourishment of Texas beaches because they were formed from sediment similar to beach sand and because they were transported by processes similar to those acting on beaches today.

Existing Information

Several types of data have been collected that can be used to directly or indirectly determine the distribution, texture, and composition of surface and shallowly-buried shelf sediments. These data, including surface samples, pipe and box cores, foundation borings, and high-resolution seismic lines, each have advantages and disadvantages in terms of assessing nonfuel mineral resources. More surface samples have been taken from the Texas continental shelf than any other type of sediment sample, but penetration is only a few centimeters and little information on the vertical extent of potential nonfuel mineral

deposits can be gained from these samples. Pipe cores achieve slightly greater penetration (up to a few meters), but are not as widespread as grab samples. Foundation borings, commissioned by petroleum companies in preparation for drilling or production activities, are perhaps best suited for determinations of vertical sediment distribution because they extend 100 m or more into the subsurface. However, uneven distribution across the Texas continental shelf and questionable visual descriptions of sediment reduce their usefulness. High-resolution seismic profiles are most useful in locating structural elements and constructing three-dimensional models of depositional systems, but they provide only indirect information on sediment texture.

Surface Samples

Investigations of surface sediment distribution along the northwest Gulf of Mexico (Texas and western Louisiana continental shelves) during Project 51 of the American Petroleum Institute (API) included collection of about 1,350 dredge samples and short gravity cores (Curray 1960). About two thirds of these samples were obtained by Scripps Institution of Oceanography in Texas offshore waters out to depths of 200 m (fig. 1). Analysis of these shallow sediments included grain size determinations (Curray 1960) and the heavy mineral suite (van Andel 1960; van Andel and Poole 1960). Textural data from cores and dredge samples were combined to produce a sediment distribution map for the Texas and Louisiana continental shelves.

In the mid-1970's, the United States Geological Survey (U.S.G.S.) completed a study of the south Texas outer continental shelf, consisting of the South Padre Island, North Padre Island, Mustang Island, and Matagorda Island areas (Berryhill 1976; Berryhill et al. 1976). During this study, many types of geological data were collected from near the State-Federal ownership boundary out to a depth of about 200 m, including surface samples, shallow cores, and seismic reflection profiles. Surface samples were taken at 264

stations along 27 dip-oriented transects; most of these samples were analyzed for grain size distribution and heavy mineral content.

Surface samples and seismic profiles on State-owned submerged lands (to 16 km offshore) were collected and analyzed by the Bureau of Economic Geology in the mid- to late 1970's (McGowen and Morton 1979). About 3,500 surface samples were collected 1.6 km apart on the Texas continental shelf; these samples were analyzed for grain-size distribution and for several geochemical constituents (White et al. 1983, 1985a, 1985b, 1985c, 1986a, 1986b, 1987).


There have been many other studies of surface sediment distribution, but these studies, such as the Sabine Bank area (Nelson and Bray 1970) and offshore from the Brazos River (Nienaber 1963) cover relatively small areas. Adequate data exist for accurate regional characterizations of sediment texture only for the south Texas outer continental shelf and the State-owned inner continental shelf.

Shallow Cores

Pipe cores, piston cores, box cores, and vibracores are characterized by relatively shallow penetration into the subsurface, ranging from a few centimeters to a few meters. They are more useful than surface samples for determining vertical dimensions of potentially economic deposits, but few systematic studies of the continental shelf have been completed. Most notable among the completed studies are one covering the entire Texas continental shelf (Curray 1960; see previous section), another covering the south Texas outer continental shelf (Berryhill et al. 1976), and a third focusing on Sabine and Heald Banks in the High Island area (Nelson and Bray 1970).

As a part of the South Texas Outer Continental Shelf study, the U.S.G.S. collected pipe cores at 90 stations and box cores at 74 stations (fig. 3) within the South Padre Island, North Padre Island, Mustang Island, and Matagorda Island areas (Berryhill et al. 1976).

Box cores penetrated about 40 cm of sediment, whereas pipe cores penetrated from less



Location of short cores on the Texas continental shelf. Figure 3.

than 30 cm to more than 2 m. Textural analyses were not done on samples from these cores, but sand lenses and other sedimentary features were noted.

In the Sabine and Heald Bank area, Nelson and Bray (1970) collected numerous short gravity cores as well as 12 rotary cores (8 of which were taken in the High Island area of the Texas continental shelf). The rotary cores penetrated 1 to 25 m below the surface, with sediment recovery ranging from less than 25 percent to 100 percent. Textural analyses were completed for 750 surface samples; textural characteristics of cored sediments were estimated from drilling characteristics and recovered sediment. In the same general area, 18 vibracores were collected in a single lease block during a study of potential archeological resources (Pearson et al. 1986). These cores achieved penetrations ranging from 5 to 12 m, with recovered lengths ranging from 3 to 6 m. Many analyses, including grain size and geochemistry, were conducted on sediment from these cores.

Foundation Borings

Foundation borings are perhaps the most useful tool for documenting the vertical distribution of near-surface sediment on the Texas shelf. These borings, commonly obtained by engineering firms under contract to oil companies preparing to drill offshore wells or build production platforms, may extend more than 100 m below the seafloor. A computerized database created at the Bureau of Economic Geology contains 410 borings from the Texas continental shelf (fig. 4 and appendix). Reports of these borings were obtained from the Houston offices of McClelland Engineers and PSI, two of the major engineering firms operating on the Texas continental shelf. The reports include visual descriptions, textural analyses, and various engineering properties of the sediments encountered in the borings. Attributes of the borings entered in the database include location, water depth, length of boring, visual description of sediments encountered, and depths of boundaries between sedimentary types. Water depths for borings in the database range from nearshore borings in 5 m of water to shelf-margin borings in 132 m of water.




Figure 4. Location of foundation borings on the Texas continental shelf.

Subsurface penetration ranges from 5 to 170 m, with most borings extending at least 30 m below the seafloor.

Drawbacks of the foundation borings are (1) uneven geographic distribution and (2) uncertainty about the accuracy of the visual sediment descriptions. Although foundation borings are located in every lease area on the Texas continental shelf, boring distribution is controlled by hydrocarbon leasing and some lease areas have more borings than others. The geographic distribution is most dense in the High Island, Matagorda, Brazos, and Galveston areas; foundation borings are sparse in the North and South Padre Island areas (fig. 4). Because sediment descriptions of the borings are commonly confirmed by textural analyses, these records should be adequate for locating significant sand and gravel deposits in the subsurface, but not for evaluating resource quality.

Geophysical Data

There are several regional geophysical surveys of the Texas continental shelf that help locate potential nonfuel mineral resources. These surveys consist of high-resolution seismic reflection data acquired by various State and Federal agencies and by oil companies. The surveys are useful because features such as buried stream channels, filled river valleys, and drowned shelf-margin deltas can be located with this information. Although lithologic information is not obtained directly from seismic data, knowledge of the types of sediment characteristic of depositional systems located by seismic profiles gives strong indirect information about lithology.

Approximately 6,500 km of high-resolution seismic data was collected on the Texas inner continental shelf (nearshore to 16 km offshore) in a cooperative effort between the U.S.G.S. and the Bureau of Economic Geology in the mid-1970's (McGowen and Morton 1979). The seismic data collected during this project consisted of 232 dip lines spaced 2.4 km apart and tied together by 2 strike lines spaced 6.5 km apart. The primary energy source was an 800 joule minisparker; some 3.5 khz subbottom profiler data were also


collected. The U.S.G.S. also collected more than 9,200 km of high-resolution seismic data on the south Texas outer continental shelf between 1974 and 1976 (fig. 5). Most of this additional seismic coverage (8,900 km) was completed using either a 1,000 to 1,500 joule Acoustipulse source or a 10,000 joule sparker source (Berryhill et al. 1976).

Regional high-resolution coverage of the Texas shelf edge and upper slope (fig. 5) was completed by the U.S.G.S. as part of a gulfwide continental slope study. Using a 400 to 1,000 joule minisparker source and a subbottom profiler, Texas shelf edge and slope (200 to 1,000 m water depth) coverage was obtained on an approximate 9-km grid (Berryhill 1987a; Suter and Berryhill 1985).

Although most of the seismic coverage was acquired for the Louisiana shelf, two high-resolution seismic surveys commissioned by the U.S.G.S. in 1979 and 1980 covered part of the Texas continental shelf (fig. 5). In 1979, a 400 joule sparker source and a 7 khz subbottom profiler were used to collect shallow subsurface information from the mid shelf to the upper slope in the High Island East Addition South Extension (Berryhill 1984b); in 1980, a similar system was used to extend the coverage from the mid-shelf to inner shelf in the High Island East area (Berryhill 1984a). North-south and east-west lines were completed on a 5.5-km grid over this area.

Regional high-resolution seismic coverage over most of the Galveston Area South Addition was conducted by Texaco in 1972 and 1973 (fig. 5). This survey consisted of 19 north-south and 5 east-west lines covering an area of about 65 by 65 km (Lewis 1984). Unlike other regional studies listed above that used sparker or Acoustipulse energy sources, this survey used a 650 cm³ airgun source.

Many other geophysical studies have been completed on the Texas continental shelf, including side scan sonar, magnetometer surveys, gravimetric surveys, and a multitude of other high-resolution seismic surveys. Most of these other seismic surveys, such as those required for lease block geohazard analysis, are of little practical use for regional characterizations of potential nonfuel mineral resources because of the effort



Location of high-resolution seismic surveys on the Texas continental shelf and slope. Figure 5.

required to obtain them and their limited geographic extent. However, they could be extremely useful once promising sites have been located.

Nonfuel Mineral Prospects

Several potentially economic nonfuel mineral deposits have been located in previous investigations and through analysis of existing offshore data. Because of the sparseness of the data, the extent and quality of the deposits are generally not well known. The potentially economic deposits can be subdivided by type and potential use (table 1). These types include shore-parallel deposits composed of sand and some shell fragments (shoreline-like deposits), thick and lobate shelf-margin deposits composed of sand and mud (shelf-margin deltas), predominantly shore-normal deposits composed of sand and, in places, gravel (ancient fluvial systems), and heavy mineral concentrations on the south Texas shelf (transgressive sheet sands). Possible economic uses for these deposits include reconstruction of eroding beaches (beach nourishment), landfill, roadbase, and in the production of various concrete products. Potentially economic concentrations of heavy minerals offshore from the Rio Grande have several industrial uses.


Shore-Parallel Sands

Many shore-parallel sandy deposits on the Texas continental shelf (fig. 6) are interpreted as shoreline or nearshore sands that mark late Pleistocene or early Holocene positions of sea level. These sandy deposits are likely to be suitable for nourishment of eroding Texas beaches because they formed from processes, conditions, and sediments similar to those forming Texas' present-day beaches. As such, they are probably relatively mature sediments composed mainly of quartz.


Heald and Sabine Banks, interpreted as submerged shoreline and shallow marine sands, are located 40 to 50 km offshore in the High Island area of the upper Texas coast in water depths of 6 to 17 m (fig. 7). These elongate surface sand deposits roughly parallel

Attributes of prospect types on the Texas continental shelf. BN = beach nourishment; Ind = industrial uses; Con = construction. Table 1.

Prospect Type	Resource	Best Example	Time of Formation	Suit B N	Suitable for BN Ind Con	for	Comments	
Shore-parallel sands	sand	Sabine and Heald Banks	post-Wisconsinan	yes	yes yes yes	yes	May contain shell	
Streamcourses	sand, gravel	Mustang Island Area	late Wisconsinan to early Holocene	Ou	no yes yes	yes	May be covered by several meters of overburden	
Shelf-margin deltas	sand	Rio Grande delta	Wisconsinan	00	i i ou	ċ	May contain silt and clay	
Transgressive sands	sand, heavy minerals	South-Padre Island Area —post-Wisconsinan—	post-Wisconsinan	yes yes yes	yes	yes	May contain shell; Rio Grande area promising	



Distribution of shore-parallel sands at the surface or shallowly buried on the Texas continental shelf. Compiled from Grady (1970); Berryhill et al. (1976); McGowen and Morton (1979); and Nelson and Bray (1970). Figure 6.



Sabine and Heald Banks, eastern Texas shelf. Location and areal extent of sandy deposits from Nelson and Bray (1970); vertical distribution of sediment from two cores published by Nelson and Bray (1970) and five foundation borings. Figure 7.


the present shoreline; Sabine Bank occurs east of Heald Bank and extends into offshore Louisiana. Much of what is known about the geology of the banks is based on short cores, grab samples, and a sonoprobe survey (Nelson and Bray 1970). In addition, the eastern part of Sabine Bank falls within an area studied by Berryhill et al. (1984) using high-resolution seismic data; the area north of Sabine Bank was studied by Pearson et al. (1986) using seismic data and vibracores. Nine soil foundation borings also penetrate the area. From these data, it is known that sands associated with these banks cover more than 1,000 km² of the sea floor and range up to 8 m thick. Cores and seismic records indicate an average thickness of about 3 m. Multiplying these values yields an estimated volume of more than 3 billion m³ of sediment within Heald and Sabine Banks. Grain size analyses performed by Nelson and Bray (1970) indicate the deposit is composed dominantly of fine to very fine sand, similar to most Texas beach sand (Bullard 1942).

Shelf-Margin Deltas

Deltas constructed at the outer shelf margin and upper continental slope during late Pleistocene lowstands of sea level contain significant accumulations of sand. Four major shelf-margin deltas have been located at the edge of the Texas shelf (fig. 8), from the ancestral Rio Grande delta to the south to deltas 'A', 'B', and 'C' to the east (Berryhill 1987b; Berryhill and Suter 1987; Morton and Price 1987; Suter and Berryhill 1985; Lewis 1984). Although these deltas each cover hundreds of square kilometers, they each also contain abundant silt and clay. Greatest concentrations of sand are likely to be found near the top of the deposits and also in the more shallow proximal deltaic areas. All of the deltas are too distant from potential markets to be economic at the present time.

Rio Grande Delta

The Rio Grande delta, located in the South Padre Island East Addition area in water depths of 45 to 200 m (fig. 8), is the largest of the shelf-margin deltaic complexes on the



Location of Wisconsinan shelf-margin deltaic complexes (after Suter and Berryhill 1985). Figure 8.

Texas continental shelf. As a sand resource, it has potential uses in the construction industry and, depending on size characteristics, could be used for beach nourishment along south Padre Island. The nearest potential market is south Padre Island, which is 55 to 100 km away.

Knowledge of the Rio Grande delta comes primarily from high-resolution seismic surveys, box cores, and pipe cores acquired by the U.S.G.S. during the South Texas Outer Continental Shelf project (Berryhill et al. 1976). Additional information on the distal edge of the delta was collected during a U.S.G.S. seismic survey of the Gulf of Mexico continental slope (Berryhill 1987b; Suter and Berryhill 1985). The shelf-margin part of the delta covers about 65 km along the shelf and about 35 km across the shelf, not including the part extending into Mexico. Pipe cores extending as much as 2 m into the delta encountered abundant shelly sand and interbedded mud and sand, but give no indication of the maximum sand thickness. The only foundation boring that penetrates the shelf-margin delta encountered 36 m of sand, silty sand, and sandy silt overlying a coarsening-upward sequence of clay to sandy silt. This sequence is probably typical of most of the shelfmargin delta, although sediments would tend to be thinner and generally coarser toward shore and thicker and generally finer toward the shelf margin. Interpretations from seismic data indicate that the delta thickness increases seaward from 20 m near the landward limit to more than 100 m near the edge of the continental shelf. Most of the delta, however, is probably composed of muddy sediments (silt and clay).

Delta 'A'

Delta 'A' is the southwesternmost of three smaller deltaic complexes located on the southern edge of the eastern Texas continental shelf (fig. 8) and may represent lowstand deltaic deposits associated with the ancestral Colorado or Brazos Rivers. Water depths over the shelf-margin phase of this deltaic complex range from about 60 m to 200 m. The delta is primarily composed of sand and mud; as a potential sand resource for beach

nourishment and the construction industry, the nearest market is Galveston at a distance of about 185 km.

The delta is located within the Brazos South and Mustang Island East Addition areas, but few cores penetrate it. Knowledge of this delta mainly comes from high-resolution seismic data collected by the U.S. Geological Survey (Berryhill et al. 1976; Suter and Berryhill 1985). From these data, it is known that the delta is composed of two vertically-stacked lobes with a total thickness of up to 100 m covering an area of about 50 km along strike and about 16 km along dip. One core taken from the eastern flank of the delta in about 200 m of water encountered a 50-m thick coarsening-upward sequence of clay to sandy clay (Sidner et al. 1978). A second core, located updip from the delta, sampled coarser sediments from the contributing fluvial system. Surficial shelly sands cover much of the delta (Suter and Berryhill 1985).

Delta 'B'

Delta 'B' is a Wisconsinan shelf-margin delta located mostly in the Galveston Area South Addition (fig. 8) in water depths of 60 to 200 m. It is composed of sand and mud; the nearest potential market for the sands contained in the delta is Galveston, which is located about 135 km away.


Knowledge of this delta comes from high-resolution seismic surveys conducted by the U.S.G.S. (Suter and Berryhill 1985) and Texaco (Lewis 1984). In addition, four foundation borings penetrate the seafloor on or near the delta. Delta B extends about 65 km in an east-west direction (along strike) and about 16 km in a north-south direction (along dip); maximum thickness is about 60 m. One boring in relatively deep water (130 m) encountered clay from the surface to a depth of 130 m; other borings in shallower water (78 to 102 m) encountered silty fine sand with thicknesses varying from 2 to 40 m. Surficial sediments of sand and silty fine sand cover the shallower-water parts of the delta (Suter and Berryhill 1985).

Delta 'C'

This multi-lobe deltaic complex is located at the southern edge of the High Island South Addition and High Island South Addition East Extension areas (fig. 8), extending east-west (strike) about 65 km and north-south (dip) about 16 km (Suter and Berryhill 1985). At least two U.S.G.S. seismic surveys have encountered part of this delta, including the Gulf of Mexico continental slope survey and the 1979 mid- to outer shelf survey which covered the eastern part of the delta (Berryhill et al. 1984). Direct knowledge of sediments associated with this delta has been obtained from seven foundation borings that penetrate it. Like the other shelf-margin deltas, this potential sand resource occurs in water depths of 60 to 200 m and is relatively remote from potential markets. The nearest local market is Galveston, about 160 km distant.

Analysis of seismic records indicates that this deltaic complex reaches a maximum thickness of more than 140 m (Suter and Berryhill 1985). Five of the seven foundation borings in the area penetrate silty fine sand, with three borings encountering sands at the surface. These surface sand deposits range from 4 to 12 m thick. These sands cap a coarsening-upward sequence of sediment that includes clays at the base, overlain by interbedded clay and silty sands. The uppermost coarsening-upward sequence is 41 to 82 m thick. This sequence is typical of deltaic depositional systems and reinforces the original seismic interpretation.

Foundation borings that penetrate the seafloor in the vicinity of the four shelf-margin deltaic complexes indicate that substantial thicknesses of sand are found near the surface of these deltas (figs. 8 and 9). Most of the foundation borings that encountered more than 7.5 m of sand in the upper 15 m of the boring are located within these deltas or their updip stratigraphic equivalents (fig. 9).




Distribution of foundation borings (shaded) containing more than 7.5 m of sand in the top 15 m of sediment below the seafloor. Figure 9.

23

Streamcourses and Valley Fill

Many sand and gravel quarries in the Texas coastal zone are located along major streams. During the Wisconsinan glaciation, extensions of these streams flowed across the exposed continental shelf to the Gulf of Mexico. Along their courses these streams deposited sediments similar to those found in onshore quarries. At the end of the last glaciation, rising sea level caused these lowstand channels and incised valleys to fill with a transgressive sequence of sediment, from relatively coarse fluvial channel deposits (sand and some gravel) to finer deltaic deposits (sand and mud) to generally fine estuarine deposits (mostly mud) and finally to open gulf deposits (shelf muds and possibly nearshore sands). Probably the only significant shallowly-buried gravel deposits on the Texas continental shelf will be found with the sands occurring in these submerged streamcourses. Unfortunately, these gravels will be at the base of the transgressive sequence, which may be tens of meters thick. The greatest chance for economically attractive sand and gravel deposits will be on the inner continental shelf, where shallow water, a relatively thin overburden, and proximity to potential markets will minimize the costs of extraction and transportation. In addition to exploitation difficulties arising from greater water depths farther offshore, gravels are likely to be less abundant and more deeply buried under late Pleistocene and Holocene deltaic, estuarine, and marine sediments.

High-resolution seismic surveys indicate that many kilometers of ancient streamcourses are preserved on the Texas continental shelf (fig. 10). Various seismic and coring surveys of Texas bays have shown where many of Texas' rivers entered the continental shelf during the last glaciation (Rehkemper 1969; Behrens 1963; Wright 1980), but these streams generally did not flow straight across the shelf to build shelf-margin deltas (Suter and Berryhill 1985). Seismic reflection data collected and interpreted by the U.S.G.S. on the south Texas outer continental shelf (Berryhill 1980, 1981a, 1981b) show several streams entering the shelf from the present-day Rio Grande to Matagorda Bay; all the streams from Copano Bay southward flowed to the Rio Grande delta (fig. 10). Seismic




Location of Wisconsinan streamcourses. Compiled from Aten (1983), Berryhill (1980, 1981a, and 1981b), Nelson and Bray (1970), and Suter and Berryhill (1985). Figure 10.

data collected by the U.S.G.S. from the eastern Texas shelf also show numerous submerged streamcourses (Berryhill et al. 1984; Suter 1987).

Relatively little seismic data exist for the continental shelf in the Brazos, Galveston, and western High Island areas (fig. 5) that would help locate ancient courses of the Trinity, Brazos, Sabine/Neches, and possibly the Colorado rivers. The Trinity, Brazos, and Colorado are currently three of the largest streams in Texas, and substantial shelf sand and gravel deposits are associated with ancient channels of these streams. Nelson and Bray (1970), using cores and sonoprobe data, found an ancient valley of the Sabine/Calcasieu system, which turns abruptly southwestward near the confluence of the Sabine and Calcasieu rivers (fig. 10). Pearson et al. (1986) studied sediments associated with this paleovalley with high-resolution seismic surveys and several vibracores. Using cores, seismic surveys, and bathymetry, Aten (1983) constructed a paleogeographic map showing the late Pleistocene and early Holocene inner shelf courses of the Sabine/Neches, Trinity, and Calcasieu rivers, and showed the ancient Sabine/Calcasieu streamcourse merging with the Trinity streamcourse 50 to 65 km southeast of Galveston (fig. 10). It is not known where these combined streams flowed gulfward from this inferred confluence. Even less is known about late Pleistocene and early Holocene streamcourses of the Brazos and Colorado rivers, although one ancient course trending southward from Matagorda Bay (Berryhill 1981b) may be related to the Colorado River system. In summary, there is very little regional seismic data in an area of the shelf that has a high probability of containing significant fluvial sand and gravel deposits.

In addition to inferred fluvial sand and gravel deposits located along streamcourses revealed by seismic data, gravels which are almost certainly fluvial in origin have been encountered in foundation borings on the Texas shelf. Of the 26 borings that encountered gravels (fig. 11), 17 were located in the Brazos or Galveston areas and are probably associated with the ancient Brazos, Colorado, and possibly the Trinity rivers. All but two of these gravel deposits were too thin or too deeply buried to be economical. One of the



Distribution of foundation borings containing gravel. Depth below seafloor and thickness of gravelly strata included for borings that encountered gravel. Figure 11.


potentially economic deposits is located in the Brazos Area and the other is in the Mustang Island Area (figs. 11 and 12)

Brazos Area Gravels

Although 9 of 75 foundation borings in the Brazos and Brazos South Areas encountered gravels, only one of these borings penetrated a significant thickness of gravelly sediment with less than 30 m of overburden. This boring, located in Brazos Block 409 about 14 km offshore from Matagorda Peninsula (fig. 11), extends 113 m into the subsurface in 19 m of water. Fine to coarse sands containing gravel and shell fragments occur between subsurface depths of 13 to 19 m; these deposits are overlain by clay containing wood fragments, sandstone fragments, and calcareous nodules. The areal extent of this gravel-bearing deposit is not known, but nearby borings in adjacent lease blocks to the northeast and southeast contain thin gravel lenses with overburden thicknesses of 8 and 33 m.

Mustang Island Area Gravels

Perhaps a more promising gravel deposit was encountered in a foundation boring located about 8 km offshore of Mustang Island in Mustang Island Block 772 (figs. 11 and 12). This boring, taken in 16 m of water, penetrated 12 m of sandy gravel underneath 16 m of a fining-upward sequence that included 12 m of dominantly silty fine sand. This boring apparently encountered basal transgressive valley-fill deposits near the confluence of ancestral Nueces, Aransas, and possibly Mission rivers, which were located through interpretation of high-resolution seismic reflection surveys conducted by the U.S.G.S. (Berryhill 1981a). Although nearby foundation borings located off the axis of these streamcourses did not encounter gravel, it is likely that similar deposits exist both upstream and downstream from this boring. Numerous sand and gravel quarries are operating in similar deposits along the Nueces River near Corpus Christi.




Sand and gravel prospect offshore of Mustang Island, Texas. Areal extent of Wisconsinan streamcourses from Berryhill (1981). Figure 12.

Heavy Minerals

The only systematic determination of heavy mineral content of Texas shelf sediments was conducted by the U.S.G.S. on the south Texas outer continental shelf (Berryhill et al. 1976). During this survey, 276 grab samples spaced on an approximate 5-km grid were analyzed for total heavy mineral content in the sand fraction. Heavy mineral concentrations determined during this study ranged from only a trace to 32 percent by weight. Heavy mineral concentrations generally increased southward toward the Rio Grande delta, with most of the higher concentrations (greater than 2 percent by weight) recorded off south Padre Island (fig. 13). Heaviest concentrations are located 16 to 72 km offshore in water depths of 20 to 100 m. The thickness of sediments with high concentrations of heavy minerals is not known because grab samples only penetrate a few centimeters below the seafloor. However, the heavy mineral deposits are probably relatively thin (less than one meter) because they occur in transgressive sandy sediments that cap the Rio Grande delta complex.

Heavy minerals in the grab samples collected by the U.S.G.S. were not identified because extensive work with the shelf heavy mineral suite was done during API Project 51 (Curray 1960; van Andel 1960; van Andel and Poole 1960). Analyses of seven samples (table 2) taken in the vicinity of the heavy mineral concentrations indicate that the suite is dominated by hornblende, epidote, zircon, and garnet (van Andel and Poole 1960). Minor amounts of staurolite, tourmaline, and kyanite are also present. These minerals have largely been brought to the shelf by the Rio Grande, which carries an assemblage of heavy minerals (table 2) similar to that found on the shelf (van Andel 1960).



Heavy mineral concentrations (in weight percent) in the sand fraction of seafloor samples collected from the Matagorda Island, Mustang Island, North Padre Island, and South Padre Island lease areas (adapted from Berryhill et al. 1976). Figure 13.

Table 2. Heavy mineral suite in sand fraction of Rio Grande fluvial deposits and on the continental shelf, South Padre Island Area. River abundances are averaged; shelf abundance given as range among seven samples. Data from van Andel (1960) and van Andel and Poole (1960).

	River Abundance	Shelf Abundance
Mineral	(percent)	(percent)
Epidote	22	4 - 19
Hornblende	20	17 - 38
Basaltic hornblende	10	2 - 7
Tourmaline	2 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	0 - 2
Zircon	5	4 - 17
Garnet	6	2 - 17
Staurolite		0 - 3
Kyanite	3	0 - 1
Others	4 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	0 - 7

Potential Markets for Nonfuel Minerals

Many of the nonfuel mineral prospects located on the Texas continental shelf are located too far from potential markets to economically compete with abundant local, onshore deposits. However, two nonfuel resources that could be economically competitive in the future are sand for beach nourishment and sand and gravel for use in the concrete and construction industry.

Beach Nourishment (Sand)

Demand

Beach nourishment, the artificial restoration of a beach by adding sediment to offset beach erosion, can be attempted where substantial human investments in recreation, residence, or industry would be damaged by continued erosion. Long-term erosional trends of Texas beaches and heavy beach use near population centers makes beach nourishment an attractive alternative to other methods of shoreline stabilization. Beach nourishment has been considered for Galveston Island to re-create a beach that once existed

seaward of the Galveston seawall, to offset high rates of erosion (averaging up to 3 m/yr since the 1850's; Morton 1974) on beaches west of the seawall, to replace the estimated 1 million cubic yards of sand eroded from the western part of the island during a recent hurricane (Morton and Paine 1985), and to replace contaminated beach sand removed from the island after oil from the wrecked tanker *Alvenus* washed ashore in 1984.

Recent extensive development along south Padre Island has placed hotels, residences, and businesses near a beach that is eroding rapidly. Since 1867, average annual rates of erosion at the southern tip of Padre Island have been as high as 5 m/yr (Morton and Pieper 1975); recent rates as high as 6 m/yr (Paine and Morton 1988) indicate that erosion is likely to continue. As the shoreline retreats, endangered structures will either be destroyed, moved (if possible), or will be protected by engineered structures such as seawalls, groin fields, and breakwaters or by beach nourishment. Because the principal industry in this area is tourism, beach nourishment will likely be the chosen alternative.

Sources and Cost

Size of the material to be added to an eroding beach is of critical importance. If the material is too fine, it will erode rapidly; if too coarse, the aesthetics of the beach will not be preserved. One of the most promising sources of sand for nourishment of Texas' gulf beaches are shore-parallel sand bodies formed from similar materials and in a similar manner to today's Texas beaches and nearshore sands; thus they are closer to ideal size parameters than are onshore fluvial or deltaic sands. In addition, sands dredged offshore can be transported to the beach over water rather than hauled over land by trucks.

There are potential sources of beach-compatible sands located offshore from both south Padre Island and Galveston Island. Sand is particularly abundant off south Padre Island (figs. 6 and 9), where the postglacial sea-level rise has caused reworking of the sand-rich Rio Grande delta and produced a transgressive sand sheet across much of the

south Texas shelf. Sand is not as abundant offshore from Galveston, yet potential sources such as Heald Bank (65 km distant) and Sabine Bank (95 km distant) do exist.

Despite the attractiveness of some offshore sands for beach nourishment, recent studies have shown that offshore sources are more expensive to exploit than are nearby onshore sources. At Galveston Island, offshore sources of sand much nearer Galveston Island than Heald Bank were considered for a beach nourishment project, but were rejected in favor of compatible sand that could be piped or trucked from a site at the eastern end of the island (U.S. Army Corps of Engineers 1983). Costs of sand obtained in this manner were budgeted at \$6.75 per cubic yard; sand obtained by hopper dredge from nearby offshore sources was more than three times as expensive (\$21 per cubic yard). It is clear that as long as nearby compatible sands are available, use of offshore sand will not be economically feasible for beach nourishment.

Construction and Industry (Sand and Gravel)

Demand

There are diverse industrial and construction uses for sand and gravel. Industrial sand is used as an abrasive, as a refractory material in metal casting, as a propping agent in hydraulic fracturing of hydrocarbon reservoirs, and in glassmaking. Sand and gravel are also used by the construction industry in the making of concrete, as road base, and as fill; together these uses make the sand and gravel industry the second largest nonfuel mineral industry in the United States (Davis and Tepordei 1985).

By far the largest sand and gravel market on the coast of Texas is the Houston Metropolitan Area (HMA). During 1985, the HMA consumed an estimated 19.3 to 25.0 million tons of aggregate (Bureau of Mines 1987a). Other Texas population centers that consume smaller but substantial quantities of aggregate are Corpus Christi on the central coast and Brownsville on the southern coast. As abundant local supplies of sand and

gravel are exhausted, each of these areas may look to nearby offshore sources of sand and gravel.

Sources and Cost

Houston, Corpus Christi, and Brownsville are all located within major stream basins (Trinity/San Jacinto, Nueces, and Rio Grande). Fluvial sand and gravel similar to deposits found along these streams on land can also be expected to occur offshore along the downstream continuations of these streams. Many of these drowned streamcourses have been located by seismic surveys (see section on streamcourses and valley fill) and the presence of sand and gravel has been verified in some areas by coring.

Cost of sand at onshore pits and quarries is relatively low, ranging from \$1 to \$3 per ton in Houston (Bureau of Mines 1987a) to about \$4 per ton in the Corpus Christi area. Gravel is more expensive, ranging from \$4 to \$8 per ton in Houston to about \$8.50 per ton in Corpus Christi. The relatively low cost of sand and gravel at the quarry is offset by high transportation costs, making local sources much cheaper than distant sources. As nearby sources are depleted, delivered costs of sand and gravel to each of the metropolitan areas will rise and may increase interest in offshore sand and gravel deposits.

Despite the probable abundance of near surface sand and gravel on the Texas continental shelf, these deposits must be competitive with equally abundant sand and gravel on land. In a recent study of the potential for offshore sand and gravel production in the Houston area, it was estimated that despite the large consumption rate, more than 40 years of on-land supply remained (Bureau of Mines 1987a). Similar abundances in areas of lower demand, such as Corpus Christi and Brownsville, will last even longer.

Conclusions

There are abundant sand, gravel, and heavy mineral deposits on the Texas continental shelf. Significant sand accumulations at or near the seafloor occur as shore-

parallel sands and transgressive sheet sands that were deposited during the post-Wisconsinan rise in sea level and in shelf-edge deltas built during the late Wisconsinan sea-level lowstand. Fluvial sand and gravel occur along Wisconsinan streamcourses across the continental shelf; however, these basal valley-fill deposits may be covered by tens of meters of overburden. Surface accumulations of heavy minerals occur on the south Texas continental shelf offshore from the Rio Grande.

Potential markets for sand and gravel mined offshore exist in Texas. Sand such as that contained in drowned shoreline and nearshore deposits have the greatest near-term economic potential because they can be used for beach nourishment projects which would not require expensive overland transport. Industrial and construction sand and gravel, though abundant offshore, are also abundant onshore. With onshore supplies adequate for 40 years or more, near-term exploitation of offshore sand and gravel for industry and construction is not likely.

Recommendations

It is not anticipated that nonfuel minerals on the Texas continental shelf (principally sand and gravel) will become generally economic in the near future because the onshore supply is adequate for many years. As long as this remains true, demand for offshore deposits will be low. However, specific local accumulations, such as sand particularly suitable for nearby beach nourishment, could become economic at any time. Uneven distribution of sediment samples, cores, and high-resolution seismic coverage makes a comprehensive inventory of potentially economic deposits impossible, but has led to the discovery of some deposits. The following recommendations reflect the combination of low and sporadic demand, sparse data, and marginal economics for nonfuel minerals on the Texas continental shelf.

Recommendation 1. Leasing the Texas continental shelf for nonfuel mineral extraction should be done in a manner that will accommodate anticipated sporadic, single-user demand for specific offshore deposits rather than multi-user competition for widely-distributed resources.

Recommendation 2. Fill existing data gaps with a regional high-resolution seismic survey of the Brazos, Galveston, and western High Island areas. Seismic coverage is adequate for the remainder of the continental shelf. The recommended seismic survey will reveal potentially significant accumulations of sand and gravel along Wisconsinan courses of the Brazos, Colorado, and Trinity rivers.

Recommendation 3. If the Minerals Management Service anticipates needing to demonstrate the economic potential of offshore deposits, then characterization studies are recommended for three sites: Heald Bank sands, sand and gravel offshore from Mustang Island, and sand and heavy mineral concentrations off the Rio Grande. These studies should be tailored for each site, but would include surface samples, cores, and high-resolution seismic surveys. Of these three sites, Heald Bank has the highest potential for use. Heald Bank sands are attractive for beach nourishment because (1) there is a nearby market at Galveston Island, (2) size requirements for beach nourishment are strict, suitable on-land deposits are limited, and Heald Bank is composed of sediment similar to that on Galveston Island, and (3) offshore sand may have a transportation advantage over truck-hauled sand from distant on-land borrow sites.

References and Bibliography

- Aten, L. E., 1983. Indians of the upper Texas coast. New York, N.Y.: Academic Press. 370 pp.
- Ballard, R. C., and Uchupi, E., 1970. Morphology and Quaternary history of the continental shelf of the Gulf coast of the United States. Bulletin of Marine Science 20:547-559.
- Bandy, O. L., Frericks, W. E., and Vincent, E., 1973. Pleistocene-Holocene boundary and Wisconsinan substages, Gulf of Mexico. Geological Society of America Memoir 136: 277-316.
- Behrens, E. W., 1963. Buried Pleistocene river valleys in Aransas and Baffin bays,

 Texas. Port Aransas, Tx.: Publications of the Institute of Marine Science 9:7-18.
- Berryhill, H. L., Jr., 1976. Environmental studies, south Texas Outer Continental Shelf, 1976—Geology. Report PB 277-337/AS. Springfield, Va.: National Technical Information Service. 625 pp.
- Berryhill, H. L., Jr., 1977a. Shallow subsurface sediments. Pp. 167-229 in Berryhill, H. L., Jr., editor, Environmental studies, south Texas outer continental shelf (1976): Geology: Report PB 277-337. Springfield, Va.: National Technical Information Service.

- Berryhill, H. L., Jr., 1977b. Environmental studies, south Texas Outer Continental Shelf, 1975—An atlas and integrated summary. Report prepared for the U.S. Bureau of Land Management: U.S. Geological Survey. 303 pp.
- Berryhill, H. L., Jr., 1978a. Environmental studies, south Texas Outer Continental Shelf, 1977—Geology. Report PB 289-144/AS. Springfield, Va.: National Technical Information Service. 306 pp.
- Berryhill, H. L., Jr., 1978b. South Texas continental shelf and continental slope: Late Pleistocene/Holocene evolution and sea-floor stability. Open-File Report 78-514.

 U.S. Geological Survey. 91 pp.
- Berryhill, H. L., Jr., 1980. Map showing paleogeography of the continental shelf during the low stand of sea level, Wisconsinan glacial epoch, Port Isabel 1° x 2° quadrangle, Texas. Map I-1254-E. U.S. Geological Survey. Scale 1:250,000.
- Berryhill, H. L. Jr., 1981a. Map showing paleogeography of the continental shelf during the low stand of sea level, Wisconsinan glaciation, Corpus Christi 1° x 2° quadrangle. Map I-1287-E. U.S. Geological Survey. Scale 1:250,000.
- Berryhill, H. L. Jr., 1981b. Map showing paleogeography of the continental shelf during the low stand of sea level, Wisconsinan glacial epoch, Beeville 1° x 2° quadrangle. Map I-1288-E. U.S. Geological Survey. Scale 1:250,000.
- Berryhill, H. L., Jr., 1984a. Location of seismic profiles, Port Arthur sheet, Texas-Louisiana. Sheet 1, OCS Map Series MMS 84-0003. New Orleans, La.: Minerals Management Service. Scale 1:250,000.

- Berryhill, H. L., Jr., 1984b. Location of seismic profiles, Bouma Bank sheet. Sheet 1, OCS Map Series MMS 84-0028. New Orleans, La.: Minerals Management Service. Scale 1:250,000.
- Berryhill, H. L., Jr., 1987a. Use of high-resolution reflection profiling techniques for identifying major late Quaternary continental shelf/slope facies. Pp. 1-10 in Berryhill, H. L., Jr., editor, Late Quaternary facies and structure, northern Gulf of Mexico: interpretations from seismic data. Studies in Geology #23. Tulsa, Ok.: American Association of Petroleum Geologists.
- Berryhill, H. L., Jr., 1987b. The continental shelf off south Texas. Pp. 11-79 in

 Berryhill, H. L., Jr., editor, Late Quaternary facies and structure, northern Gulf of

 Mexico: interpretations from seismic data. Studies in Geology #23. Tulsa, Ok.:

 American Association of Petroleum Geologists.
- Berryhill, H. L., Jr., Owen, D. E., and Suter, J. R., 1984. Shallow geology of the continental shelf and upper continental slope, Port Arthur sheet, Texas-Louisiana.

 OCS Map Series MMS 84-0003. New Orleans, La.: Minerals Management Service. Scale 1:250,000.
- Berryhill, H. L., Jr., Owen, D. E., Trippet, A. R., and Mihalyi, D., 1984. Distribution of sediments of probable early and late Wisconsinan age, Bouma Bank sheet.

 Sheet 2, OCS Map Series MMS 84-0028. New Orleans, La.: Minerals

 Management Service. Scale 1:250,000.

- Berryhill, H. L, Jr., Shideler, G. L., Holmes, C. W., Hill, G. W., Barnes, S. S., and Martin, R. G., Jr., 1976. Environmental studies, south Texas Outer Continental Shelf, 1975—Geology. Publication PB-251341. Springfield, Va.: National Technical Information Service. 393 pp.
- Berryhill, H. L., Jr., and Suter, J. R., 1987. Deltas. Pp. 131-190 in Berryhill, H. L., Jr., editor, Late Quaternary facies and structure, northern Gulf of Mexico: interpretations from seismic data. Studies in Geology #23. Tulsa, Ok.: American Association of Petroleum Geologists.
- Berryhill, H. L., Jr., and Trippet, A. R., 1980. Map showing post-Wisconsin sedimentation patterns and faulting in the Port Isabel 1° x 2° quadrangle, Texas. Map I-1254-D. U.S. Geological Survey. Scale 1::250,000.
- Berryhill, H. L., Jr., and Trippet, A. R., 1981a. Map showing post-Wisconsin sedimentation patterns and faulting in the Corpus Christi 1° x 2° quadrangle, Texas.

 Map I-1287-D. U.S. Geological Survey. Scale 1:250,000.
- Berryhill, H. L., Jr., and Trippet, A. R., 1981b. Map showing post-Wisconsin sedimentation patterns and faulting in the Beeville 1° x 2° quadrangle, Texas. Map I-1288-D. U.S. Geological Survey. Scale 1:250,000.
- Bouma, Arnold H., 1972. Distribution of sediments and sedimentary structures in the Gulf of Mexico. Pp. 35-36 in Rezak, Richard, and Henry, V. J., editors, Contributions on the geological and geophysical oceanography of the Gulf of Mexico. Volume 3. Texas A&M University Oceanography Studies. Houston: Gulf Publishing Company.

- Brotherton, Mark A., 1982. Geology of aggregate resource in the Brazos and Colorado floodplains on the coastal plains of Texas. M.S. Thesis. College Station, Tx.:

 Texas A&M University.
- Bullard, F. M., 1942. Source of beach and river sands on gulf coast of Texas.

 Geological Society of America Bulletin 53,:1021-1044.
- Bureau of Mines, 1979. Minerals in the economy of Texas. State Mineral Profile. Washington, D.C.: U.S. Department of the Interior. 21 pp.
- Bureau of Mines, 1987a. An economic reconnaissance of selected sand and gravel deposits in the U.S. Exclusive Economic Zone. Open File Report OFR 3-87. Washington, D.C.: Bureau of Mines. 113 pp.
- Bureau of Mines, 1987b. An economic reconnaissance of selected heavy mineral placer deposits in the U.S. Exclusive Economic Zone. Open File Report OFR 4-87.

 Washington, D.C.: Bureau of Mines. 112 pp.
- Cogen, W. M., 1940. Heavy mineral zones of Louisiana and Texas Gulf coast sediments. American Association of Petroleum Geologists Bulletin 24:2069-2101.
- Coleman, James M., Prior, David B., and Roberts, Harry H., 1986. Geologic development and characteristics of the continental margins, Gulf of Mexico. Gulf Coast Association of Geological Societies Transactions 36:61-64.

- Crisp, J. A., 1985. Sources and distribution of coarse silts on the South Texas

 Continental Shelf. M.S. Thesis. College Station, Tx.: Texas A&M University.

 81 pp.
- Curray, J. R., 1960. Sediments and history of Holocene transgression, Continental Shelf, northwestern Gulf of Mexico. Pp. 221-226 in Shepard, F. P., Phleger, F. B., and van Andel, T. H., editors, Recent sediments, northwest Gulf of Mexico. American Petroleum Institute Project 51. Tulsa, Ok.: American Association of Petroleum Geologists.
- Curray, J. R., 1965. Late Quaternary history, continental shelves of the United States.

 Pp. 723-735 in Wright, H. E., Jr., and Frey, D. G., editors, The Quaternary of the United States. Princeton, N.J.: Princeton University Press.
- Davies, D. K., and Ethridge, F., 1975. Sandstone composition and depositional environment. American Association of Petroleum Geologist Bulletin 59:239-264.
- Davis, L. L., and Tepordei, V. V., 1985. Sand and gravel. Preprint from Bureau of Mines Bulletin 675. Washington, D.C.: Bureau of Mines. 15 pp.
- Devine, S. B., Ferrell, R. E., Jr., and Billings, G. K.,1973. Mineral distribution patterns, deep Gulf of Mexico. American Association of Petroleum Geologists Bulletin 57:28-41.
- Ewing, M., Ericson, D. B., and Heezen, B. C., 1958. Sediments and topography of the Gulf of Mexico. Pp. 995-1053 in Weeks, E., editor, Habitat of Oil. Tulsa, Ok.:

 American Association of Petroleum Geologists.

- Flores, R. M., and Shideler, G. L., 1978. Factors controlling heavy-mineral variations on the South Texas outer continental shelf, Gulf of Mexico. Journal of Sedimentary Petrology 48:269-280.
- Frazier, D. E., 1974. Depositional episodes: their relationship to the Quaternary stratigraphic framework in the northwestern portion of the Gulf basin. Geological Circular 74-1. The University of Texas at Austin, Bureau of Economic Geology. 28 pp.
- Grady, J. R., 1970. Chart of distribution of sediment types—northern Gulf of Mexico.

 National Oceanic and Atmospheric Administration, National Marine Fisheries

 Service, Gulf Coastal Fisheries Center.
- Halper, F. B., 1984. The effect of storms on sediment resuspension and transport on the outer continental shelf of the northwest Gulf of Mexico. Ph. D. dissertation.College Station, Tx.: Texas A&M University. 132 pp.
- Hawkins, J. W., 1983. Sources of relict sand on northeast Texas continental shelf:

 M.S. thesis. College Station, Tx.: Department of Oceanography, Texas A&M

 University. 63 pp.
- Holmes, C. W., 1971. Zirconium on the continental shelf—possible indicator of ancient shoreline deposition. Professional Paper 750-C. U.S. Geological Survey. Pp. C7-C12.

- Holmes, C. W., 1973. Distribution of selected elements in surficial marine sediments of the northern Gulf of Mexico continental shelf and slope. Professional Paper 814.U. S. Geological Survey. 7 pp.
- Holmes, C. W., 1976. Distribution, regional variation, and geochemical coherence of selected elements in the sediments of the central Gulf of Mexico. Professional Paper 928. U.S. Geological Survey. 24pp.
- Kahn, A. S., Latta, L. A., Oakes, R. L., Pert, D. N., Sweet, W. E., Smith, N., and Wade, E. J., 1975a. Geological and operational summary, Continental Offshore Stratigraphic Test (COST) No. 1, South Padre Island East Addition, offshore South Texas. Open File Report No. 75-174. U.S. Geological Survey. 32 pp.
- Kahn, A. S., Latta, L. A., Oakes, R. L., Pert, D. N., Sweet, W. E., Smith, N., and
 Wade, E. J., 1975b. Geological and operational summary, Continental Offshore
 Stratigraphic Test (COST) No. 2, Mustang Island, offshore South Texas. Open
 File Report No. 75-259. U.S. Geological Survey. 32 pp.
- Kunzendorf, H., 1986. Marine mineral exploration. Amsterdam, The Netherlands: Elsevier. 300 pp.
- Lewis, D. L., 1984. Pleistocene seismic stratigraphy of the Galveston South Addition, offshore Texas. M.S. thesis. Houston, Tx.: Rice University. 152 pp.
- Lindquist, P., 1978. Geology of South Texas shelf banks. M.S. thesis. College Station, Tx.: Texas A&M University. 138 pp.

- Mazzullo, J., and Crisp, J., 1985. Sources and dispersal of coarse silt on the South Texas continental shelf. Marine Geology 69:131-148.
- Mazzullo, J., and Withers, K. D., 1984. Sources, distribution, and mixing of Late Pleistocene and Holocene sands on the South Texas Continental Shelf. Journal of Sedimentary Petrology 54:1319-1334.
- McGowen, J. H., and Morton, R. A., 1979. Sediment distribution, bathymetry, faults, and salt diapirs, submerged lands of Texas. Special Report. The University of Texas at Austin, Bureau of Economic Geology. 31 pp.
- McKelvey, V. E., 1986. Subsea mineral resources. Bulletin 1689-A. U.S. Geological Survey. Washington, D. C.: U.S. Government Printing Office. 106 pp.
- Minerals Management Service, 1984. Resource evaluation program report: geological and geophysical data acquisition in the outer continental shelf. OCS Report MMS-84-0035. Reston, Va.: U.S. Department of the Interior. 30 pp.
- Moore, W. S., 1982. Late Pleistocene sea-level history. Pp. 481-506 in Ivanovitch, M., and Harmon, R. S., editors, Uranium series disequilibrium: applications to environmental problems. Oxford, England: Clarendon Press.
- Morton, R. A., 1974. Shoreline changes on Galveston Island (Bolivar Roads to San Luis Pass), an analysis of historical shoreline changes of the Texas Gulf shoreline.

 Geological Circular 74-2. The University of Texas at Austin, Bureau of Economic Geology. 34 pp.

- Morton, R. A., Jirik, L. A., and Foote, R. Q., 1985a. Facies analysis of hydrocarbon bearing sediments, offshore Texas: Geological Circular 85-2. The University of Texas at Austin, Bureau of Economic Geology. 31 pp.
- Morton, R. A., Jirik, L. A., and Foote, R. Q., 1985b. Structural cross sections,

 Miocene Series, Texas continental shelf. Cross Sections. The University of Texas

 Bureau of Economic Geology. 8 pp.
- Morton, R. A., and Paine, J. G., 1985. Beach and vegetation-line changes at Galveston Island, Texas: erosion, deposition, and recovery from Hurricane Alicia.

 Geological Circular 85-5. The University of Texas at Austin, Bureau of Economic Geology. 39 pp.
- Morton, R. A., and Pieper, M. J., 1975. Shoreline changes on Brazos Island and south Padre Island (Mansfield Channel to mouth of the Rio Grande): an analysis of historical changes of the Texas Gulf shoreline. Geological Circular 75-2. The University of Texas at Austin, Bureau of Economic Geology. 39 pp.
- Morton, R. A., and Price, W. A., 1984. Influence of sea level on erosional, depositional, and submergence patterns of the Texas Coast. Abstract. Tulsa, Ok.:

 Society of Economic Paleontologists and Mineralogists 1984 midyear meeting. Pp. 67.
- Morton, R. A., and Price, W. A., 1987. Late Quaternary sea-level fluctuations and sedimentary phases of the Texas coastal plain and shelf. Pp. 181-198 in Nummedal, D., Pilkey, O. H., and Howard, J. D., editors, Sea level fluctuations

- and coastal evolution. Special Publication No. 41. Tulsa, Ok.: Society of Economic Paleontologists and Mineralogists.
- Morton, R. A., and Winker, C. D., 1979. Distribution and significance of coarse biogenic and clastic deposits on the Texas inner shelf. Gulf Coast Association of Geological Societies Transactions 29:306-320.
- Nelson, H. F., and Bray, E. E., 1970. Stratigraphy and history of the Holocene sediments in the Sabine-High Island area, Gulf of Mexico. Pp. 48-77 in Morgan,
 J. P., and Shaver, R. H., editors, Deltaic sedimentation, modern and ancient.
 Special Publication 15. Tulsa, Ok: Society of Economic Paleontologists and Mineralogists.
- Nienaber, J. H., 1963. Shallow marine sediments offshore from the Brazos River, Texas. Port Aransas, Tx.: The University of Texas, Marine Science Institute Publication 9:311-372.
- Ohl, J. P., and McBride, M. W., 1987. The mineral industry of Texas in 1985. Mineral Resource Circular No. 79. The University of Texas at Austin, Bureau of Economic Geology. 20 p.
- Paine, J. G., and Morton, R. A., 1988. Shoreline and vegetation-line movement, Texas gulf coast, 1974 to 1982. Geological Circular 88-5. The University of Texas at Austin, Bureau of Economic Geology (in press).

- Parker, R. H., and Curray, J. R., 1956. Fauna and bathymetry of banks on continental shelf, northwest Gulf of Mexico. American Association of Petroleum Geologists

 Bulletin 40:2428-2439.
- Pearson, C. E., Kelley, D. B., Weinstein, R. A., and Gagliano, S. M., 1986.

 Archeological investigations on the outer continental shelf: a study within the

 Sabine River valley, offshore Louisiana and Texas. Report prepared for Minerals

 Management Service. Baton Rouge, La.: Coastal Environments, Inc., 314 pp.
- Poag, C. W., 1972. Shelf-edge submarine banks in the Gulf of Mexico: paleoecology and biostratigraphy. Gulf Coast Association of Geological Society Transactions 22:267-287.
- Poag, C. W., 1973. Late Quaternary sea levels in the Gulf of Mexico. Gulf Coast Association of Geological Societies Transactions 23:394-400.
- Poag, C. W., and Valentine, P. C., 1976. Biostratigraphy and ecostratigraphy of the Pleistocene basin Texas-Louisiana continental shelf. Gulf Coast Association of Geological Societies Transactions 26:185-256.
- Price, W. A., 1954. Dynamic environments: reconnaissance mapping, geologic and geomorphic, of continental shelf of Gulf of Mexico. Gulf Coast Association of Geological Societies Transactions 4,:75-107.
- Pyle, C. A., 1977. Late Quaternary geologic history of the South Texas Continental Shelf. M.S. thesis. College Station, Tx.: Department of Oceanography, Texas A&M University. 72 pp.

- Rehkemper, L. J., 1969. Sedimentology of Holocene estuarine deposits, Galveston Bay. Pp. 12-52 in Lankford, R. R., and Rogers, J. J. W., compilers, Holocene geology of the Galveston Bay area. Houston, Tx: Houston Geological Society.
- Rezak, R., 1985. Local carbonate production on a terrigenous shelf. Gulf Coast Association of Geological Societies Transactions 35:477-483.
- Rezak, R., Bright, T. J., and McGrail, D. W., 1985. Reefs and banks of the northwestern Gulf of Mexico: their geological, biological, and physical dynamics. New York, NY: John Wiley & Sons. 259 pp.
- Rezak, R., and Henry, V. J., editors, 1971. Contributions on the Geological and Geophysical Oceanography of the Gulf of Mexico: Volume 3. Texas A&M University Oceanographic Studies. Houston, Tx.: Gulf Publishing Co. 303 pp.
- Rittenhouse, G., 1943. The transportation and deposition of heavy minerals. Geological Society of America Bulletin, 54:1725-1780.
- Roemer, L. B., and Bryant, W. R., 1977. Structure and stratigraphy of late Quaternary deposits on the outer continental shelf. Technical Report. College Station, Tx:

 Department of Oceanography, Texas A&M University. 169 pp.
- Shepard, F. P., 1973. Submarine Geology. Third edition. New York, NY.: Harper and Row. 517 pp.

- Shideler, G. L., 1976. Textural distribution of sea-floor sediments, south Texas outer continental shelf. Journal of Research, U.S. Geological Survey 4:703-713.
- Shideler, G. L., 1977. Late Holocene sedimentary provinces, South Texas Outer

 Continental Shelf. American Association of Petroleum Geologists Bulletin 61:708722.
- Sidner, B. R., Gartner, S., and Bryant, W. R., 1978. Late Pleistocene geologic history of Texas outer continental shelf and upper continental slope. Pp. 243-266 in Bouma, A. H., Moore, G. T., and Coleman, J. M., editors, Framework, facies, and oil-trapping characteristics of the upper continental margin. Studies in Geology 7. Tulsa, Ok.: American Association of Petroleum Geologists.
- Slatt, R. M., 1984. Continental shelf topography: Key to understanding distribution of shelf sand-ridge deposits from Cretaceous Western Interior seaway. American Association of Petroleum Geologist Bulletin 68:1107-1120.
- Stetson, H. C., 1953. The sediments of the western Gulf of Mexico: Part I— the continental terrace of the western Gulf of Mexico; its surface sediments, origin, and development. Massachusetts Institute of Technology and Woods Hole

 Oceanographic Institute papers in physical oceanography and meteorology 12(4):3-45.
- Suter, J. R., 1987. Ancient fluvial systems and Holocene deposits, southwestern

 Louisiana continental shelf. Pp. 131-190 in Berryhill, H. L., Jr., editor, Late

 Quaternary facies and structure, northern Gulf of Mexico: interpretations from

- seismic data. Studies in Geology #23. Tulsa, Ok.: American Association of Petroleum Geologists.
- Suter, J. R., and Berryhill, H. L., Jr., 1985. Late Quaternary shelf-margin deltas, northwest Gulf of Mexico. American Association of Petroleum Geologist Bulletin 69:77-91.
- Sweet, W. E., Reed, J. C., Leyendecker, C. L., and Khan, A. S., 1986. Correlation of Cenozoic sediments, Gulf of Mexico, Outer Continental Shelf: Galveston Area, offshore Texas, to Vermilion Area, offshore Louisiana. Abstract. American Association of Petroleum Geologists Bulletin 70:654.
- Tatum, T. E., 1979. Shallow geologic features of the upper continental slope, northwestern Gulf of Mexico. Technical Report 79-2-T. College Station, Tx.: Department of Oceanography, Texas A&M University. 60 pp.
- Thayer, P. A., La Rocque, A., and Tunnell, J. W., Jr., 1974. Relict lacustrine sediments on the inner continental shelf, southeast Texas. Gulf Coast Association of Geological Societies Transactions 24:337-347.
- U.S. Army Corps of Engineers, 1983. Galveston County shore erosion study: feasibility report and environmental impact statement. Draft, 3 volumes.Galveston, Tx.: Galveston District, U.S. Army Corps of Engineers. Not consecutively paginated.

- U.S. Congress, Office of Technology Assessment, 1987. Marine minerals: exploring our new ocean frontier. OTA-O-342. Washington, D.C.: U.S. Government Printing Office. 347 pp.
- U.S. Geological Survey, 1984. A national program for the assessment and development of the mineral resources of the United States Exclusive Economic Zone. Circular 929. Alexandria, Va.: U.S. Geological Survey. 308 pp.
- van Andel, T. H., 1960. Sources and dispersion of Holocene sediments, northern Gulf of Mexico. Pp. 34-55 in Shepard, F. P., Phleger, F. B., and van Andel, T. H., editors, Recent sediments, northwest Gulf of Mexico. American Petroleum Institute Project 51. Tulsa, Ok.: American Association of Petroleum Geologists.
- van Andel, T. H., and Poole, D. M., 1960. Sources of recent sediments in the northern Gulf of Mexico. Journal of Sedimentary Petrology 30:91-122.
- White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., Littleton, T. G., McGowen, J. H., Nance, H. S., and Schmedes, K. E., 1983. Submerged lands of Texas, Corpus Christi area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands. Special Publication. The University of Texas at Austin, Bureau of Economic Geology. 154 pp.
- White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., Littleton, T. G., McGowen, J. H., Nance, H. S., and Schmedes, K. E., 1985a. Submerged lands of Texas, Galveston-Houston area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands. Special Publication. The University of Texas at Austin, Bureau of Economic Geology. 145 pp.

- White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., Littleton, T. G., McGowen, J. H., and Nance, H. S., 1985b. Submerged lands of Texas, Bay City-Freeport area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands. Report prepared for the Minerals Management Service. The University of Texas at Austin, Bureau of Economic Geology. 214 pp.
- White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., Littleton, T. G., McGowen, J. H., and Nance, H. S., 1985c. Submerged lands of Texas, Port Lavaca area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands. Report prepared for the Minerals Management Service. The University of Texas at Austin, Bureau of Economic Geology. 309 pp.
- White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., Littleton, T. G., McGowen, J. H., Nance, H. S., and Schmedes, K. E., 1986a. Submerged lands of Texas, Brownsville-Harlingen area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands. Special Publication. The University of Texas at Austin, Bureau of Economic Geology. 138 pp.
- White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., Littleton, T. G., McGowen, J. H., and Nance, H. S., 1986b. Submerged lands of Texas, Kingsville area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands. Report prepared for the Minerals Management Service. The University of Texas at Austin, Bureau of Economic Geology. 271 pp.
- White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., Littleton, T. G., McGowen, J. H., and Nance, H. S., 1987. Submerged lands of Texas,

- Beaumont-Port Arthur area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands. Special Publication. The University of Texas at Austin, Bureau of Economic Geology. 138 pp.
- Williams, S. J., Prins, D. A., and Meisburger, E. P., 1979. Sediment distribution, sand resources, and geologic character of the Inner Continental Shelf off Galveston, County, Tx. Miscellaneous Report No. 79-4. Fort Belvoir, Va.: U.S. Army Corps of Engineers, Coastal Engineering Research Center. 159 pp.
- Winker, C. D., 1979. Late Pleistocene fluvial-deltaic deposition, Texas coastal plain and shelf: M.S. thesis. Austin, Tx.: The University of Texas at Austin. 187 pp.
- Winker, C. D., 1982. Cenozoic shelf margins, northwestern Gulf of Mexico. Gulf Coast Association of Geological Societies Transactions 32:417-448.
- Withers, K. D., 1984. Grain shape variance in Late Pleistocene and Holocene fluvial and shelf sands in the Northwest Gulf, and the relationship to source and shelf paleogeography. M.S. thesis. College Station, Tx.: Texas A&M University. 50 pp.
- Woodbury, H. O., Murray, I. B., Jr., Pickford, P. J., and Akers, W. H., 1973.

 Pliocene and Pleistocene depocenters, outer continental shelf, Louisiana and Texas.

 American Association of Petroleum Geologists Bulletin 57:2428-2439.
- Wright, S. S., 1980. Seismic stratigraphy and depositional history of Holocene sediments on the central Texas gulf coast. M.S. thesis. Austin, Tx.: The University of Texas at Austin. 123 pp.

Appendix

Lon, X, W refers to either longitude, X-coordinate, or distance (in feet) from eastern edge of lease block. Lat, Y, S refers to either latitude, Y-coordinate, or distance (in feet) from northern edge of lease block.

					Water	Boring		
Lease Area	Block	Name	Lon. X. W	Lat. Y. S		(ft)	Source	Date
Brazos	337	Brazos 37	1902	14762	50.0	40.0	McClelland Engineers	1/1/75
Brazos	340	Brazos 60	13291	2764	43.0	304.0	McClelland Engineers	2/18/82
Brazos	340	Brazos 64	12140	7080	50.0	305.5	McClelland Engineers	5/5/82
Brazos	14 V 24 V 240 V 340 V 340 V 3	CB-1			52.0	304.0	National Soil Services	6/20/78
Brazos	341	Brazos 30	13208	7508	0.09	40.0	McClelland Engineers	1/1/73
Brazos	341	Brazos 38	13959	8555	29.0	40.0	McClelland Engineers	1/1/75
Brazos	364	Brazos 52	15327	6253	65.0	256.0	McClelland Engineers	10/26/80
Brazos	364	Brazos 53	8265	10019	71.0	120.0	McClelland Engineers	10/30/80
Brazos	365	Brazos 39	6171	12285	63.0	40.0	McClelland Engineers	1/1/75
	376	Brazos 82	3040	4080	56.0	200.0	McClelland Engineers	5/16/87
9 Brazos		Brazos 29	14906	2200	75.0	40.0	McClelland Engineers	1/1/73
Brazos		Brazos 40	11014	9906	84.0	40.0	McClelland Engineers	1/1/75
Brazos	397	Brazos 72	9193	5713	80.0	381.0	McClelland Engineers	2/10/84
Brazos		Brazos 81	12567	12713	78.0	181.0	McClelland Engineers	11/1/85
Brazos	A ^{rg}	Brazos 1	15540	006	47.0	109.5	Greer & McClelland	10/1/48
Brazos		Brazos 2	4905	10840	20.0	210.0	McClelland Engineers	8/1/66
Brazos		Brazos 50	831	15146	61.0	371.0	McClelland Engineers	1/1/80
Brazos	417 1	Brazos 41	6993	3436	95.0	40.0	McClelland Engineers	1/1/75
Brazos	430	Brazos 42	1103	2514	0.96	40.0	McClelland Engineers	1/1/75
Brazos	430	Brazos 44	4719	14882	0.86	150.0	McClelland Engineers	1/1/75
Brazos	437	CB-1			0.89	256.0	National Soil Services	5/23/80
Brazos	438	Brazos 45	11742	1301	26.0	277.0	McClelland Engineers	1/1/78
Brazos	438	Brazos 59	15336	479	26.0	250.0	McClelland Engineers	6/21/81
Brazos	438	Brazos 61	2020	528	59.0	302.0	McClelland Engineers	1/27/82
Brazos		Brazos 67	11325	972	27.0	308.0	McClelland Engineers	8/6/82
Brazos	440	Brazos 4	9840	1000	52.0	226.0	McClelland Engineers	1/1/66
Brazos	440	Brazos 78			59.0	316.0	McClelland Engineers	6/21/85
Brazos		Brazos 3	10840	5500	50.0	165.0	McClelland Engineers	1/1/66
Brazos		Brazos 58	5683	13833	58.0	305.5	McClelland Engineers	11/7/81
Brazos	446	Brazos 62	0430	14557	53.5	338.0	McClelland Engineers	//13/81

	Date	2/12/82	5/4/82	5/1/82	3/12/80	5/1/85	12/8/83	1/1/70	9/3/85	1/1/10	8/2/81	1/1/68	1/1/70	1/1/70	1/1/68	3/1/81	1/13/81	1/1/79	1/1/19	1/1/79	1/1/81	1/1/72	1/1/75	1/1/75	1/1/73	1/1/75	1/29/84	1/28/81	12/18/84	12/15/83	8/24/84	10/8/84	3/14/84		_		F1 1/11/11	-	1/1//4
	Source	McClelland Engineers	McClelland Engineers	McClelland Engineers	National Soil Services	McClelland Engineers	McClelland-Engineers	McClelland Engineers		McClelland Engineers		McClelland Engineers	National Soil Services	National Soil Services	McClelland Engineers																								
Boring	(ft)	300.0	79.0	301.0		255.0					-			T	-						_	_			_		- 1	. '		- '		_,						80.0	
Water		53.0	57.0	57.0	56.0	59.0	73.0	86.0	70.0	86.0	110.0	102.0	88.0	-91.0	97.0	121.0	121.0	129.0	125.5	123.0	141.0	46.0	16.0	23.0	44.5	$\frac{30.0}{20.0}$	38.0	142.0	161.0	165.0	164.0	160.0	166.0	157.0	165.0	183.0	178.0	1/6.0	100.0
	Lat. Y. S	14557	14549	14623		14159	738	14852	3982	8214	10983	13041	10516	10550	7289	14865	9401	9311	10639	1457	4613	3759	2250	2816	3940	3016	2240	7930	11000	7033	7473	4216	13963	8017	1998	1000		2/° 56 20.11"	1809
	Lon, X, W	6436	6712	6943		13352	4601	843	70	11076	15254	7759	126	140	8328	4131	8838	12752	487	8280	13332	1378	2309	2989	3975	3279	1014	7598	5500	9703	26/1	14523	13532	8585	6311	14840	Č	95° 59° 07.82"	C7011
	Name	Brazos 63	Brazos 65	Brazos 66	CB-1	Brazos 77	Brazos 70	Brazos 8	Brazos 79	Brazos 9	Brazos 57	Brazos 5	Brazos 10	Brazos 11	Brazos 6	Brazos 56	Brazos 54	Brazos 48	Brazos 46	Brazos 47	Brazos 51	Brazos 28	Brazos 43	Brazos 35	Brazos 31	Brazos 36	Brazos 69	Brazos 55	Brazos 75	Brazos 71	Brazos 73	Brazos 74	Brazos 76	Brazos 33		Brazos 12	CB-1	, , , , , , , , , , , , , , , , , , ,	Brazos 32
	Block	446	446	446	446	449	452	474	489	495	505	206	510	510	538	542	A-007	A-019	A-020	A-020	A-039	8386	s405	s412	s415	s438	s468	A-047	A-050	A-052	A-052	A-052	A-065	A-070	A-076	A-084	A-102	A-102	A-105
	Lease Area		Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos	Brazos South	Brazos South	Brazos South	Brazos South	Brazos South	Brazos South	Brazos South	Brazos South				

					Water Depth	Boring Lenoth		
Lease Area	Block	Name .	Lon, X, W	Lat. Y. S	(ft)	(ft)		Date
Brazos South Brazos South	A-105 A-105	CB-1			196.0	59.0 66.0	National Soil Services	1/22/12
Brazos South	A-105	CB-2			196.0	44.0	Soil	103/12
Brazos South	A-106	Brazos 49	13321	15037	197.0	375.0	McClelland Engineers	1/17/9
Brazos South		Brazos 34	12826	3484	202.0	401.0	McClelland Engineers	1/1/75
Brazos South		Brazos 68	5082	3718	206.0	303.0	McClelland Engineers	9/17/83
Brazos South	A-133	Brazos 80	8283	9563	203.0	400.0	McClelland Engineers	11/29/85
Galveston	100	Galveston 65	6820	3732	36.0	302.5	McClelland Engineers	12/16/79
Galveston	102	Galveston 62	9365	9269	33.0	357.0	McClelland Engineers	3/11/79
Galveston	102	Galveston 63	5840	4345	33.0	277.5	McClelland Engineers	3/30/79
Galveston	102	Galveston 64	∞ ;	4179	30.0	321.0	McClelland Engineers	4/15/79
Galveston	1 2	CB-1	94° 32′ 23.46″	29° 19' 37.39"	43.0	300.0	National Soil Services	7/20/77
Galveston	<u>4</u>	CB-1			48.0	300.0	National Soil Services	7/22/77
Galveston	4	Galveston 22	250	8700	20.0	200.0	McClelland Engineers	1/1/56
Galveston	146	CB-1			36.0	292.0	National Soil Services	<i>TV</i> 9/ <i>T</i>
Galveston	241	Galveston 70	4745	2082	0.09	304.0	McClelland Engineers	2/18/83
Galveston	247	Galveston 6	9200	14385	49.0	214.0	McClelland Engineers	1/1/55
Galveston	248	Galveston 1	9015	14135	51.0	150.0	Greer and McClelland	1/1/48
Galveston	249	Galveston 2	11700	975	46.0	120.0	Greer & McClelland	12/1/54
Galveston	249	Galveston 4	5725	1595	51.0	157.5	McClelland Engineers	1/1/55
Galveston	249	Galveston 5	8025	2670	49.0	117.0	McClelland Engineers	1/1/55
Galveston	249	Galveston 7	4140	755	49.0	320.0	McClelland Engineers	1/1/55
Galveston	253	Galveston 8	12140	10440	0.99	139.0	McClelland Engineers	1/1/56
Galveston	255	Galveston 43	11503	2618	64.0	223.0	McClelland Engineers	1/1/71
Galveston	727	Galveston //	13939	14346	57.0	300.0	McClelland Engineers	3/23/87
Calveston	8/7 8/7	Galveston 3	1650	8200	55.0	145.0	Greer and McClelland	1/1/48
Calveston	000	Calveston 31	6612	0300	0.70	529.0	McClelland Engineers	1/1/64
Galveston	703 203	Galveston 28	3340 10050	9930	0.60	271.0	McClelland Engineers	1/1/64
Galveston	20%		1320	3375	0.89	273.0	McClelland Engineers	1/1/64
Galveston	296	Galveston 34	4215	2155	0.89	243.0	McClelland Engineers	1/1/64
Galveston	300	Galveston 71	11116	3838	0.09	368.0	McClelland Engineers	4/0/84
Galveston	308	Galveston 76			69.0	151.0	McClelland Engineers	11/14/85
Galveston	310	Galveston 49	2956	14701	61.0	271.0	McClelland Engineers	
Galveston	310	Galveston 50	4372	15626	64.0	26.0	McClelland Engineers	
Galveston	310		2874	2491	59.0	305.0	McClelland Engineers	1/29/82 🛒
Galveston	334		5851	824	64.0	26.0	McClelland Engineers	1/1/75 鬥
Galveston	334	Galveston 52	7304	1820	63.5	26.0	McClelland Engineers	1/1/75

]	DE	RA]	FΤ	٠,,	
	Date	1/1/75	6/22/84	6/23/84	5/19/79	11/14/79	1/1/73	8/20/84	1/1/73	1/1/75	1/1/75	1/1/75	1/1/75	1/1/75	1/1/73	1/1/73	1/1/73	10/14/85	91/9/9	8/22/80		1/1/59	10/9/80	11/29/82	7/11/78	7/16/78	1/1/75	1/1/68	1/1/75	2/19/79	7/29/83	10/4/85	1/1/68	11/15/81	6/13/84	7/12/84	1/1//1	18/C1/11	
	Source	McClelland Engineers	McClelland Engineers	McClelland Engineers	National Soil Services	National Soil Services	McClelland Engineers	National Soil Services	McClelland Engineers	National Soil Services, Inc	McClelland Engineers	National Soil Services	McClelland Engineers																										
Boring Length	(ft)	278.0	299.0	302.0	363.0	356.0	40.0	403.0	40.0	335.0	330.0	150.0	150.0	150.0	330.0	121.0	120.0	304.0	150.0	356.0	150.5	107.5	300.5	327.0	471.5	401.5	432.0	251.5	37.0	249.0	301.0	308.0	249.5	401.0	184.0	222.0	321.0	401.3 174.0) ,
Water Depth	(ft)	61.0	100.0	0.86	0.86	0.96	94.0	102.0	102.0	0.66	0.86	102.0	106.0	101.0	111.0	115.0	113.0	113.0	26.0	27.0	26.5	31.0	169.0	161.0	177.5	186.0	433.0	35.0	42.0	36.0	38.0	40.0	37.0	35.0	32.0	30.0	42.0	44.0	
	Lat. Y. S.	2876	13584	7550			6317	1939	12371	12454	12961	10491	4624	2481	1794	10126	326	9149	602,555.85	2197	598,067.57	4000	11156	7897	2009	10568	9264	14547	15373		12413	2026	3046	473	94° 22' 57.3"		5030	617,285	1314.13
	Lon, X, W	8830	11040	11140			14955	2001	4413	13199	12569	5346	14884	7156	10138	9138	2875		3,390,695.87		3,387,676.40	3000	6941	4906	6500	10768	2738	14515	1150		6522	2934 (E)	10833		29° 30' 0.62"		14820	3,540,634	
	Name	Galveston 53	Galveston 72	Galveston 73	CB-1	CB-1	Galveston 48	Galveston 74			Galveston 55		Galveston 56	Galveston 57	Galveston 44	Galveston 45		Galveston 75		Galveston 66	CB-1	Galveston 23	Galveston 67	Galveston 69	Galveston 61	Galveston 60	Galveston 59	High Island 32	High Island 83	CB-1			Sland (Island	Island	High Island 183	High Island 160		
	Block	334	386	386	391	393	418	424	429	429	429	429	460	460	4 4 4 4	464	464	465	s174	s175	s182	s226	A-126	A-127	A-131	A-157	A-248	002	800	010	022	022	024	026	027	027	020	051	F .
	Lease Area	Galveston	Galveston	Galveston	Galveston	Galveston	Galveston	Galveston	Galveston	Galveston	Galveston	Galveston	Galveston	Galveston	Galveston	Galveston	Galveston	Galveston	, Galveston	Galveston	Galveston	Galveston	Galveston South	High Island	High Island	High Island	High Island			High Island	High Island	High Island	High Island	riign Island High Island					

																																				DF	R.S	F	Γ	
		Date	4/15/81	2/18/84	1/1/69	3/25/85	1/1/76	1/1/16	1/1/74	4/29/87	1/1/64	1/1/66	7/12/80	277/81	1/29/19		1/1/56	9/22/77	1/1/16	1/1/16	1/1/16	1/1/76	1/1/75	1/1/60	1/1/66	8/26/82	1/1/16	1/1/60	8/25/85	12/21/82	2/21/86	2/26/86	3/9/87	1/1/56	7/30/56	1/1/56	7/27/56	7/28/56	7/28/56	7/28/56
			McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers		McClelland Engineers		McClelland Engineers																														
Boring	Length	(11)	278.5	414.0	329.0	405.0	303.0	348.0	276.0	350.0	369.0	270.0	439.0	394.0	355.0	191.0	134.0	284.5	29.0	206.0	49.0	28.0	451.0	250.0	250.0	81.0	317.0	245.0	332.0	0.66	299.4	77.0	305.0	200.0	0.09	191.0	80.0		80.0	80.0
Water	Depth		43.0	42.0	37.0	44.0	42.0	50.0	45.0	50.0	50.0	50.0	49.0	50.0	51.0	50.0	50.0	53.0	51.0	51.0	52.0	52.0	50.0	52.0	51.0	58.0	55.0	62.0	61.0	0.99	80.0	84.0	83.0	82.0	82.0	81.0	81.0	82.0	81.0	81.0
	D 22 1-1	Lat. Y. S	5272	1595	15840	2572		14340	8394	10256	12945	11340	5466	618	6318	550,720	7100	15175					7082	8838	7840	15230	6198		619	829	13164	7735	11830	13200	13200	10600	10600	12600	0098	8200
	'A A A'	Lon, A, W	9759	2223	2501	1501		1542	100	13304	14520	100	4001	8175	14178	3,485,360	099	333					400	5277	15640	4195	6553		13417	10143	5101	3122	4776	13200	13700	7000	009	2600	2600	4600
	N. S. see	' اده	Island	High Island 168	Island	High Island 173	High Island 97	High Island 104	High Island 52	High Island 187	High Island 28		High Island 149	High Island 152	High Island 137		sland	Island	Island	High Island 106	Island	High Island 108	High Island 99	High Island 26	High Island 30		High Island 112	High Island 27	High Island 176	High Island 166	High Island 184	High Island 181	High Island 186	High Island 8	High Island 9	High Island 21	Island	Island	High Island 24	High Island 25
	Disch	DIOCK	055	890	071	880	860	110	110	134	135	136	137	138	139	140	141	142	154	154	154	154	154	161	161	170	193	236	A-020	A-033	A-052	A-053	A-068	A-072	A-072	A-073	A-073	A-073	A-073	A-073
		Lease Area	High Island	High Island		High Island	High Island	High Island	High Island	High Island	High Island	High Island	High Island	High Island	High Island	High Island		High Island				High Island	High Island	High Island	High Island				High Island		High Island	High Island	High Island	High Island						

	1/1/56	7/26/56	7/26/56	8/9/56	2/25/86	1/1/56	8/6/56	95/9/8	95/9/8	95/9/8	8/8/56	8/5/56	877/56	8/9/26	1/1/55	4/10/81	1/1/56	1/1/75	1/1/78	1/1/75	1/1/69	1/1//1	8/2/84	11/23/83	1/1/75	1/1/75	1/1/75	8/8/82	8/7/82	1/1/75	1/1/75	S	1/1/75 및	1/1/75 💆	10/30/82	1/1/74	S//I/I
Solution N	McClelland Engineers	McClelland Engineers	McClelland Engineers			McClelland Engineers	National Soil Services	McClelland Engineers	McClelland Engineers		McClelland Engineers	McClenand Engineers	McClelland Engineers		McClelland Engineers																						
Boring Length	193.0	80.0	80.0	80.0	61.2	198.0	80.0	80.0	78.0	68.0	81.0	70.0	78.0	78.0	200.0	303.0	199.0	307.5	193.0	337.0	275.5	293.0	321.0	347.0	16.0	16.0	16.0	310.0	300.0	17.0	15.0	15.0	15.0	16.0	305.0	455.0	10.0
Water Depth	82.0	82.0	82.0	80.0	79.0	85.0	82.0	83.0	84.0	84.0	82.0	83.0	85.0	85.0	92.0	27.5	27.0	29.0	45.0	50.0	50.0	48.0	21.0	71.0	74.0	78.0	86.0	123.0	119.0	123.0	126.0	131.0	135.0	147.0	156.5	154.0	140.0
× 18.7	13200	13,200	13200	2400	2896	14600	9400	4100	9400	9400	4100	12400	9300	12000	2640	2150	750	2400		10745	7,00	2001	0/23 0/00/77	1000	-	4462	899	15106	986	9478	1242	3860	8134	14134	4857	7947	2411
Lon X. W	13200	12,200	14200	4300	6117	13200	13200	13200	7900	8200	2006	1300	2600	1300	2640	4630	200	1625		13841	1760	1500	1001 7117	191	9837	1110	2356	2036	4567	480	1740	3056	4254	5546	6205	7925	00/0
Name	High Island 5	High Island 6		High Island 20	High Island 180	High Island 10	High Island 12	High Island 13	High Island 14	High Island 15	High Island 18	High Island 11	High Island 16	High Island 19	-	High Island 154	High Island 4	High Island 74	CB-1	High Island 70	High Island 34	Light Island 170	High Island 177	High Island 171	High Island 86	Island	High Island 88	Island	Island	Island	Island	Island	Sland	High Island 93		High Island 5/	Dialic
Block	A-073	A-073	A-073	A-074	A-075	A-077	A-077	A-077	A-077	A-077	A-077	A-078	A-078	A-078	A-104	2087	s095	s140	914	119	671	120	A-178	A-193	A-218	A-228	A-228	A-244	A-244	A-248	A-255	A-255	A-255	A-255	th A-262	th A 264	
ease Area	High Island	High Island	High Island	High Island	Island	High Island	Island	Island	High Island	High Island	High Island	High Island	High Island			High Island		Island		Island	High Island East		Liigh Island East High Island Fast		High Island East	High Island East	Island	High Island East	Island		High Island East	High Island East	High Island East	High Island East	High Island East South A-262	High Island East South	Island Last Sou

																•																			DI	R.A	F	Γ	
		Date	1/1/75	10/7/81	3/15/84	1/1/75	5/25/18	5/24/78	10/13/84	11/23/77	170001	1/1/75	1/1/75	1/1/75	111177	12/26/79	1/1/76	1/1/76	1/1/73	1/1/74	9/26/78	1/1/73	1/1/74	1/1/75	12/23/73	1/1/73	1/1/75	1/1/75	1/1/74	1/1//4	1/1//4	9//6/1	1/1/16	1/1/74	1/1/75	1/1/77	1/1/17	1/1/74	11111111
			McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	National Soil Services	National Soil Services	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	National Soil Services	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	MCCICIIANU ENBINEEIS
Boring	Length	(ft)	16.0	353.0	353.0	323.0	314.0	315.0	340.5	423.0	388.0	455.0	424.0	436.0	318.0	353.0	429.5	374.5	382.0	498.5	360.0	380.0	335.5	334.0	72.0	408.0	408.0	420.0	430.5	450.3	381.0	407.0	339.0	392.0	392.4	334.5	150.0	350.0	2001
Water	Depth	(ij)	151.0	142.0	166.0	164.0	164.0	163.0	170.0	172.0	167.0	192.0	213.0	210.0	213.0	216.0	217.0	217.0	234.0	236.5	227.0	227.0	225.0	227.0	257.0	266.0	260.0	257.0	237.0	0.167	230.0	235.5	237.0	239.0	235.6	277.0	277.0	316.0	707.0
		Lat. Y. S	7257	3094	12832	8361	6092	7775	514 (N)	28° 22' 24.418"	28° 23' 42.525"	4000	8565	8739	11931	2097	6892	7113	10843	8093	10628	6084	14331	15537			12703	19871	C707	722	7386	3395	5412	5100	3007	6146	6427	9049 4402	774
		Lon, X, W	7730	15219	9322	34	9085	8912	1777	93° 53' 27.932"	· 93° 53' 57.445"	12709	9011	8838	2500	6505	5955	5747	14351	7852	11548	6029	10433	9592			1033	7000	13030	12030	111136	4812	9220	10178	3122	403 (East)	530 (East)	7007	
	,	Name	High Island 95	High Island 155		High Island 82	_	High Island 131	High Island 175	CB-2	CB-3	High Island 77	High Island 80	High Island 81	High Island 117	High Island 148	High Island 110		Island			High Island 37	Island	High Island 69	B-1	High Island 49	High Island 72	Island	High Island 56	Island	High Island 125	Island	High Island 109	Island	High Island 78	High Island 115		High Island 53	
		Block	ast South A-264	High Island East South A-267	High Island East South A-268	High Island East South A-269	ist South A-273	South	ist South A-281	1st South A-283	ist South A-283	East South A-298	South	East South A-309	High Island East South A-313	ist South A-315	ist South A-317	ist South A-317				South	South		1st South A-330	1st South A-330	ist South A-330	ist South A-330	Island East South A-334		South			South	st South A-343	East South A-349	st South A-349	st South A-350	
		Lease Area	High Island East	High Island E	High Island Ea	High Island E		High Island Ea	High Island East	High Island East South	High Island East South	High Island Ea	High Island Ea	High Island Ea	High Island Ea	High Island East South	High Island Ea	Island	High Island Ea	High Island East South	High Island Fact		High Island East South	High Island East	Island	Island	Island	High Island East South	High Island East South										

Lease Area Block	Name	Lon, X, W	Lat, Y, S	Water Depth (ft)	Boring Length (ft)	Source	Dafe
sland East South	High Island 119	92	366	257.0	401.5	McClelland Engineers	7/26/77
South	High Island 133	10359	200	311.0	411.0	McClelland Engineers	6/17/78
High Island East South A-368	High Island 40	8142	7367	329.0	72.0	McClelland Engineers	111/73
High Island East South A-368	High Island 41	3275	3390	320.0	71.0	McClelland Engineers	1/1/73
	High Island 42	11827	11589	338.0	72.0	McClelland Engineers	1/1/73
Island East South	High Island 43	8820	7198	331.0	71.0	McClelland Engineers	1/1/73
Island East South	High Island 44	10827	9333	333.0	72.0	McClelland Engineers	1/1/73
Island East South	High Island 45	9358	8855	336.0	72.0	McClelland Engineers	1/1/73
Island East South	High Island 46	8814	4004	319.0	50.5	McClelland Engineers	1/1/73
South	High Island 47	8955	3863	320.0	51.0	McClelland Engineers	1/1/73
South	Island	8800	2600	325.0	61.0	McClelland Engineers	1/1/73
South	High Island 67	2568	5248	352.0	391.5	McClelland Engineers	1/1/74
Island East South	High Island 141	9171	1114	329.0	419.0	McClelland Engineers	6/17/19
Island East South	Island	11848	23	323.0	404.5	McClelland Engineers	1/1/74
Island East South	High Island 157	11162	3664	312.0	373.0	McClelland Engineers	1/27/82
Island East South A	High Island 179	14406	6859	341.0	500.0	McClelland Engineers	11/16/85
Island East South	High Island 164	5454	5450	350.0	501.0	McClelland Engineers	9/23/82
Island East South	Island 1	9637	6302	410.0	248.0	McClelland Engineers	4/1/78
Island East South	High Island 128	6086	6177	405.0	446.5	McClelland Engineers	3/31/78
Island East South	High Island 129	9567.	6389	408.0	250.0	McClelland Engineers	4/1/78
Island South	CB-1			135.0	328.5	National Soil Services	1/5/79
Island South	High Island 144	11736	13246	153.0	314.0	McClelland Engineers	10/6/79
Island South	High Island 135	12004	4829	182.0	376.0	McClelland Engineers	7/29/78
Island South	Island 1	10027	7631	162.0	446.5	McClelland Engineers	4/17/79
Island South	Island	12866	4945	160.0	321.0	McClelland Engineers	4/27/79
Island South	Island 1	3764	3790	163.0	358.0	McClelland Engineers	10/6/79
Island South	Island	220	5513	161.0	427.0	McClelland Engineers	5/25/78
Island South	Island			172.0	326.0	McClelland Engineers	1/1/67
Island South	Island	9541	7926	196.0	104.0	McClelland Engineers	11/7/80
Island South	High Island 59	3671	6216	206.0	512.0	McClelland Engineers	1/1/74
	Island	13904	8392	188.0	302.0	McClelland Engineers	5/21/79
-	Island	2431	7630	187.0	390.0	McClelland Engineers	10/29/81
Island South	Island	12826	13077	177.0	456.0	McClelland Engineers	
Island South	Island	1363	473	171.0	206.0	McClelland Engineers	
Island South	Island	4967	6910	177.0	476.0	McClelland Engineers	
Island South	High Island 147	20	5534	193.0	411.0	McClelland Engineers	1/28/80 🗐
Island South	High Island 150	13,382	13,560	182.0	420.0	McClelland Engineers	5/2/80 ≒
High Island South A-511	High Island / I	418/	13190	195.0	478.0	McClelland Engineers	1/1/5

				14.1																														I		A	FT	
	Date	1/1/75	5/2/83	8/13/77	1/1/14	1/1/75	2/13/85	1/1/75	1/1/76	1/1/16	8/16/78	1/1/17	1/1/74	8/26/77	1/1/74	1/1/74	1/1/75	1/1/16	1/1/77	7124777	1/1/74	8/18/79	8/19/79	2/24/78	11/13/73	10/4/74	9/30/13	1/1/76	3/9/19	1/1/76	2/9/67	1/20/82	1/1/67	2/15/76	2/10/77	5/18/78	08/07/0	2/5/80
	Source	McClelland Engineers		McClelland Engineers	National Soil Services	National Soil Services	National Soil Services	McClelland Engineers	National Soil Services, Inc	National Soil Services	National Soil Services	McCielland Engineers	McClelland Engineers																									
Boring	Lengin (ft) S	438.0 N		308.0	473.0 N	437.0 N	399.5 N	406.0				452.0 N	366.5 N	450.0 N	373.0 N	326.0 N						_ O		-		339.0 N	_	,'		77.	· · ·				7.		200.0	
Water	ndəri (L)	196.0	204.0	210.0	222.0	220.0	202.0	191.0	199.0	203.0	230.0	244.0	270.0	260.0	277.0	250.0	256.0	332.0	307.0	296.0	337.0	291.0	292.0	276.0	298.0	298.0	345.0	340.0	333.0	40.0	58.0	57.0	55.0		61.0	61.0	0.79	71.0
	Lat. Y. S	12937	3664	2269	7935	11411	7085	14664	11760	7879	8467	771	13736	0	5910	13665	5326	15843/3	<i>L</i> 996	10245	13587	3849	4080	8517			(7949	796	0669	11839					KEOA	10040	14180
	Lon, X, W	6092	4291	6032	7928	7932	6378	6820	4467	7189	7190	1758	15693	15840	12635	8575	9468	2985	3816	3197	4504	11471	11383	13088				3964	C08C	14310	1631					1716	471J 12920	12305
	Name	High Island 85	Island	High Island 124	High Island 61	High Island 73	High Island 174	High Island 84	Island	High Island 98	High Island 136	High Island 114	High Island 65	High Island 121	High Island 50	High Island 51		High Island 102		Island	_	High Island 145	High Island 146	High Island 126	B-1				High Island 140	Matagorda 4	Matagorda 3	Matagorda 16	Matagorda 2	Boring No. 1		CB-1	Matagorda 10	Matagorda 9
	Block	A-511	A-515	A-517	A-519	A-519	A-526	A-531	A-536	A-537	A-542	A-548	A-553	A-555	A-555	A-555	A-561	A-563	A-563	A-563	A-563	A-567	A-567	A-571	A-572	A-572	A-573	A-573	A-582	444	481	485	485	520	520	220	976	527
	ease Area			High Island South	Island	High Island South	Island	High Island South	Island	High Island South	High Island South	High Island South	Island	Island	sland	Island	Island	Island	Island		Island				High Island South	Matagorda Island	Matagorda Island	Matagorda Island	Matagorda Island Matagorda Island	Matagorda Island								

Name		McClelland Engineers 1/25/87	McClelland Engineers 2/3/78	National Soil Services 9/9/77	McClelland Engineers 5/9/84	m)			10	McClelland Engineers 6/26/83	McClelland Engineers 12/12/83		National Soil Services 3/11/78	7	McClelland Engineers 2/10/79	McClelland Engineers 6/26/82				McClelland Engineers 11/23/82		9	McClelland Engineers 11/30/83				-		National Soll Services 8/31//2 McClelland Engineers 1/1/60	12		∞	McClelland Engineers 8/1/82
Block Name Lon, X, W Lat, Y, S	Boring Length (ft)	254.5	308.0	300.0	304.0	357.0	149.5	74.0	97.0	70.0	288.0	86.5	350.5	303.0	299.0	350.0	302.0	343.0	392.5	399.5	89.0	349.5	390.0 405.0	150.0	158.0	65.5	324.5	300.0	393.0	301.0	0.10	308.0 308.0	245.0	299.0	30.0	0 250.0
Block Name Lon, X, and the state S57 Matagorda 29 S68 Matagorda 13 1 1 1 1 1 1 1 1 1	\(\sigma \)			40				08								04	73	-									tean a		-	7 - 1						i alif Period Period
B10ck and a 557 and a 558	X.	7.3	2882			130	1697		4215		1219	9215	7527	6853	1544			10271	4174	12655	2644	13589	4030 1501		9340	1828	2,679,956	467	m	7 401	2,461,202.00				2,458,774.71	
				CB-1	Matagorda				-	Matagorda	· ·					٠,	Ĭ.,	Matagorda 7	Matagorda 1			Matagorda	Matagorda	CB-1	_	7.7										Mustang Island
그는 그들은 살아가 되어 가장 살아 아이를 보고 다른 모든 모든 모든 모든 모든 모든 물을 쓸 쓸 쓸 쓸 쓸 쓸 쓸 쓸 쓸 때	Area Block					Matagorda Island 565	Matagorda Island 566			· · · · · · · · · · · · · · · · · · ·			Matagorda Island 657															s pı								

																																			DI	R.A	LF.	Γ	
	Date	5/19/84	3/3/85	4/1/87	9/25/84	8/24/83	1/1/75	8/25/84	11/27/85	11/17/79	3/30/80	3/26/80	11/7/85	9/11/84	1/1/57	9/13/73	4/16/83	4/14/82	1/1/54	1/1/57	1/1/54	1/1/56	11/7/82	1/1/77	1/1/16	1/1/77	10/30/82	3/28/87	6/23/81	3/22/84	1/1/83	9/23/84	1/26/83	7/13/73	11/13/79	11/13/79	1/5/14	1/3/74	10/3/84
	Source	McClelland Engineers	McClelland-Engineers	McClelland Engineers	National Soil Services	National Soil Services	McClelland Engineers	Greer & McClelland	Greer & McClelland	Greer & McClelland	McClelland Engineers	McClelland Engineers	McClelland Engineers	McClelland Engineers	National Soil Services	McClelland Engineers																							
Boring	(ft)	100.0	199.0	306.0	342.0	302.0	435.5	402.0	400.0	400.0	393.5	376.5	141.0	245.0	59.0	254.0	203.0	204.5	161.0	130.0	121.5	89.5	400.0	448.5	463.0	439.0	303.0	369.0	249.0	370.0	390.0	400.0	134.0	233.5	300.0	300.0	254.0	254.0	320.0
Water	(ft)	55.0	48.0	129.0	183.0	0.89	165.0	123.0	210.0	273.0	213.0	212.0	268.0	281.0	40.0	46.0	47.0	42.0	40.0	44.0	38.0	42.0	258.0	258.0	260.0	264.0	281.0	322.0	61.0	230.0	257.0	242.0	244.0	77.5	82.0	82.0	0.09	60.0	199.0
	Lat. Y. S.	6169	6525	14357		1135			13124	403	13043	11425	189	7011	3700		0.629,069	2918	3750	2783	1200	1000		5338	2480	9519	,	9306	1194	7162	13425	1378	3729					27.7.1	14/5/
	Lon, X, W	4286	1104	15283	1 (2055			13247	4963	14060	12787	7113	573	2500		2,425,035.0	4572	650	2691	2500	2750		12774	12796	827		8712	_				7594					501	10C
	Name	Island			Island			Mustang Island 27	Mustang Island 29	Mustang Island 12	Mustang Island 13	Mustang Island 14	Mustang Island 28		Mustang Island 5	Boring No. 1	CB-1	Mustang Island 16	Mustang Island 2	Mustang Island 4	Mustang Island 1	Island 3	Mustang Island 18	Mustang Island 10	Mustang Island 9		Mustang Island 19	Mustang Island 33				North Padre Island 5	e Island	Boring No. 1	CB-1		Boring No. 1	Well #1	South Facile Island 1
	Block	773	773	781	784	797	831	847	A-11	A-16	A-20	A-20	A-25	A-36	s881	2883	s926	s943	s945	s947	s948	s952	A-065	A-085	A-085	A-086	A-152	A-164					East A-72	1048	1064	1064	1066	1000	•
	Lease Area	Mustang Island	Mustang Island	Mustang Island	Mustang	S Mustang Island	Mustang Island	Mustang Island	Mustang Island East	North Padre Island	North Padre Island East	North Padre Island East			South Padre Island	South Faure Island E.																							