

**FLUVIAL-ESTUARINE SEDIMENTATION
TEXAS GULF COAST**

Literature Synthesis

by

William A. White and Thomas R. Calnan

Prepared for
Texas Parks and Wildlife Department
Resource Protection Division
in accordance with Interagency Contract (88-89) 0820

Bureau of Economic Geology
W. L. Fisher, Director
The University of Texas at Austin
Austin, Texas 78713

September 1988

CONTENTS

Introduction	1
Background and Scope of Study.....	1
Texas Bay-Estuary-Lagoon Systems.....	2
Origin of Texas Estuaries.....	4
General Setting	9
Climate	9
Salinity	19
Bathymetry	21
Relative Sea-Level Rise.....	22
Eustatic Sea-Level Rise.....	24
Subsidence	25
Subsidence in the Houston-Galveston Area.....	28
Faulting and Subsidence.....	30
Effects of Subsidence and Faulting on Texas Coastal Wetlands.....	33
Characteristics of Major Rivers Discharging into Coastal Basins.....	39
Historical Discharge and Loads.....	44
Effect of Reservoir Development on River Sediment Load.....	46
Sediment Trapping.....	54
Effects Downstream from Reservoirs.....	61
Classification of Rivers by Sediment Load.....	64
Deltaic Sedimentation	66
Delta Development.....	66
Development of Deltaic Marshes.....	74

Marsh Sediments.....	75
Marsh Degradation.....	77
Life Cycle of a Subdelta in a Delta System.....	79
Marsh Maturation and Aggradation.....	81
Biodeposition	95
Texas Deltaic Marshes--Development and Current State.....	101
Colorado River Delta.....	101
Recent Changes in State of the Colorado River Delta.....	111
Guadalupe River Delta.....	115
Recent Changes in State of the Guadalupe River Delta.....	118
Trinity River Delta.....	120
Recent Changes in State of the Trinity River Delta.....	127
Recent Changes in State of Wetlands in Other Texas Fluvial-Deltaic Areas.....	129
San Jacinto River.....	129
Neches River.....	131
Lavaca River.....	136
Nueces River.....	137
Bay-Estuary-Lagoon Sedimentation.....	140
Sediment Composition.....	140
Sources of Sediment.....	148
Turbidity	152
Effects of Marine Grasses on Sedimentation.....	158
Trap Efficiencies of Bays.....	164
Importance of Episodic Events on Fluvial-Estuarine Sedimentation.....	168
Are Texas Bays Getting Shallower or Deeper?.....	174
Contribution of Fluvial Sediment to Bay Sedimentation.....	180

The Filling of Lavaca Bay -- An Example.....	180
Trends in Relative Sea-Level Rise and Fluvial-Estuarine Sedimentation.....	183
Corpus Christi-Nueces Bay System.....	185
San Antonio-Espiritu Santo Bay System.....	186
Lavaca-Matagorda Bay System.....	189
Galveston-Trinity Bay System.....	193
Macrobenthos-Sediment Relationships.....	201
Introduction	201
Trophic Structure.....	204
Biomass and Density.....	206
Benthic Assemblages.....	218
Marshes	221
Mudflats and Sandflats.....	222
Species Diversity and Species Richness.....	223
Summary	232
Summary and Preliminary Conclusions.....	235
References	240
Appendix: Annotated bibliography.....	277

FIGURES

1.	Bay-estuary-lagoon and major fluvial-deltaic systems along the Texas Gulf Coast.....	3
2.	Sea-level changes during the past 18,000 years, as interpreted by various authors.....	5
3.	Fluvial environments and facies deposited along the incised lower Trinity River valley, Galveston-Houston area.....	7
4.	Fluvial and deltaic systems of the Colorado and Brazos Rivers in the Bay City-Freeport area.....	8
5.	Development of the Texas shoreline through the late Pleistocene falling sea level stage.....	10
6.	Average annual precipitation in Texas.....	18
7.	Subsidence at selected sites along the Texas coast based on tide gauge records.....	26
8.	Cumulative area in the Texas Coastal Zone affected by land-surface subsidence in excess of 30 cm (1 ft) between 1943 and 1973.....	27
9.	Land-surface subsidence in the Houston-Galveston area.....	29
10.	Subsidence as reflected by differences in mean monthly tide stage relative to the gage record at a designated base station.....	31
11.	Subsidence over an oil and gas field.....	32
12.	Changes in the distribution of wetlands near Jones Bay and Swan Lake near Galveston.....	35
13.	Block diagram of changes in wetlands that can occur along an active surface fault.....	36
14.	Changes in the distribution of wetlands affected by surface faults.....	37
15.	Land subsidence profile based on benchmark-releveling data along State Highway 87 on Bolivar Peninsula.....	38
16.	River and coastal basins in Texas.....	40
17.	Structural elements that affect Texas coastal rivers and Basins.....	41
18.	Longitudinal channel profiles for major Texas rivers.....	42
19.	Characteristics of some major coastal rivers in Texas.....	43

20.	Annual discharge, silt load, and drainage area for some major Texas rivers.....	45
21.	Annual streamflow and suspended sediment load of the Trinity River.....	47
22.	Suspended-sediment load for the Trinity and San Jacinto Rivers for the period 1936 through 1975.....	48
23.	Annual streamflow and suspended load of the Brazos River.....	49
24.	Comparison of suspended-sediment loads of the Brazos and Colorado Rivers, and Rio Grande.....	49
25.	Annual streamflow and suspended load of the Colorado River.....	50
26.	Suspended load of the Colorado River at Austin and Columbus.....	50
27.	Annual streamflow and suspended load of the Lavaca River.....	51
28.	Annual streamflow and suspended load of the Navidad River.....	51
29.	Annual streamflow and suspended load of the Guadalupe River.....	52
30.	Annual streamflow and suspended load of the San Antonio River.....	52
31.	Annual streamflow and suspended load of the Nueces River.....	53
32.	Concentration of suspended-sediment load for the Nueces River.....	53
33.	Reservoirs located on the Gulfward half of Texas.....	55
34.	Water discharge and suspended-sediment load for the Colorado River at Yuma, Arizona.....	56
35.	Suspended-sediment load (percent by weight) of the Trinity River at Romayor, and cumulative authorized water storage in reservoirs of the Trinity River basin.....	57
36.	Annual suspended sediment loads of the Brazos, Trinity, and Colorado Rivers, with declines in load related to reservoir development.....	58
37.	Trap efficiency curve.....	60
38.	Comparison of trap efficiency curves from Brune (1953) and Churchill (1948).....	60
39.	Pre-dam and post-dam longitudinal profiles of the Brazos River below Lake Whitney Dam, indicating net deposition along several miles of the channel.....	63
40.	Modern depositional systems and sediment dispersal patterns, northwest Gulf of Mexico.....	65

41.	Major components of a river system.....	67
42.	Development of mid-channel shoals and branching distributaries.....	71
43.	Components of a delta developing in shallow water.....	71
44.	Seaward migration of depositional environments in high-constructive deltas, typical of bay-head deltas.....	72
45.	Development of deltaic facies, including constructional and destructional cycles.....	73
46.	Relative percentages of sand, silt, and clay in salt marsh sediments, Sapelo Island, Georgia.....	76
47.	Development and deterioration of deltaic marshes.....	78
48.	Evolutionary sequence of the Mississippi River Delta distributaries.....	80
49.	Effect of subsidence from sediment compaction on distributary and interdistributary areas.....	82
50.	Growth and development of the Guadalupe delta at the head of San Antonio Bay.....	83
51.	Seasonal variations in aggradation rates of salt-water marshes in Barataria Basin, Louisiana.....	90
52.	Percentages of inorganic (mineral) sediments in marsh soils in Mississippi delta basins arranged in order of increasing age.....	92
53.	Relationship between rates of marsh aggradation and relative sea-level rise between 1940 and 1980.....	93
54.	Comparison of sediment delivery to near-shore areas along the eastern U.S. and Mississippi Delta coasts between the early-1900's and 1970.....	96
55.	Theoretical biodeposition cycle in an estuary.....	98
56.	Comparison of size and shape of the Trinity, Guadalupe, and Colorado River deltas with the Atchafalaya River delta and crevasse-splay deposits of the modern Mississippi River.....	102
57.	Historical development of the Colorado River delta between 1908 and 1941.....	104
58.	Increase in area of the Colorado River delta for the period 1908-1953.....	105
59.	Deterioration or destruction of the southeast lobe of the Colorado River delta plain, as determined by comparing shoreline positions on aerial photographs taken in 1943, 1952, and 1965.....	107

60.	Cross sections of the southeast lobe of the Colorado River delta.....	108
61.	Changes in marsh distribution between 1956 and 1979 on the Colorado River delta at Tiger Island Cut.....	112
62.	Map showing idealized spatial relationship of depositional environments of the Guadalupe River delta prior to formation of the Traylor Cut subdelta.....	116
63.	Generalized cross section of the Guadalupe delta showing facies.....	116
64.	Shoreline changes along the Traylor Cut subdelta, based on aerial photographs, 1929 to 1982.....	119
65.	Changes in the distribution of wetlands between 1957 and 1979 on the seaward half of the Guadalupe River delta.....	121
66.	Trinity River delta.....	122
67.	Rates of shoreline change along Trinity Bay and Trinity delta, 1930 to 1982.....	128
68.	Changes in the distribution of wetlands between 1956 and 1979 of a subsiding segment of the San Jacinto river near Houston, Texas.....	130
69.	Aerial photograph of the lower reaches of the San Jacinto River.....	132
70.	Cumulative reservoir storage capacity for selected rivers along the Texas coast.....	133
71.	Changes in the distribution of wetlands between 1956 and 1978 in the lower Neches River valley near the head of Sabine Lake.....	135
72.	Changes in the distribution of wetlands between 1958 and 1979 at Menefee Flat in the Lavaca River valley.....	138
73.	Index for sediment distribution maps shown in figures 74-79.....	141
74.	Distribution of sediments in Sabine Lake.....	142
75.	Distribution of sediments in Galveston Bay.....	143
76.	Distribution of sediments in Matagorda Bay.....	144
77.	Distribution of sediments in San Antonio and selected bays to the southwest.....	145
78.	Distribution of sediments in Corpus Christi and Nueces Bays.....	146
79.	Distribution of sediments in southern Laguna Madre.....	147
80.	Conceptual model of sediment sources and associated deposition relative to the magnitude of the source.....	150

81.	Depositional features typical of Texas bay-estuary systems related to sediment sources and processes.....	151
82.	Sediment trap efficiency curves of Brune (1953) applied to estuaries.....	165
83.	Sediment transport by the Colorado River.....	170
84.	Model depicting possible effect of frontal passage on the transportation and distribution of fine-sediment from the bay-estuary-lagoon system to the South Texas shelf.....	173
85.	Depth changes in Texas bays.....	176
86.	Galveston Bay depth changes determined by comparing bathymetric surveys ranging from 1852 to 1867 with surveys in 1934.....	181
87.	Map showing depth to Pleistocene surface beneath Lavaca and Matagorda Bays.....	182
88.	Cross section down Lavaca River valley showing sediments that have accumulated in Lavaca Bay above the Pleistocene surface.....	184
89.	Paleobathymetric curve over the past 10,000 years for Lavaca Bay.....	184
90.	Lavaca-Matagorda bay system depth changes determined by comparing bathymetric surveys ranging from 1856 to 1872 with surveys in 1934.....	191
91.	Bathymetric changes for the Trinity-Galveston bay system for the period 1968-1977.....	195
92.	Annual-average volume of sediments dredged from Anahuac Channel along the Trinity River.....	199
93.	Relationship of sediment volume dredged from Anahuac Channel with annual streamflow along the Trinity River.....	200
94.	Conceptual model illustrating the role of the estuarine benthos in food chain dynamics and nutrient recycling.....	202
95.	Average measures for sediment characteristics and benthic macroinfaunal biomass at three stations in Corpus Christi Bay from 1981 to 1983.....	217
96.	Density (numbers of individuals/m ²) by month of benthic macroinvertebrates at sandy and sandy mud stations in Espiritu Santo and Lavaca Bays.....	220
97.	Scattergram of total species and percent sand in the bays of the Galveston-Houston area.....	229
98.	Scattergram of total species and percent sand in the bays of the Port Lavaca area.....	230

TABLES

1.	Dimensions and average tidal range of bays-estuaries-lagoons.....	11
2.	Generalized characteristics of active coastal processes and conditions in the Beaumont-Port Arthur area.....	13
3.	Generalized characteristics of active coastal processes and conditions in the Galveston-Houston area.....	14
4.	Generalized characteristics of active coastal processes and conditions in the Bay City-Freeport area.....	15
5.	Generalized characteristics of active coastal processes and conditions in the Port Lavaca area.....	16
6.	Generalized characteristics of active coastal processes and conditions in the Corpus Christi area.....	17
7.	Factors influencing deltaic sedimentation.....	70
8.	Marsh aggradation (vertical accretion) rates measured in coastal Louisiana and along the U. S. Atlantic coast.....	87
9.	Rates of biodeposition of mud by some suspension and deposit feeders on the Texas coast.....	99
10.	Minimum, maximum, and mean turbidity measurements.....	155
11.	Comparisons of predicted trapping efficiencies of other estuaries with those in Texas.....	166
12.	Relation of bay fill to suspended sediment load.....	178
13.	Bay-estuary sedimentation and net deposition rates.....	179
14.	Annual average suspended load of the San Antonio and Guadalupe Rivers combined.....	187
15.	Feeding type of benthic macroinvertebrate species.....	207
16.	Biomass and density of benthic macroinvertebrates.....	211
17.	Mean density of benthic macroinvertebrates in marine grasses.....	212
18.	Mean density of benthic macroinvertebrates versus percent sand.....	213
19.	Mean density versus mean percent sand for benthic macroinvertebrate assemblages.....	214
20.	Biomass and density of benthic macroinvertebrates in marshes, mudflats, and sandflats.....	215
21.	Mean percent sand, species diversity, and mean number of species of benthic macroinvertebrate assemblages.....	226

22.	Mean percent sand, species diversity, and mean number of benthic macroinvertebrates in marine grassflat assemblages.....	227
-----	---	-----

FLUVIAL-ESTUARINE SEDIMENTATION TEXAS GULF COAST

Literature Synthesis

INTRODUCTION

Deltaic and associated alluvial areas at the mouths of rivers that discharge into the bay-estuary-lagoon system along the Texas coast are the sites of extensive salt-, brackish-, and fresh-water marshes that are essential components of biologically productive estuarine systems. These bay-head depositional systems are constructed primarily by fluvial sediments, sediments transported and deposited by the major rivers that enter estuarine waters. The loss of over 10,000 acres of wetlands in alluvial and deltaic areas of the Neches and San Jacinto Rivers has emphasized the need to examine in more detail the processes that establish and maintain, as well as degrade these important natural resources along the Texas coast.

Background and Scope of Study

This report, which is a synthesis of published and unpublished data that focuses on fluvial-deltaic and estuarine sedimentation, and associated interactive processes, is part of a study funded by the Texas Parks and Wildlife Department and Texas Water Development Board with funds allocated by the Texas Legislature for comprehensive studies of the effects of freshwater inflows on the bays and estuaries of Texas. Another phase of this project involves field studies of the Trinity and Colorado River deltas to investigate, among other processes, current rates of sedimentation (aggradation) in deltaic marshes (White and Calnan, in preparation).

Most of the fresh water inflow studies, past and ongoing, of Texas estuaries have focused on inundation, cycling and exchange of nutrients, salinity patterns, and production of fisheries (TDWR, 1982). A significant part of the past research effort

has dealt with the need to inundate deltaic wetlands (through fresh water inflows) in order to export nutrients into the estuarine system. Although habitat maintenance was one of the objectives of the investigations, little emphasis was placed on the geological processes that play a critical role in the construction of the deltaic and alluvial systems on which the biologically productive wetlands develop.

Among the objectives of this study is to focus on the sedimentary and associated interactive processes that develop, maintain, and/or degrade the environments. Information is provided on the present and historical (including geologic) role of fluvial sediments--sediments carried by rivers--in developing and maintaining estuarine habitats, with emphasis on wetlands, marine grassflats, and benthic communities. Among the interactive processes that are analyzed are: riverine discharge and associated sediment loads, aggradation (vertical accretion) rates, erosion rates, subsidence (both natural and human-induced), sea-level rise, sediment supply from sources other than rivers, sediment sinks, biodeposition, and effects of human activities on sediment budgets.

TEXAS BAY-ESTUARY-LAGOON SYSTEMS

LeBlanc and Hodgson (1959) distinguish between estuaries and lagoons along the Texas coast by differences in origin and by differences in alignment with respect to the Gulf shoreline. Estuaries, which have formed as a result of valley entrenchment, are generally aligned perpendicular to the coast, while lagoons, which formed as a result of shoreline processes, are generally aligned parallel to the coast (fig. 1). Texas coastal water bodies, although perhaps most properly termed coastal lagoons (Morton and McGowen, 1980), have been variously referred to as bays, estuaries, and lagoons in the literature. For simplification in this report, the terms "bays" and "estuaries" are generally used interchangeably.




Figure 1. Bay-estuary lagoon and major fluvial deltaic systems along the Texas Gulf Coast (Modified from LeBlanc and Hodgson, 1959)

To properly understand the current processes that affect modern estuarine environments, it is helpful to look briefly at past processes and events that have developed and helped shape the estuaries and associated deltaic areas along the Texas coast.

Origin of Texas Estuaries

Texas estuaries have evolved as deeply eroded valleys were flooded by a rising sea level (Price, 1933, 1947). Sea levels have fluctuated dramatically during the past million years as a result of alternating cooling and warming climatic cycles that have produced glacial and interglacial periods. During periods of glaciation, large amounts of water are locked up in continental ice sheets resulting in dramatic drops in worldwide sea level (fig. 2). Over the past 2.5 to 3 million years, evidence shows that there has been at least 8 major cycles in which sea level has fallen (producing a lowstand) and risen (highstand) (Beard and others, 1982). The last major cycle is pertinent to this discussion because it has had the most profound effect on our modern bay-estuary-lagoon system and Gulf shoreline.

During the most recent major period of glaciation and lowstand (ending about 18,000 years ago) sea level along the Texas coast was approximately 400 ft feet below today's level (LeBlanc and Hodgson, 1959; Curray, 1960); this placed the shoreline between 50 and 140 miles offshore (LeBlanc and Hodgson 1959). During the lowstand, the base levels of rivers along the Texas coast, as well as throughout the world, were lowered and extensive down-cutting and erosion formed deep valleys. The valleys cut by rivers along the Texas coast range from about 50 to 150 ft (15 to 45 m) deep (relative to today's sea level) along the valley axes near the gulfward margin of the bay shorelines (Shepard and Moore, 1955, Fisk, 1959, Behrens, 1963, McEwen, 1969, Byrne, 1975, and Wright, 1980). As sea level rose the valleys were flooded.




Figure 2. Sea-level changes during the past 18,000 years, as interpreted by various authors. All curves show a general trend of rising sea level although the authors interpret minor sea-level fluctuations differently. A generalized curve would show sea level 4,500 years ago to have been about 15 lower than at present. (From Fisher and others, 1973.)

Approximately 4,500 years before present (B.P.) the rise in sea level slowed as it approached today's level. During the rise in sea level, rivers were meandering within their valleys depositing large point-bar sand bodies and extensive overbank mud sheets (Fisher and others, 1972). The erosion of the valleys and subsequent deposition was recorded by most streams, including the Trinity River, for example (fig. 3). Deposition of sediment in the entrenched valleys by rivers like the Trinity could not keep pace with sea-level rise, and a major portion of their valleys was drowned producing estuaries like Trinity Bay. Trinity Bay and other Texas bays have since been enlarged by shoreline erosion, and the deeper parts of the submerged valleys have been filled slowly by bay sediment. The major sediment depositional centers in the estuaries are the bayhead deltas. These active depositional features have filled much of the lower stream valleys and have advanced (prograded) over bay muds at the heads of the bays. Along the gulfward reach of the estuaries, a series of barrier islands and peninsulas has formed, restricting the exchange of marine and estuarine waters to relatively narrow tidal inlets.

In contrast to rivers like the Trinity, three Texas rivers--Brazos, Colorado, and Rio Grande--filled their estuaries with sediments and constructed broad deltaic plains that protrude into the Gulf of Mexico (Price, 1947; LeBlanc and Hodgson, 1959). The fact that some rivers have filled their valleys and others have not is related to the nature of their drainage basins and sediment supply (Fisher and others, 1972). The depositional patterns developed by the Brazos and Colorado Rivers indicate that, in the past, they merged to form a single alluvial system that rapidly filled their valleys and estuaries (LeBlanc and Hodgson, 1959; McGowen and others, 1976) (fig. 1). The Colorado River has more recently abandoned its ancestral courses (which included Caney Creek) and now flows in a more narrow alluvial valley that intersects Matagorda Bay southwest of its previous course near the Brazos (LeBlanc and Hodgson, 1959, Wilkinson and Basse, 1978) (fig. 4). The Brazos River discharges




Figure 3. Fluvial environments and facies deposited along the incised lower Trinity River valley, Galveston-Houston area. (From Fisher and others, 1972.)




Figure 4. Fluvial and deltaic systems of the Colorado and Brazos Rivers in the Bay City-Freeport area. (From McGowen and others, 1976.)

into the Gulf near Freeport, and has built a small delta at its mouth. The evolution of the Texas coastal shoreline is depicted in figure 5.

General Setting

The Texas Gulf shoreline is nearly 370 miles (595 km) long and consists of seven major estuarine systems (fig. 1). The total open water surface area of the estuaries at mean high water (MHW) is 1,532,430 acres (620,634 hectares) (Diener, 1975). Many of the bays are comparatively small, having surface areas of less than 2,000 acres (809 hectares), but some are very large with surface areas of over 100,000 acres (40,469 hectares) at MWH (table 1). These systems are characterized by diverse climatic conditions and hydrologic features. Tables 2 through 6 provide a summary of some of the coastal processes and climatic conditions that affect these diverse bay-estuary lagoon systems and their adjacent Gulf shorelines.

Climate

The bay-estuary-lagoon system along the Texas coastal zone is affected by a diverse climatic setting that systematically changes down the coast. Climate along the upper Texas coast in the Beaumont-Port Arthur and Galveston-Houston areas is humid (Thorntwaite, 1948) (tables 2 and 3). Average annual precipitation ranges from approximately 50 inches (127 cm) in the Beaumont-Port Arthur area to 40 inches (102 cm) in the Galveston-Houston area (fig. 6). Between 1931 and 1960, the upper Texas coast had from 5 inches (13 cm) to more than 12 inches (30 cm) of excess moisture from precipitation after evaporation and plant transpiration (Fisher and others, 1972; Fisher and others, 1973). Temperatures generally range from average winter lows in the mid- 40's (°F) (7° to 9°C) to average summer highs in the low to mid- 90's (°F) (33° to 35°C). Two principal wind regimes dominate the Texas Coastal Zone -- persistent, southeasterly winds from March through November






Figure 5. Development of the Texas shoreline through the late Pleistocene falling sea level stage (A), the early recent sea level stage (B), and recent standing sea level stage (C). (From LeBlanc and Hodgson, 1959.)

Table 1. Dimensions and average tidal range of bays-estuaries-lagoons. Modified from Diener, 1975).

Location	Surface area		Depth at mean low water**		Average
	Mean low water	Mean high water*	Maximum	Average	tidal
	-----Acres-----		-----Feet-----		
<u>Sabine Lake</u>					
Sabine Lake	43,960	44,830	24	5.1	0.2
Sabine Pass	1,360	1,360	40		
<u>Galveston Bay</u>					
East Bay	33,370	33,690	12	3.3	1.2
Trinity Bay	83,310	86,240	17	5.2	1.0
Galveston Bay (upper)	69,890	70,080	42	5.7	1.0
Galveston Bay (lower)	89,380	90,390	44	6.5	1.4
Lake Anahuac (Turtle Bay)	4,660	4,850	5	2.1	--
Scott-San Jacinto Bay	3,230	4,310	40	1.8	1.0
Clear Lake	1,260	1,280	14	2.7	0.9
Dickinson Bay	1,520	1,540	6	2.1	0.7
Moses Lake (Dollar Bay)	2,130	2,140	36	5.2	0.5
Offats Bayou	1,180	1,200	28	14.5	1.0
Jones Lake	1,040	1,050	2	1.6	1.0
West Bay	44,390	45,420	25	3.9	0.9
Chocolate Bay	4,890	4,920	12	2.6	0.6
Bastrop-Oyster Bay	9,690	10,410	20	3.2	0.7
<u>Matagorda Bay</u>					
East Matagorda Bay	37,810	39,080	5	3.4	0.4
Matagorda Bay	167,570	170,130	36	8.0	0.7
Oyster Lake	2,450	2,570	12	2.7	0.5
Tres Palacios Bay	9,440	9,860	12	4.1	0.6
Turtle Bay	1,280	1,760	5	2.5	0.6
Carancahua Bay	12,160	12,300	7	3.8	0.5
Salt, Redfish Lakes	920	950	4	1.2	0.5
Keller Bay	4,770	4,850	8	3.2	0.6
Lavaca Bay	39,970	40,080	36	4.2	0.7
Swan Lake	860	880	3	1.4	0.1
Lavaca River Estuary	740	760	13	8.0	0.2
Chocolate Bay	1,440	1,760	12	2.7	0.5
Powderhorn Lake	2,890	2,970	4	2.2	0.7
Cedar Lakes Complex	3,760	3,840	12	2.1	0.5

Table 1 (cont.)

Location	Surface area		Depth at mean low water**		Average
	Mean low water	Mean high water*	Maximum	Average	tidal
	-----Acres-----		-----Feet-----		
<u>San Antonio Bay</u>					
Espiritu Santo Bay	38,940	40,630	14	5.9	0.3
San Antonio Bay	76,530	77,700	12	4.6	0.3
Guadalupe Bay	2,070	2,090	9	2.7	0.2
Mission Lake	1,820	2,400	--	--	--
Hynes Bay	6,580	6,610	3	2.4	0.2
Ayers Bay	2,220	2,550	12	3.2	0.3
Mesquite Bay	8,080	9,220	12	3.4	0.2
<u>Copano Bay</u>					
St. Charles Bay	8,410	8,730	6	3.6	0.2
Mission Bay	3,760	3,760	2	1.9	0.1
Copano Bay	41,740	42,930	9	3.7	0.3
Port Bay	1,650	2,000	9	2.2	0.2
Mission Lake	100	100	--	--	--
Aransas Bay	56,220	59,220	25	7.8	0.4
<u>Corpus Christi</u>					
Redfish Bay	9,630	13,420	17	2.0	0.4
Corpus Christi Bay	73,820	75,560	40	10.5	0.7
Nueces Bay	18,470	18,550	3	2.2	0.4
Oso Bay	5,070	5,070	15	1.6	0.9
<u>Laguna Madre</u>					
Upper Laguna Madre	47,240	68,360	12	2.8	0.7
Lower Laguna Madre	175,160	329,740	26	4.7	1.0
South Bay-La Badilla					
Grande Complex	4,380	7,300	36	1.5	1.5
Baffin Bay	31,870	32,610	12	7.7	0.5
Alazan Bay	13,860	14,750	4	2.9	0.5
Cayo del Infernillo	700	1,630	2	0.7	0.5
Laguna Salada	3,230	3,530	6	2.8	0.5
Cayo del Grullo	4,470	8,470	6	2.8	0.5

*Does not include peripheral marsh areas.

**Exclusive of navigation channels.

Table 2. Generalized characteristics of active coastal processes and conditions in the Beaumont–Port Arthur area. From White and others (1987).

Climatic zone:	Humid (<i>Thorntwaite, 1948</i>)
Average annual precipitation:	51.5 to 55.7 inches/yr (130.8 to 141.5 cm/yr) (<i>Fisher and others, 1973</i>)
Dominant wind directions:	Southeasterly, northerly (<i>Fisher and others, 1973</i>)
Astronomical tidal range: Gulf shoreline (Sabine Pass jetty) Diurnal range: Bay shoreline (average, Sabine Lake):	2.5 ft (0.8 m) (<i>U.S. Department of Commerce, 1978</i>) 0.2 ft (6 cm) (<i>Diener, 1975</i>)
Tidal current velocities: Sabine Pass Average maximum flood: Average maximum ebb:	2.7 ft/s (0.8 m/s) (<i>U.S. Department of Commerce, 1983</i>) 2.9 ft/s (0.9 m/s) (<i>U.S. Department of Commerce, 1983</i>)
Wave height (Gulf): (Caplan, Texas) Onshore wave height:	Between 2.5 and 3.5 ft (0.8 and 1.1 m) about 65% of the time (<i>U.S. Army Corps of Engineers, 1956</i>)
Direction of net longshore sediment transport:	Southwesterly (<i>Fisher and others, 1973</i>)
Maximum recorded hurricane surge height on open coast: At Sabine Pass: Near High Island:	6.7 ft (2.0 m) above MSL (<i>U.S. Department of Commerce unpublished data</i>) 4.2 ft (1.3 m) MSL (<i>Bodine, 1969</i>)
Hurricane probability:	12% in any one year (<i>Simpson and Lawrence, 1971</i>)
Gulf shoreline change, Sabine Pass to Bolivar Roads from 1882 to 1974:	Net rates are minor or moderate except for extreme net accretion of 26 and 28 ft/yr (7.9 to 8.5 m/yr) at points adjacent to the jetties at Sabine Pass and Galveston Harbor. Net erosion occurred at most other coastal points and ranged from 1 to 17.4 ft/yr (0.3 to 5.3 m/yr) and averaged 6.4 ft/yr (2.0 m/yr) (<i>Morton, 1975</i>).
Subsidence: Sabine Pass: Estimated rate based on tide-gauge records: Magnitude related to withdrawal of oil, gas, associated ground water, and solution mining of sulfur:	0.5 inch/yr (1.25 cm/yr) during 1960–1969 (<i>Swanson and Thurlow, 1973</i>) Generally less than 0.5 ft (0.15 m) but locally exceeding 1 ft (0.3 m) during 1918–1977 (<i>Ratzlaff, 1980</i>)

Table 3. Generalized characteristics of active coastal processes and conditions in the Galveston-Houston area. From White and others (1985).

Climatic zone:	Humid (<i>Thorntwaite, 1948</i>)
Average annual precipitation:	41.8 to 51.5 inches/yr (106.2 to 130.8 cm/yr) (<i>Fisher and others, 1972</i>)
Dominant wind directions:	Southeasterly, northerly (<i>Fisher and others, 1972</i>)
Average wind speed (in 1978 at Texas City):	6.8 knots (12.6 km/hr) (<i>Shew and others, 1981</i>)
Astronomical tidal range: Gulf shoreline (Galveston Pleasure Pier) Mean diurnal: Bay shoreline (mean):	2.1 ft (0.6 m) (<i>U.S. Department of Commerce, 1978</i>) 0.5 to 1.4 ft (0.2 to 0.4 m) (<i>Diener, 1975</i>)
Tidal current velocities: Bolivar Roads Average maximum flood: Average maximum ebb:	3.3 knots (1.7 m/sec) (<i>Bernard and others, 1959</i>) 4.3 knots (2.2 m/sec) (<i>Bernard and others, 1959</i>)
Wave height (Gulf): (Caplan, Texas) Onshore wave height:	Between 2.5 and 3.5 ft (0.8 and 1.1 m) about 65% of the time, (<i>U.S. Army Corps of Engineers, 1956</i>)
Direction of net longshore sediment transport:	Southwesterly (<i>Fisher and others, 1972</i>)
Maximum hurricane surge height on open coast:	12.7 ft (3.9 m) above MSL (<i>Bodine, 1969</i>)
Hurricane frequency:	12% in any one year (<i>Simpson and Lawrence, 1971</i>)
Gulf shoreline change, Bolivar Roads to San Luis Pass from 1850-52 to 1973-74:	Total gain from accretion of 1,074 acres and loss from erosion of 1,183 acres; net loss of 109 acres (<i>Morton, 1977</i>)
Subsidence: Pasadena - Houston Ship Channel area:	8.5 to 9 ft (2.6 to 2.7 m) during 1906-1973 (<i>Ratzlaff, 1980</i>)
Faulting: Houston metropolitan area:	Offset by at least 160 faults (<i>Verbeek and Clanton, 1981</i>)

Table 4. Generalized characteristics of active coastal processes and conditions in the Bay City-Freeport area. From White and others (in press a).

Climatic zone:	Subhumid (<i>Thorntwaite, 1948</i>)
Mean annual precipitation:	40.6 to 49.2 inches (103.1 to 124.9 cm) (<i>McGowen and others, 1976</i>)
Dominant wind directions:	Southeasterly, northerly (<i>McGowen and others, 1976</i>)
Astronomical tidal range: Gulf shoreline (Freeport Harbor) Diurnal range: Mean: Bay shoreline (Matagorda Bay):	 1.8 ft (0.5 m) (<i>U.S. Department of Commerce, 1978</i>) .09 ft (0.3 m) (<i>U.S. Department of Commerce, 1978</i>) 0.5 to 0.7 ft (0.2 m) (<i>McGowen and Brewton, 1975</i>)
Direction of net longshore sediment transport:	Southwesterly (<i>McGowen and others, 1976</i>)
Estimated maximum hurricane surge height at Freeport:	9.5 ft (2.9 m) above MSL (<i>Bodine, 1969</i>)
Hurricane frequency:	Probability of occurrence along 50-mi (80.5-km) segment of coast in Bay City-Freeport area: 14% in any one year (<i>Simpson and Lawrence, 1971</i>)
Net rate of Gulf shoreline erosion over period of about 120 yr: Matagorda Peninsula: New Brazos River delta to Brown Cedar Cut:	 About 2 to 3 ft/yr (0.6 to 0.9 m/yr), on average, although exceeding 10 ft/yr (3 m/yr) just south of Brown Cedar Cut (<i>Morton and others, 1976</i>) 12.7 ft/yr (3.9 m/yr), on average (<i>Morton and Pieper, 1975</i>)
Subsidence: Freeport: Estimated rate based on tide-gauge records: Magnitude related to withdrawal of ground water:	 0.44 inch/yr (1.12 cm/yr) during 1959-1971 (<i>Swanson and Thurlow, 1973</i>) Generally less than 0.5 ft (0.15 m) but locally exceeding 2 ft (0.6 m) during 1906-1973 (<i>Ratzlaff, 1980</i>)

**Table 5. Generalized characteristics of active coastal processes and conditions, Port Lavaca area.
From White and others (in press b).**

Climatic zone:	Subhumid (<i>Thorndwaite, 1948</i>)
Mean annual precipitation:	32 to 39 inches (81.3 to 99.1 cm) (<i>McGowen and others, 1976</i>)
Dominant wind direction:	Southeast, north (<i>McGowen and others, 1976</i>)
Astronomical tidal range:	
Gulf shoreline (Pass Cavallo)	
Diurnal range:	1.4 ft (0.4 m) (<i>U.S. Department of Commerce, 1978</i>)
Mean:	0.7 ft (0.2 m) (<i>U.S. Department of Commerce, 1978</i>)
Bay shoreline (Port Lavaca)	
Diurnal range:	0.7 ft (0.2 m) (<i>U.S. Department of Commerce, 1978</i>)
Mean:	0.3 ft (0.1 m) (<i>U.S. Department of Commerce, 1978</i>)
Direction of net longshore sediment transport (Gulf shoreline):	Southwesterly (<i>McGowen and others, 1976</i>)
Estimated peak hurricane surge height on open coast near Port O'Connor:	12.3 ft (3.7 m) m.s.l. (<i>Bodine, 1969</i>)
Hurricane frequency:	Probability of occurrence along 50-mi (80.5-km) segment of coast in Port Lavaca area is 9 percent in any one year (<i>Simpson and Lawrence, 1971</i>)
Net rate of Gulf shoreline accretion or erosion over period of 117 yr:	
Matagorda Island:	Less than 1 ft (0.3 m) of average annual accretion in southern half of island; accretion rates in the northern half range from 1.1 to 9.1 ft (0.3 to 2.8 m) per year. Erosion rates near Pass Cavallo range from 5.1 to 17.3 ft (1.6 to 5.3 m) per year (<i>Morton and Pieper, 1976</i>)
San José Island:	Less than 1.5 ft (0.5 m) of average annual erosion and accretion (<i>Morton and Pieper, 1976</i>)

Table 6. Generalized characteristics of active coastal processes and conditions in the vicinity of Mustang Island, Corpus Christi area (modified from White and Galloway, 1977).

Climatic zone:	Dry subhumid (<i>Thorntwaite, 1948</i>)
Mean annual precipitation:	30 to 32 inches/yr (76 to 81 cm/yr) (<i>Carr, 1967</i>)
Dominant wind directions:	Southeasterly; north-northeasterly (<i>Lohse, 1955</i>)
Average wind speed (in 1980):	12.8 mi/hr (20.6 km/hr) (<i>U. S. Dept. of Commerce, 1980a</i>)
Direction of net sand transport by winds:	Northwesterly (<i>Hunter and others, 1972</i>)
Astronomical tidal range: Gulf shoreline (Port Aransas) Mean diurnal Maximum diurnal Bay shoreline, mean	 1.5 ft (0.48 m) (<i>Hayes, 1965</i>) 2.5 ft (0.76 m) (<i>Collier and Hedgpeth, 1950</i>) Approx. 0.5 ft (0.15 m) (1 ft [0.3 m] lower than Gulf) (<i>Watson and Behrens, 1976</i>)
Tidal current velocities: Aransas Pass Average maximum flood Average maximum ebb Fish Pass Average maximum Usual value	 2.0 ft per second (fps) (0.6 m/s) 1.5 fps (0.5 m/s) (<i>U. S. Dept. of Commerce, 1980b</i>) 3 fps (0.9 m/s) Below 2 fps (0.6 m/s) (<i>Defehr and Sorensen, 1973</i> ; <i>Watson and Behrens, 1976</i>)
Wave height (Gulf): Usual height Mean height	 Below 4 ft (1.2 m) (<i>Davis and Fox, 1972</i>) 2.6 ft (0.8 m) (<i>Watson and Behrens, 1976</i>)
Longshore current velocities (Gulf): Range Average	 0 to 3.9 fps (0 to 1.2 m/s) (<i>Davis and Fox, 1972</i>) 0.38 fps (0.1 m/s) (fall) and 0.7 fps (0.2 m/s) (winter) (<i>Davis and Fox, 1972</i>)
Direction of net longshore sediment transport:	Southwesterly (<i>Lohse, 1955</i> ; <i>Behrens and Watson, 1974</i>)
Average rate of Gulf shoreline erosion over period of about 100 years (Mustang Island):	2.0 ft/yr (0.6 m/yr) (<i>Morton and Pieper, 1977</i>)
Maximum hurricane surge height recorded at Aransas Pass (1919 to 1977):	11.5 ft (3.5 m) (1919) (<i>Price, 1956</i>)
Hurricane frequency: Probability of occurrence along 50-mile (80.5-km) segment of coast in Corpus Christi area	 7% in any one year (<i>Simpson and Lawrence, 1971</i>)




Figure 6. Average annual precipitation in Texas. (Modified from Larkin and Bomar, 1983.)

and short-lived but strong northerly winds from December through February (Fisher and others, 1972; Fisher and others, 1973).

Climate along the middle Texas coast from the Bay City-Freeport area to the Corpus Christi area is subhumid to dry subhumid (Thorntwaite, 1948) (tables 4, 5, and 6). Average annual precipitation ranges from near 50 inches (127 cm) in the Bay City-Freeport area to 32 inches (81 cm) in the Corpus Christi area (fig. 6). Between 1931 and 1960, the middle Texas coast had from 4 inches (10 cm) of excess moisture from precipitation after evaporation in the eastern part of the Bay City-Freeport area to a precipitation deficit of about 12 to 16 inches (30 to 41 cm) in the Corpus Christi area (McGowen and others, 1976a, 1976b; Brown and others, 1976). Temperatures generally range from winter minimum lows of 46 to 48°F (8° to 9°C) to average maximum summer highs in the low to mid- 90's (33° to 35°C).

Along the lower Texas coast in the Kingsville and Brownsville-Harlingen, climate is semiarid (Thorntwaite, 1948). The annual rainfall ranges from 26 inches (66 cm) along the southern coastline in the Brownsville-Harlingen area to 34.5 inches (87.6 cm) along the northern coastline of the Kingsville area (Brown and others, 1977, 1980). Between 1931 and 1960 the average annual mean free-air temperature in the area was about 73 to 73.5°F (Brown and others, 1977, 1980).

Salinity

Water salinities vary considerably both between bay-estuary-lagoon systems and within each system, in part because of the regional variations both in fresh-water inflows from rivers and streams and in salt water interchange from tidal passes. Compounding the complexity in each system are seasonal and cyclic climatic variations that produce substantially higher than normal salinities during dry periods and lower than normal salinities during wet periods. Salinities fluctuate over a wider range in shallow bay margins than in mid-bay areas, but average salinities are generally lower in the bay margins. High surface salinities in the bays are recorded during periods of low rainfall and stream discharge.

Average salinities in Texas estuaries range from a low of about 2 parts per thousand (ppt) in the Sabine-Neches estuary (Armstrong, 1982) to over 54 ppt in the upper part of Cayo del Grullo in the Baffin Bay system (Brown and others, 1977). On the upper coast, salinities in Sabine Lake generally range from less than 10 ppt in the upper part of the lake to between 10 and 20 ppt in the tidally influenced lower part (Fisher and others, 1972). Salinity decreases with increasing distance from Sabine Pass, such that salinity is slightly lower in the central part of Sabine Lake than in the lower open-bay area.

Of the bays in the Galveston Bay system, salinities are generally highest in West Bay, followed, in order of decreasing average salinity, by Galveston, East, and Trinity Bays. Average salinities in West Bay are generally more than 15 ppt and range into the 30's, which is in marked contrast to Trinity Bay, where average salinities range from less than 5 to about 10 ppt (White and others, 1985). Salinities in Trinity Bay can drop to 0 ppt or exceed 25 ppt.

Salinity data for the Matagorda Bay system, including Matagorda, East Matagorda, Tres Palacios, Carancahua, and Lavaca Bays, can be found in Ward and others (1980) and Jones and others (1986). Data for San Antonio Bay are reported in Matthews and others (1974). Salinities in Lavaca and Matagorda Bays generally increase toward Pass Cavallo. Average salinities in Matagorda Bay are generally above 20 ppt and range into the 30's. In East Matagorda Bay salinities range from 15 ppt near the Colorado River delta to 17.4 in the northeastern part of the bay. Matthews and others (1974) found that salinities in upper San Antonio Bay ranged from approximately 0.5 to 9.0 ppt and in the most gulfward region from approximately 6.0 to 26.0 ppt.

In the Corpus Christi area, salinities are generally highest in upper Laguna Madre, followed, in order of decreasing average salinity, by Corpus Christi, Redfish, Aransas, Nueces, and Copano Bays (Holland and others, 1975; Brown and others,

1976; and Hildebrand and King, 1978). Average salinities in upper Laguna Madre are generally above 30 ppt and range into the 40's and occasionally higher, which is in marked contrast to Copano Bay, where average salinities range from about 10 to 15 ppt increasing toward the mouth of the bay. Average monthly median salinities in the upper part of Copano Bay fluctuate around 14.4 ppt, and in the upper part of Nueces Bay fluctuate around 21.7 ppt (Texas Department of Water Resources, 1981a). Monthly mean salinities in mid-Corpus Christi Bay vary, as demonstrated by measurements in 1973 that show a high of 35 ppt in February and a low of 15.6 ppt in October (Holland and others, 1975).

Average salinities for the bay-estuary-lagoon system in the Kingsville area, including Baffin, Alazan, Cayo del Grullo, and that part of upper Laguna Madre near Baffin Bay, are probably the highest on the Texas coast. Calculated average surface salinities of Baffin Bay and associated water bodies range from a low of just less than 50 ppt in Laguna de los Olmos to over 54 ppt in Cayo del Grullo (Brown and others, 1977). Average surface salinities for eight stations in Baffin Bay range from approximately 40 ppt in May 1966 to 70 ppt in December 1964 (Behrens, 1966).

In the Brownsville area, salinities generally increase from the southern end of lower Laguna Madre at Port Isabel to north of Port Mansfield (Brown and others, 1980; Espey, Huston and Associates, Inc., 1981). Salinities in the Port Isabel area range from 23 to 36 ppt and are influenced by the exchange of Gulf water through Brazos Santiago Pass. Salinity at the northern end of lower Laguna Madre ranges from 20 to 40 ppt and averages about 38 ppt.

Bathymetry

Bathymetric data (table 1) are taken from Diener (1975) and represent averages of the most recent soundings at mean low water (MLW) exclusive of navigation channels. Average depths range from 0.7 ft (0.2 m) in Cayo del Infernillo of the Baffin Bay system to 14.5 ft (4.4 m) in Offats Bayou of the Galveston Bay system.

Average depths of the larger bay-estuary-lagoon systems range from 2.8 ft (0.9 m) in upper Laguna Madre to 10.5 ft (3.2 m) in Corpus Christi Bay. Many of the bays are shallow with average depths of less than 4 ft (1.2 m). Maximum depths occur in the navigation channels and near the tidal passes.

Tides

Astronomical tidal variations in Texas estuaries are small compared to estuaries of the Atlantic and Pacific Coasts. In the Gulf of Mexico the principal variations in the tide are due to changing declination of the moon (U. S. Department of Commerce, 1979). Tidal range in the northwest Gulf of Mexico during maximum declination of the moon is about 2.6 ft (0.8 m) and at minimum declination about 0.7 ft (0.2 m) (Ward and others, 1980). Meteorological events are more important than astronomical tides, as far as the estuaries are concerned, as they alternately expose and flood the greatest area of tidal flat and marsh (Collier and Hedgpeth, 1950).

The most noticeable fluctuations in bay levels are caused by direction and force of the wind or wind tides. The amount of open-bay fetch and direction of wind tides control the effectiveness of wind tidal activity (Brown and others, 1976). For example, broad fetch, as in Trinity Bay and the western arm of Matagorda Bay, and persistent southeast winds aligned with the axis of the bay, result in high wind tides that may build tide heights two to three ft (0.6 to 0.9 m) above normal (Holliday, 1973). Frontal passage, such as during a norther, can also drastically affect the wind tides and estuarine water levels.

Relative Sea-Level Rise

The bay-estuary-lagoon system is affected by many interactive processes. One of the most important at work along the Texas coast today is relative sea-level rise.

Stated very simply, for example, if coastal wetlands do not receive and trap sufficient sediments (organic or inorganic) so that the aggradation rate (vertical accretion) is equal to or greater than the rate of relative sea-level rise, the wetlands will ultimately be lost and replaced by open water. If bay-estuary-lagoon sedimentation rates do not keep pace with rates of relative sea-level rise, water depths will increase through time; if the sedimentation rates exceed relative sea-level rise, water depths will decrease and the bay or estuary may eventually fill with sediment.

Relative sea-level rise as used here refers to a rise in sea level with respect to the surface of the land, whether it is caused by actual sea level rise or land-surface subsidence; the current general trend along the Texas coast involves both of these processes working together.

Sea-level fluctuations occur for a variety of reasons and on broad spatial and temporal scales. Nummedal (1983) reviewed sea-level fluctuations and how they are affecting the coast of Louisiana, and Morton and Price (1987) present information on Late Quaternary sea-level fluctuations and their relation to the shallow-water depositional complexes on the Texas coastal plain and shelf. Nummedal (1983) notes that sea-level changes can be categorized into two fundamental groups that operate on global and local scales. Among the global factors are (1) the volume of the ocean basins controlled by sea floor spreading rates, sedimentation, and opening and closing of marginal seas, and (2) the volume of oceanic water which has changed in response to glaciation and, possibly, water temperature. Local factors include subsidence of continental margins, movement of the land surface along faults, compaction due to dewatering of sediments, and many atmospheric factors.

Various methods have been used to measure changes in mean sea level; a primary method during historic time is to compare records from tide gauges and examine the trends over as long a period as the records allow (Marmer, 1954; Swanson and Thurlow, 1973). Using this method along the Gulf coast, Swanson

and Thurlow (1973) concluded that subsidence is an important factor with regard to mean sea-level rise.

The discussion below will touch briefly on eustatic (global) sea-level rise but will focus principally on compactional subsidence, natural and human-induced, because these processes appear to be the most significant with respect to the Texas coast.

Eustatic Sea-Level Rise

It is generally accepted that sea level is rising on a worldwide (eustatic) basis (Hicks, 1978, Gornitz and others, 1982), apparently in response to a global warming trend resulting from increases in atmospheric CO₂ and the resulting "greenhouse" effect (Hansen and others, 1981), which can add volume to the oceans through glacial melt water (Etkins and Epstein, 1982; Meier, 1984) and perhaps thermal expansion (Gornitz and others, 1982). The worldwide rate of sea-level rise, based on tide-gauge records over the past century, is about 1.2 mm per year (Gornitz and other, 1982). For the Texas coast, this rate is not nearly as significant as relative sea-level rise due to compactional subsidence (Swanson and Thurlow, 1973).

Subsidence

There are many causes of subsidence (Nummedal, 1983) including regional downwarping or tilting of the earth's crust due to loading, which is significant over a geologic time frame along the Texas coast (Winker, 1979), but is not as significant when viewed over an historic time frame. Holdahl and Morrison (1974) reported slight subsidence along the Gulf Coast region ranging between 0.0 and 1.5 mm/yr, in addition to anomalous subsidence in Houston and Corpus Christi areas (discussed in succeeding sections). The most significant subsidence along the Texas coast appears to be due to compactional subsidence, especially as affected by subsurface fluid withdrawal (water, oil, gas, and, locally, sulfur).

Using tide-gauge records along the Texas and Louisiana coasts, and comparing them with records from the more stable (tectonically or geologically) Florida coast,


Swanson and Thurlow (1973) concluded that subsidence is occurring along the Texas Coast at rates of from 0.5 to 1.2 cm/yr. Furthermore, they found that rates for the period from 1959 to 1971 are higher than rates before 1959 (1948-1959) (fig. 7). Highest rates along the Texas Coast are at Sabine Pass, Freeport, and Port Aransas (1.12 to 1.28 cm/yr) and the lowest rate is at Port Isabel (0.49 cm/yr). Subsidence in the Freeport area was believed to be due to the delta environment of the Brazos River, and subsidence in the area of Pier 21 at the mouth of West Bay near Galveston to faulting and withdrawal of oil and gas (Swanson and Thurlow, 1973). Swanson and Thurlow (1973) noted that their data, which show an overall tendency toward subsidence along the entire coast, supported Shepard and Moore (1960), who suggested that much of the Texas coast could be subsiding due to increasing sediment overburden and compaction of underlying fine sediment.

Compactional subsidence occurs as sediments are consolidated, generally as a result of compressive forces from overlying material (sediments and water) and the dewatering of the compacting sediments either naturally or as influenced by withdrawal of fluids.

The highest rates of subsidence along the Texas coast have been caused by withdrawal of underground fluids, principally water (Winslow and Doyel, 1954; Gabrysch, 1969; Brown and others, 1974; Gabrysch and Bonnet, 1975; Kreitler, 1977; Ratzlaff, 1980; Gabrysch, 1984). Production of oil and gas can also cause subsidence (Pratt and Johnson, 1926; Kreitler, 1977; Gustavson and Kreitler, 1976, and Verbeek and Clanton, 1981). Extreme local subsidence has occurred in relation to sulfur mining around salt domes along the Texas Coast (Ratzlaff, 1980, and Mullican, 1988).

Brown and others (1974) reported that along the Texas Coastal Zone, the rates of subsidence, both in terms of area impacted and drops in surface elevation, have progressively increased since 1940 (fig. 8).

Subsidence Estimated From Tide Gages



Swanson & Thurlow (1973)

ESTIMATED RATE OF SUBSIDENCE

LOCATION	TIME PERIOD	RATE, mm/yr
Sabine Pass	1960-1969	12.5
Galveston (PP)	1959-1970	8.6
Freeport	1959-1971	11.2
Port Aransas	1959-1969	12.8
Port Isabel	1959-1971	4.9


 QA 7531

Figure 7. Subsidence at selected sites along the Texas coast based on tide gauge records. (From Swanson and Thurlow, 1973.)




Figure 8. Cumulative area in the Texas Coastal Zone affected by land-surface subsidence in excess of 30 cm (1 ft) between 1943 and 1973. (From Brown and others, 1974.)

Subsidence in the Houston-Galveston Area

The most extensive subsidence, both in terms of vertical and areal magnitude, due to withdrawal of fluids is in the Houston-Galveston area (Ratzlaff, 1980), where more than 2.7 m (9ft) and possibly as much as 3.0 m (10 ft) of subsidence has occurred in the vicinity of the Houston Ship Channel between 1906 and 1978 (almost 2.7 m [9 ft] of which occurred during 1943 to 1978) (Gabrysch, 1984). Maximum subsidence is in the center of a subsidence "bowl" that encompasses an area from near Freeport (where another smaller "bowl" is centered) to an area north of the Trinity River delta (fig. 9). Average maximum rates of subsidence at the center of the "bowl" have been as high as 122 mm/yr (0.4 ft/yr) for the period 1964 to 1973 (Gabrysch and Bonnet, 1975).

According to Gabrysch and Bonnet (1975), subsidence due to withdrawal of ground water from an artesian aquifer results from a decrease of hydraulic pressure and attendant movement of water from clays to adjacent sands leading to compaction of the clays (Gabrysch and Bonnet, 1975). Most of the compaction is permanent because of the inelastic nature of the clay; thus, even with total recovery of artesian pressure, less than 10 percent rebound can be expected (Gabrysch and Bonnet, 1975).

Methods of measuring subsidence, by the USGS, include conventional leveling, extensometers, and tide gauges (Gabrysch, 1985). Conventional leveling is the most frequently used method, and involves comparing the elevations of benchmarks that have been measured at different times using precise leveling techniques. Borehole extensometers have been used at specific locations to determine small changes in elevations; extensometers can provide very precise, continuous records with information on the compacting interval, but they are costly to install and have small areal application (Gabrysch, 1985). Subsidence can be determined by comparing tide gauge records from two different stations, but this method is less precise than leveling and




Figure 9 Land-surface subsidence in the Houston-Galveston area, 1943-1973. (From Gabrysch and Bonnet, 1975.)

extensometers. Gabrysch (1985) reported that evaluation of tide gauge data from five stations in Galveston Bay and Buffalo Bayou indicated that elevation change of less than 150 mm (0.5 ft) and perhaps as little as 30 mm (0.1 ft) could be detected (fig. 10).

Other parts of the Texas coastal region, in addition to the Houston-Galveston and Freeport areas, where subsidence has occurred include (1) Beaumont-Port Arthur where an area over Spindletop Dome has subsided about 1.5 m (5 ft) between 1925 and 1977, and an area near Port Acres has subsided approximately 0.9 m (3 ft) between 1959 to 1977, primarily due to withdrawal of oil and gas and associated ground water, (2) Jefferson and Matagorda Counties inland from Matagorda Bay where subsidence of more than 0.46 m (1.5 ft) occurred during 1943 to 1973 as a result of ground-water withdrawals, and (3) the western part of Corpus Christi where more than 1.5 m (5 ft) of subsidence occurred during 1942 to 1975 due to withdrawals of oil, gas, and associated ground water (Ratzlaff, 1981).

Faulting and Subsidence

In some areas along the Texas Coast, subsidence may be accompanied by active surface faults. A good example is a fault in the Saxet oil and gas field west of Corpus Christi (Price, 1933). The fault has produced a 2 m (6 ft) scarp at the surface (Gustavson and Kreitler, 1976). Profiles constructed from releveled lines across benchmarks show rapid increases in subsidence at the fault (fig. 11). Subsidence rates during the period from 1950 to 1959 were 70 mm/yr (0.22 ft/yr), which was an increase of almost twice the rate of 40 mm/yr (0.14 ft/yr) during the period of 1942 to 1950 (fig. 11). Gustavson and Kreitler (1976) theorized that an increase in gas production from 1950 to 1959 may have been responsible for compaction of shallow reservoir sands on the downthrown side of the fault leading to differential subsidence and accelerated fault movement. Evidence of the fault can be seen where





Figure 10. Subsidence as reflected by differences in mean monthly tide stage relative to the gage record at a designated base station (Galveston Railroad Causeway in the above case). Change in the difference between mean water level as measured by tide gauges at stations a few miles apart reflects the change in elevation between the stations. (From Gabrysch, 1984.)

Land Subsidence Over Saxet Oil and Gas Field Corpus Christi, Texas



From Kreitler (1977)

QA 7526
Bureau of Economic Geology

Figure 11. Subsidence over an oil and gas field. (From Kreitler, 1977.)

it crosses highways and other structures. Lower elevations in the subsidence bowl inhibit drainage of surface water locally and promote ponding of water.

Although the fault in the Saxet field in the Corpus Christi area is a good example of an active surface fault, the major zone of surface faulting along the Texas coast is in the Houston-Galveston area where 95 linear miles (150 km) of faulting has been reported (Reid, 1973; Brown and others, 1974). Surface faults correlate with, and appear to be extensions of, subsurface faults in many areas (Weaver and Sheets, 1962; Van Siclen, 1967; Kreitler, 1977; Verbeek and Clanton, 1981). Most of the surface faulting in the Houston metropolitan area has apparently taken place during the last few decades (Verbeek and Clanton, 1981), largely due to fluid withdrawal (water, oil, and gas), which has reinitiated and accelerated fault activity (Reid, 1973; Kreitler, 1977; Verbeek and Clanton, 1981).

The range in measurable vertical displacement of surface traces of faults is from 0 to 3.9 m (12 ft) (Reid, 1973). Rates of fault movement commonly range between 5 mm/yr and 20 mm/yr (0.2 in/yr and 0.8 in/yr) (Verbeek and Clanton, 1981), but many exceed 40 mm/yr (1.6 in/yr) (Van Siclen, 1967; Reid, 1973; Everett and Reid, 1981). Movement along surface faults apparently occurs episodically (Reid, 1973). Highways, railroads, industrial complexes, airports, homes, and other structures placed on active faults in the Houston area have undergone millions of dollars worth of damage annually (Clanton and Verbeek, 1981).


Effects of Subsidence and Faulting on Texas Coastal Wetlands

Subsidence in the Houston-Galveston area has had a significant effect on wetlands in the area (Johnston and Ader, 1983; White and others, 1985). One of the most dramatic examples of wetland losses due to subsidence is along the San Jacinto River. More than 560 hectares (1,389 acres) of fluvial woodlands, swamps, and marshes were displaced by open water between 1956 and 1979 (White and others, 1985). The lower reach of the San Jacinto River, near its confluence with

Buffalo Bayou and the Houston Ship Channel, is in an area of subsidence (fig. 9) caused mostly by ground-water withdrawal (Gabrysch and Bonnet, 1975).




The change in wetlands along the lower San Jacinto River valley (discussed more thoroughly in a later section) is pronounced because of the proximity of the valley to the center of maximum subsidence. However, wetlands associated with other streams and valleys located around the Trinity and Galveston Bay system are also changing as a result of human-induced subsidence and accompanying relative sea-level rise. Replacement of marshes by open water is occurring along the bay margins as well (fig. 12). Changes in the distribution of wetlands as a result of natural compactional subsidence has also been reported along the Texas coast (Donaldson and others, 1970; McGowen and Brewton, 1975; White and others, 1988).

Faults have affected marshes from the Freeport area to Sabine Lake (White and others, 1985, 1987, and in press a). As vertical displacement occurs along a fault that intersects a marsh, more frequent and eventually permanent inundation of the wetland surface on the downthrown side of the fault can lead to replacement of marsh vegetation by open water if marsh sedimentation rates do not keep pace with submergence rates (fig. 13). This has occurred at several locations along the upper Texas coast as exemplified by a marsh system on the bay side of Bolivar Peninsula, where approximately 500 hectares (1,230 acres) of salt-water marsh has been replaced primarily by "barren" shallow subaqueous flats and open water (fig. 14). In this area, at least two surface faults intersect marsh substrates. Benchmark releveled profiles along State Highway 87 indicate the faults are active; a marked increase in subsidence occurs on the downthrown side (fig. 15). More than 25 faults that cross wetlands along the upper coast (Freeport area to Sabine Pass) have been identified on aerial photographs. One fault has affected wetlands that have developed on modern fluvial-deltaic deposits along the lower Neches River valley at




Index Map

EXPLANATION



1956



1979

QA1774

Figure 12. Changes in the distribution of wetlands near Jones Bay and Swan Lake near Galveston. Note increase in open water in 1979, apparently as a result of subsidence. (From White and others, 1985.)

Example of Changes in Wetlands Along a Fault







Figure 13. Block diagram of changes in wetlands that can occur along an active surface fault. There is generally an increase in low marshes, shallow subaqueous flats, and open water on the downthrown side of the fault relative to the upthrown side.



Wetland Map Unit	1956		1979		Net Change	
	Acres	Hectares	Acres	Hectares	Acres	Hectares
Water	140	57	1130	458	+990	+401
Marsh	3480	1409	2560	1037		
Marsh-covered tidal channels	640	259				
TOTAL MARSH	4120	1668	2560	1037	-1560	-631
Barren tidal flats			270	109	+270	+109
Dredged spoil/uplands	520	211	720	292	+200	+81
TOTAL	4780	1936	4680	1896	-100*	-40*

*Most of the loss in total area is probably due to erosion of marsh along the margins of East Bay. Historical monitoring indicates that this shoreline is undergoing erosion at rates of up to 3 ft (1 m) per year (Paine and Morton, in preparation).
(Areas were calculated using a square-count method; smallest squares used were equivalent to 6.4 acres, or 2.6 hectares).

QA 1774

Figure 14. Changes in the distribution of wetlands affected by surface faults. Increases in the areal extent of water in 1979 relative to 1956 are apparently related to localized subsidence and active faults (D = downthrown side of fault, U = upthrown side). (From White and others, 1985.)

Land Subsidence Profile Along Bolivar Peninsula




Figure 15. Land subsidence profile based on benchmark-releveling data along State Highway 87 on Bolivar Peninsula. The increase in subsidence along the profile indicates it crosses an active fault, probably an extension of the fault with the NE-SW strike in figure 14. (Profile from C. W. Kreitler, unpublished data.)

the head of Sabine Lake (White, 1987); this area is discussed in more detail in a later section of this report.

Characteristics of Major River Systems Discharging into Coastal Basins

Major coastal rivers and state-wide drainage basins are shown in figures 1 and 16, respectively. Slopes of the Cenozoic Gulf Coastal Plain are relatively steep across the San Marcos Arch but more gentle across the Houston and Rio Grande Embayments (fig. 17). The gradients of rivers reflect the slopes of these different tectonic provinces (Morton and Donaldson, 1978). Rivers such as the Nueces and Guadalupe on the central Texas coast were affected by uplift along the San Marcos Arch and have steeper gradients than rivers crossing the Houston Embayment (Sabine, Neches, Trinity, and Brazos Rivers) and Rio Grande Embayment (Rio Grande River) (fig. 18).

Winker (1979) reported that modern rivers that cross the Texas coastal plain can be characterized in terms of drainage basin, discharge, and sediment load; these parameters allow calculation of ratios that further define the nature of the river systems (fig. 19). As indicated by Winker, the dominance of the Rio Grande, Colorado, and Brazos over other Texas rivers is clearly reflected in drainage basin area, average annual discharge, and sediment load; the relatively systematic decline in runoff depths toward the southwest reflects the climatic gradient (fig. 6). The fact that average denudation rates show a more complex pattern than runoff depths may be the result of human modification of the drainage basins (Winker, 1979; the data he used was pre-1960).

As stated by Milliman and Meade (1983), two basic methods are used to estimate the amount of sediments transported by rivers to the oceans (and estuaries): one method estimates the denudation of the land (as illustrated in the bottom graph in fig. 19), and the other estimates the mass carried by the rivers. Of these two,





Figure 16. River and coastal basins in Texas. (From Texas Water Development Board.)



(MURRAY, 1961, & HARDIN, 1962)

Figure 17. Structural elements that affect Texas coastal rivers and basins. (From Wright 1980, after Murray 1961 and Hardin, 1962.)




Figure 18. Longitudinal channel profiles for major Texas rivers. (From Morton and Donaldson, 1978.)




Figure 19. Characteristics of some major coastal rivers in Texas. (From Winker, 1979.)

the denudation method yields a much larger estimate of sediment load because it includes a large amount of sediment that never reaches the oceans (Milliman and Meade, 1983). Factors controlling denudation, a term commonly used as a synonym for erosion, include (1) size of drainage basin, (2) precipitation and vegetation, (3) elevation and relief, (4) rock types, and (5) man (Ritter, 1967).

The Soil Conservation Service, U.S. Department of Agriculture, has estimated the amount of sediment eroded from Texas land areas (based on general land use and soils maps) using the universal soil loss equation (Greiner, 1982). The universal soil loss equation uses factors related to those listed above for denudation including rainfall, soil erodibility, topography, crop management, and erosion control. In addition, the Soil Conservation Service has investigated sedimentation by water in Texas (USDA, 1959; Greiner, 1982). A comparison of previous sedimentation surveys (median date of which is 1947) with a study in 1979 for various lakes and reservoirs in Texas indicated rates of sediment accumulation had declined (Greiner, 1982). Greiner attributed the lower rates in sedimentation to several factors including changes in land use, by noting there have been (1) significant decreases in amount of cropland (a large producer of sediment), (2) continuous implementation of soil conservation measures since 1935, and (3) construction of flood-prevention dams and other trapping elements since about 1954. Effects of reservoirs on stream sediment loads is discussed in a following section.

Historical Discharge and Load

A comparison of discharge and sediment load of the major Texas rivers, based on early records (up to 1954), shows that the Trinity, Brazos, Colorado, and Rio Grande had (in the past) the highest average annual discharge, ranging from about 2.7 million acre-feet to slightly more than 5 million acre-feet; these same rivers also had the highest annual silt loads, ranging from the Brazos with 20,148 acre-feet to the Trinity with 3,622 acre-feet (fig. 20). The Trinity River load is only about 18%




Figure 20. Annual discharge, silt load, and drainage area for some major Texas rivers. Based on data prior to 1955. (From LeBlanc and Hodgson, 1959.)

of that of the Rio Grande. The high sediment loads characterizing the Rio Grande and the ancestral Brazos-Colorado couplet contributed to the filling of their respective paleovalleys (see preceding section on origin of bays), and to the progradation of deltas into the Gulf. Suspended sediment load transported by the other streams was considerably lower, ranging from 88 acre-feet per year for the Lavaca River to 636 acre-feet for the Sabine River. None of these rivers, including the Trinity, have completely filled their paleovalleys (estuaries), but have constructed deltas at the head of the bays (bayhead deltas).

Historical trends in streamflow and sediment loads of major Texas coastal Rivers are shown in figures 21 to 32. Major reductions in sediment load are apparent for the Trinity, Brazos, Colorado, Nueces, and Rio Grande. For example, for the Brazos River, Curtis and others (1973) reported the annual sediment load to be about 16 million tons; as noted by Milliman and Meade (1983), this load is one-half of that (32 million tons) presented in an earlier report by Holeman (1968). The annual average sediment load of the Brazos River presented by Winker (1971) was 31 million tons (fig. 19) for the period of 1925 to 1947 (Richmond station). The average annual load for the Brazos from 1947 to 1979 was approximately 12 million tons per year (fig. 23), or less than 40 percent of that for the earlier period. Although reductions in river sediment load can be partly attributed to land use changes and to the continuous implementation of soil conservation measures in the drainage basins since 1935 (Greiner, 1982), the apparent major contributing factor to the decreased sediment supply in many of the streams in Texas is reservoir development (fig. 33).

Effect of Reservoir Development on River Sediment Load

The effect of reservoirs on sediment load is well documented. One of the most often cited examples is the Colorado River that discharges into the Gulf of California (Milliman and Meade, 1983). The average annual suspended-sediment load of the




Figure 21. Annual streamflow and suspended sediment load of the Trinity River. (Data from Stout and others, 1961; Adey and Cook, 1964; Cook, 1967; Cook, 1970; Mirabal, 1974; Dougherty, 1979; and unpublished records from Texas Water Development Board, available through TNRIS.)




Figure 22. Suspended-sediment load for the Trinity and San Jacinto Rivers for the period 1936 through 1975. (From Morton and McGowen, 1980.)




Figure 23. Annual streamflow and suspended load of the Brazos River. Suspended load is plotted by tons $\times 10^{-1}$. (Source of data same as for figure 21.)




Figure 24. Comparison of suspended-sediment loads of the Brazos and Colorado Rivers, and Rio Grande; gaging stations on the Brazos at Richmond, on the Colorado near Eagle Lake, and on the Rio Grande at Brownsville. Both the Brazos and Colorado show a decrease in suspended load. The load at Brownsville was monitored only after the completion of Falcon Dam (1954). (From McGowen and others, 1977.)




Figure 25. Annual streamflow and suspended load of the Colorado River. (Source of data same as for figure 21.)




Figure 26. Suspended load of the Colorado River at Austin and Columbus. (Source of data same as for figure 21.)




Figure 27. Annual streamflow and suspended load of the Lavaca River. (Sources of data same as for figure 21.)




Figure 28. Annual streamflow and suspended load of the Navidad River. (Sources of data same as for figure 21.)




Figure 29. Annual streamflow and suspended load of the Guadalupe River. (Sources of data same as for figure 21.)




Figure 30. Annual streamflow and suspended load of the San Antonio River. (Sources of data same as for figure 21.)




Figure 31. Annual streamflow and suspended load of the Nueces River. (Sources of data same as for figure 21.)




Figure 32. Concentration (by weight) of suspended sediment for the Nueces River. The Three Rivers measuring station is upstream from Lake Corpus Christi, and the Mathis station is downstream. No data for dashed segments. Data from sources given in figure 21. (From Morton and Paine, 1984.)

Colorado River at Yuma, Arizona, for the period 1911-1916 was about 235 million tons; as a result of reservoir development and increased use of water for irrigation, the suspended-sediment load was reduced (for 1965-1967) to an average annual discharge of about 0.153 million tons, or about seven-hundredths of a percent of the past value (Curtis and others, 1973) (fig. 34). It should be noted that Meade (1969) (in discussing the Atlantic Coastal Plain) concluded that because of the vast amount of sediment contributed to river systems by human activities such as agriculture (estimated to have increased sediment yield by ten-fold) and urbanization, that even though sediment is partly intercepted by reservoirs on rivers, the sediment loads reaching the Atlantic Coast are larger than if the drainage areas were still in a natural state.

Sediment Trapping

Reservoir development in the drainage basins of the Nueces, Trinity, Brazos, and Colorado Rivers in Texas show significant effects on sediment load and discharge (figs. 32, 35, and 36). A considerable amount of data has been accumulated on reservoir sedimentation and trapping efficiency in order to determine reservoir life expectancy (Vanoni, 1975; Strand and Pemberton, 1982). The trap efficiency of a reservoir is a measure of the amount of inflowing sediment that is deposited or retained in the reservoir. Large reservoirs (storage capacities of greater than 10,000 acre-ft) trap virtually 95 to 100 percent of the incoming sediment (Vanoni, 1975; Leopold and others, 1964). Williams and Wolman (1984) citing the U. S. Army Corps of Engineers (1960) reported that the trapping efficiency of Denison Dam on the Red River (Texas and Oklahoma) was 99.2 percent during the first 12 years after closure. The amount retained is primarily controlled by two factors, the average velocity of flow through the reservoir and the sediment size; clay may remain in suspension long enough to pass through the reservoir, but sand will not (Vanoni, 1975). Brune (1953) developed curves for estimating the trapping efficiency of reservoirs by plotting




Figure 33. Reservoirs located on the Gulfward half of Texas. (From Greiner, 1982.)




Figure 34. Water discharge and suspended-sediment load for the Colorado River at Yuma, Arizona. The sharp decline in the curves in the mid-1930's is related to reservoir development. (From Curtis and others, 1973.)




Figure 35. Suspended-sediment load (percent by weight) of the Trinity River at Romayor, and cumulative authorized water storage in reservoirs of the Trinity River basin. (Sources of data for suspended load same as in figure 21; reservoir data from Texas Water Development Board, 1973.) (From Paine and Morton, 1986.)




Figure 36. Annual suspended sediment loads of the Brazos, Trinity, and Colorado Rivers, with declines in load related to reservoir development. Sediment load along the vertical axis is concentration in percent by weight. (From USDA, 1972.)

sediment trapped in 44 reservoirs against the ratios of reservoir capacity to annual inflow (fig. 37). This concept is discussed in a later section with respect to trapping efficiency of estuaries. Churchill (1948), based on reservoirs of the Tennessee Valley Authority, presented a method of calculating reservoir trapping efficiency by relating the percentage of incoming sediment that passes through a reservoir to the reservoir's sediment index, which is the period of retention (reservoir capacity/daily inflow) divided by mean velocity of the flow through the reservoir. Strand and Pemberton (1982) compared curves of the two methods (fig. 38). Several Texas reservoirs, are included in figures 37 and 38. The trap efficiencies of Possum Kingdom on the Brazos River and Buchanan on the Colorado River are about 98 percent. The trap efficiency of Lake Corpus Christi, which is about 74 percent for 1942-1948 (Brune, 1953), has increased more recently (reference next paragraph).

The Soil Conservation Service of the U.S. Department of Agriculture (USDA, 1959) in an inventory of sedimentation in Texas, has presented data on reservoir sedimentation, including reservoir trap efficiency, within the various river basins. In addition, a considerable amount of data on sediment outflow from three to four reservoirs in Texas has been accumulated through measurements by Texas Water Development Board (Cook, 1970; Mirabal, 1974; Dougherty, 1979). Data on reservoir trap efficiency defined by ratios of reservoir capacity to reservoir inflow is presented by Cook (1970) and Mirabal (1974). Leibbrand (1987) estimated the sediment input into Lake Corpus Christi for the period of 1972-1985 to be 5,320 acre-ft (dry); the sediment output during the same period was 117 acre-ft (dry). The amount deposited in the lake was about 5,140 acre-ft (dry), or approximately 97 percent of the sediment that entered.

Detailed sedimentation surveys have been conducted in some reservoirs in Texas. For example, Govin (1973) conducted a sedimentation survey of Lake Buchanan on the Colorado River in north-central Texas. He found that about 101,400 acre-ft of




Figure 37. Trap efficiency curve. (From Brune, 1953.)




Figure 38. Comparison of trap efficiency curves from Brune (1953) and Churchill (1948). (From Strand and Pemberton, 1982.)

sediment has accumulated in the lake since 1937 when impoundment began. Among his findings are (1) average rate of sedimentation was 2,800 acre-ft/yr, which represented an annual average loss in reservoir capacity of 0.29 percent, (2) maximum accumulation occurred in the upper lake and in the deep river channel of the lower lake, (3) sediments were predominantly medium and fine-grained silt and clay from the river, and (4) the river's bed load and the coarser fraction of the suspended load must be deposited well above the lake. The average annual denudation rate of the drainage basin was calculated at a minimum of 0.11 acre-ft/mi², which Govin considered to be in reasonable agreement with a rate of 0.32 acre-ft/mi² calculated by Ritter (1967) for the Colorado River in central and west Texas.

Effects Downstream from Reservoirs

Williams and Wolman (1984) compiled a large data set detailing the downstream effect of dams by analyzing 1,817 measurements of 287 cross sections downstream from 21 dams. Although there is wide variation in the post dam water-discharge characteristics from river to river, flood peaks were generally decreased by the dams. Average annual peak discharges were decreased from 3 to 91 percent of the pre-dam values, with the average decrease being about 39 percent. Suspended sediment load decreased markedly for hundreds of kilometers downstream from dams, and on some rivers annual loads did not equal pre-dam loads anywhere downstream. The distance downstream from dams that is required for a river to regain the sediment load equivalent to the pre-dam load varies. In some major rivers, annual sediment loads may not equal pre-dam values for hundreds or thousands of km downstream, if at all. Degradation of channel beds generally occurred during the first decade or two after dam completion. The magnitude of degradation varied from negligible to approximately 7.5 m. The general trend was for bed material to coarsen as degradation proceeded, although this trend may change in later years. Channel width down-

stream from a dam can increase, decrease, or remain constant. Vegetation commonly increased downstream from dams, probably as a result of reduction in peak flows after dam closure.

Degradation or erosion downstream from dams is a common occurrence (Williams and Wolman, 1984) as noted above. However, sedimentation below dams can also occur (for example, Minter, 1976; Woolley, 1985; Hobbs, 1987). Minter (1976) concluded that among the effects caused by construction of dams on the Brazos River and its tributaries was a reduction in peak flood flows and a great reduction in the river's sediment carrying capacity. The construction of Lake Whitney on the Brazos has decreased peak floods by 50 to 65 percent, resulting in a decrease in sediment transporting power (competency) and a local build-up of gravel deposits in the channel at the mouths of tributary streams (Woolley, 1985; Hobbs, 1987). Although a reduction in sediment load below the dam has promoted channel degradation or erosion, net aggradation has occurred near tributary junctions (fig. 39). The inability of the Brazos to transport the size of the gravel delivered to it by its tributaries has led to armoring and stabilization of the clastic deltas (Hobbs, 1987).

Minter (1976) concluded that major dam and reservoir development within the Brazos River basin has significantly reduced the amount of suspended load and bed load reaching the Texas coast. It was estimated that reservoirs are trapping approximately 76 percent of all sand derived from the drainage basin. According to Minter (1976), the amount of sand lost by entrapment plus that lost due to declines in river transporting power can account for the entire increase in rates of Gulf shoreline erosion (Seelig and Sorensen, 1973; Morton and Pieper, 1975) near the mouth of the Brazos, at least since 1937.

In contrast to the findings of Minter (1976), who reported a loss in sand delivered to the coast after reservoir development, Isphording (1986), in a study of sediments deposited in Apalachicola Bay, Florida, reported an increase in sand




Figure 39. Pre-dam and post-dam longitudinal profiles of the Brazos River below Lake Whitney Dam, indicating net deposition along several miles of the channel. (From Woolley, 1985; illustration from Hobbs, 1987.)

delivered to the bay after reservoir development. Isphording found abrupt changes in sediment regimen in the bay, including increases in sand and clay but a striking decrease in silt, which he traced to construction of a number of dams on rivers that discharged into the bay. Noting that the reservoirs trap sand and silt, he hypothesized that clay is washed over the spillways and continues down the river eventually to settle out in bay waters. He attributed increases in sand in the bay to channel erosion downstream from the reservoirs. The dam nearest the bay was approximately 160 km (100 mi) upstream.

Classification of Rivers by Sediment Load

Rivers can be classified in terms of their sediment load (Schumm, 1968, 1972) as suspended load, mixed load and bed load streams. (Bed load is that part of the load transported along the bed of the channel, and is usually composed of the coarser materials such as sand, gravel, and larger material that slide, roll, or bounce [saltate] along the bottom of the stream; suspended load is that which is held in the water column, or suspended above the channel bed, and is composed primarily of the finer sized materials--mostly silt and clay but also some sand.)

Suspended-load streams transport on average less than 3 percent sand size or larger sediments (bed load), and are generally characterized by relatively narrow, deep and sinuous channels. Bed-load streams transport on average greater than 11 percent sand size and larger material, and are relatively wide, shallow, and straight. Mixed-load streams have sediment characteristics and morphology intermediate between these two end members (Schumm, 1972). The Guadalupe and San Antonio Rivers are examples of suspended load streams, and the Brazos and Colorado Rivers vary from predominantly bed-load streams in their upper reaches to mixed-load streams on the coastal plain (Morton and McGowen, 1979).

Sediment dispersal patterns and modern depositional systems along the Gulf Coast are depicted in figure 40. Deltas fed by modern rivers are significant depositional features along the coast.




Figure 40. Modern depositional systems and sediment dispersal patterns, northwest Gulf of Mexico; by A. J. Scott. (From Fisher and others, 1969.)

DELTAIC SEDIMENTATION

"Of all geologic processes, fluvial sediment transport and deltaic deposition are among the most dynamic..." (Morgan, 1970). A delta as defined by Fisher (1969) is simply "a river-fed depositional system that results in irregular progradation of a shoreline. A complex of delta lobes comprise a delta system." In a broader definition of deltas, Wright (1985) included both subaqueous and subaerial accumulations of river-derived sediments that are deposited at or near the source stream; his definition includes these deposits when reworked by waves, currents, and tides. Although modern deltas may have general similarities in that they are formed from sediment deposited as a stream loses its velocity upon reaching base level (the receiving reservoir), deltas may vary greatly in terms of their composition, size, shape, origin, and sedimentary properties (Morgan, 1970). Deltas range from those that are large-scale depositional units supplied by rivers with high discharge and large sediment loads and that prograde into oceanic waters, to those that are relatively small scale, such as bayhead deltas, that prograde into bays and estuaries (Fisher, 1969).

Delta Development

Among the variables that affect delta formation are: (1) sediment input load such as amount, rate, variation in rate, and suspended/bedload ratio; these factors are affected by climate and extent and nature of the drainage basin, (2) nature of the discharging and reservoir water bodies, especially regarding relative water densities, (3) reservoir energy such as the kind and degree of waves, currents, tides, particularly as related to the amount of sediment input, (4) depth of water into which the delta progrades, (5) nature of substrate of the receiving reservoir, which effects subsidence and storage of prograding sediments, and (6) structural nature of the receiving basin (Fisher, 1969).

The first prerequisite for a delta to occur is the existence of a major river system composed of a drainage basin within which sediments are supplied by erosion from precipitation and runoff, and individual tributaries coalesce to produce a larger trunk stream, which is housed within an alluvial valley that connects to the coastal receiving basin (fig. 41)




Figure 41. Major components of a river system (from Coleman and Wright, 1971).

A general discussion of the history of deltaic studies and delta formation is presented by Morgan (1970) from which the following general information on delta formation was derived.

As sediment-laden water from the river enters an estuary or other receiving basin, density differences cause the river water to spread and gradually mix with the water of the basin. In the mixing zone, suspended sediment, including clays that flocculate, are deposited in subaqueous levees. (Flocculation is a process that occurs as the fresher riverine water enters the more saline estuarine water. Clay particles are characterized by large surface areas that normally have a high negative surface charge. The like charges cause the individual particles to repel each other and remain dispersed. However, when the particles enter water with high concentrations of cations such as marine water, the particles become destabilized and form floccules held together by electrochemical bonding forces. The flocculation process aids sedimentation because the larger floccules settle faster than the individual smaller particles. Clay minerals flocculate and settle at different rates depending on salinities. Kaolinite and illite, under laboratory conditions, flocculate rapidly as salinity increases between 1 and 2 parts per thousand; montmorillonite flocculates over a much wider range of salinities and accordingly settles at a slower rate. Flocculated particles may deflocculate if returned to fresh water. Turbulence can bring more particles together enhancing flocculation, but flocs can also be torn apart by high turbulence that produces shear stresses exceeding the attractive forces of the flocs. This discussion on flocculation was derived principally from Nichols and Biggs, 1985; there are many other papers that consider the process of flocculation in estuaries including Postma, 1967, and Kranck, 1984).

The coarser bed-load is also moved into the receiving basin where, as current velocity decreases, it is deposited at the mouth of the river forming a river mouth bar. Most of the suspended load and some of the bed load is transported beyond the river mouth bar and deposited in an environment referred to as the delta front. Beyond the delta front is an area called the prodelta depositional environment where finer fractions of the suspended load gradually settle to the bottom forming a blanket

of clay and silt. The rates of deposition of sediments around the river mouth are not only dependent on the rate of water mixing and corresponding reductions in water velocities, but also the energy characteristics, such as strength of tides, waves, and currents, of the receiving basin. The interrelated factors that have a bearing on deltaic sedimentation are summarized in table 7.

As deposition at the river mouth continues, the subaqueous levees and river mouth bar eventually become subaerial. Deposition of the river mouth bar causes the channel to divide, or bifurcate, and new subaqueous levees are constructed along the distributaries (fig. 42). The process of subdivision repeats itself in geometric progression (2, 4, 8, etc.) if not opposed by wave erosion and longshore currents (Russell, 1967). The result is a delta, characterized by branching distributaries (fig. 43) that progrades seaward (fig. 44).

Successive flooding and deposition of sediment continues to build up the delta and develop the environments depicted in figures 45 A and B. Coarser sediments (sand) characterize the channels and channel mouth bar deposits, sand and silt characterize the delta front sediments, and clay and silty clay the prodelta deposits. During floods along the distributary channel, coarser sediments are deposited closest to the channel as the river currents diminish rapidly when the flood water leaves the channel. Finer sediments are carried into the flood basins or interdistributary basins where they are deposited in slack water. The natural levees formed along the channel are composed of silt and sandy silt, sediments finer than those in the active channels but coarser than the floodbasin deposits.

Deltas formed at the head of estuaries, for example the Guadalupe and Trinity deltas along the Texas coast, prograde over muds that were deposited in the estuarine environment. As the delta moves seaward, the fine grained prodelta sediments override the bay muds (silt and clay) and they themselves are overridden by the successive deltaic deposits including the delta front sands (fig. 44). However, the dis-

Table 7. Factors influencing deltaic sedimentation (From Morgan, 1970.)

RIVER REGIME (Variations influence sediment load and transport capacity)	Flood stage	Sediment load	Quantity of suspended load and bed load (that is, stream capacity) increases during flood
		Particle size	Particle size of suspended load and bed load (that is, stream competence) increases during flood
	Low river stage	Sediment load	Stream capacity diminishes during low river stage
		Particle size	Stream competence diminishes during low river stage
COASTAL PROCESSES	Wave Energy		High wave energy with resulting turbulence and currents erode, rework, and winnow deltaic sediments
	Tidal range		High tidal range distributes wave energy across an extended littoral zone and creates tidal currents
	Current strength		Strong littoral currents, generated by waves and tides, transport sediment alongshore, offshore, and inshore
STRUCTURAL BEHAVIOR (With respect to sea level datum)	Stable area		Rigid basement precludes delta subsidence and forces deltaic plain to build upward as it progrades
	Subsiding area		Subsidence through structural downwarping coupled with sediment compaction allows delta to construct over- lapping sedimentary lobes as it progrades
	Elevating area		Uplift of land (or lowering of sea level) causes river distribu- taries to cut downward and rework their sedimentary deposits
CLIMATIC FACTORS	Wet area	Hot or warm	High temperature and humidity yield dense vegetative cover, which aids in trapping sediment transported by fluvial or tidal currents
		Cool or cold	Seasonal character of vegetative growth is less effective in sediment trapping; cool winter temperature allows seasonal accumulation of plant debris to form delta plain peats
	Dry area	Hot or warm	Sparse vegetative cover plays minor role in sediment trapping and allows significant aeolian processes in deltaic plain
		Cool or cold	Sparse vegetative cover plays minor role in sediment trappings; winter ice interrupts fluvial processes; seasonal thaws and aeolian processes influence sediment transportation and deposition




Figure 42. Development of mid-channel shoals and branching distributaries. (From Russell, 1967.)




Figure 43. Components of a delta developing in shallow water. (From Gould, 1970.)




Figure 44. Seaward migration of depositional environments in high-constructive deltas, typical of bay-head deltas (Fisher and Brown, 1972). (From Scruton, 1960.)




Figure 45. Development of deltaic facies, including constructional and destructional cycles. (From Frazier, 1967.)

tributary channels may cut down into the bay muds and occupy a lower elevation than the bay floor (Donaldson and others, 1970).

As noted by Morgan (1970, p. 112), deltaic distributaries "prograde when the distributary finds a shorter route of steeper gradient to base level; as a rule this occurs when the river crevasses or breaks through its own levees into an adjacent bay." Typical longitudinal gradients before crevassing may be from 0.10 to 0.15 ft/mi (1.8 to 2.8 cm/km); the cross-levee gradient at the point of the crevasse may exceed 5 ft/mi (100 cm/km) (Morgan, 1970). This steep gradient produces high velocities and turbulence that allows the crevassing stream to scour a channel through its levee and deposit a sediment splay (crevasse splay) in the adjacent bay water (fig. 45 C). If the channel is scoured deeply enough it will continue to supply sediment to the newly formed deposit during the succeeding low-water stages. Morgan (1970) notes that subdeltas of the modern Mississippi River are classic examples of crevasses that have filled shallow bays along principal distributaries.

Development of Deltaic Marshes

As the delta environments aggrade, salt, or brackish to fresh-water marsh plants become established in the intertidal and higher areas. The plants help baffle currents and trap sediments, thus building up the marsh (Redfield, 1972; Gleason and others, 1979; Stumpf, 1983). In intertidal areas, natural levees initially aggrade more rapidly than backmarshes located in interdistributary basins, but as the levees reach higher elevations (along tidal channels), flooding and depositional events are less frequent and aggradation on the back marshes may start to catch up, thus producing a more level surface (Redfield, 1972). This general trend toward decreasing accretion rates and increasing elevations of the marsh surface has not been supported by some investigations (Oenema and DeLaune, 1988).

The higher marsh substrates generally contain more sand and silt than the lower marshes, which are typically characterized by mud (Kanes, 1970; Frey and Basan, 1985; fig. 46). In addition to being topographically above the lower marsh, the higher marsh is older and is influenced more by terrestrial conditions (Frey and Basan, 1985). The lower marsh is often intertidal and more frequently inundated by waters carrying suspended sediments from the bay-estuary-lagoon system.

Marsh Sediments

Frey and Basan (1985) noted that very few marshes along the Gulf coast have been studied in detail geologically, and suggest that the marsh sediments (except for the Mississippi delta complex) seem to be similar to those of the southeastern Atlantic coast, which were described as dominantly inorganic with insignificant amounts of peat. Suggested reasons for the absence of thick peat deposits are (1) tidal flushing, (2) rapid degradation of plant material by intense biologic activity, and (3) extremely slow rates of coastal warping or submergence (Frey and Basan, 1985). Comparison of total organics in sediments in Texas brackish marshes and salt marshes (White and Calnan, in preparation) indicate lower concentrations than reported by Frey and Basan (1985) for Georgia salt marshes.

Inorganic sediments of a representative southern Coastal Plain salt marsh (near Sapelo Island) contained approximately constant proportions of silt and clay, where maximum amounts were 60 % and 55 %, respectively; sand is uniformly low in the low marsh with sand and muddy sand predominant in the high marsh (Frey and Basan, 1985) (fig. 46).

In a study of a delta lobe on the eastern half of the Colorado River delta along the Texas coast, Kanes (1970) reported that the low-marsh sediments were characteristically finer grained than the higher marsh. He suggested that the sediments on the higher marsh were derived from the river during floods, while the low-marsh




Figure 46. Relative percentages of sand, silt, and clay in salt marsh sediments, Sapelo Island, Georgia. (From Frey and Basan, 1985, after Edwards and Frey, 1977.)

sediments were derived from turbid bay waters. In a limited number of samples collected from salt marshes on the western half of the Colorado River delta, White and Calnan (in preparation) found clay (ranging up to 80 %) to be significantly more abundant than silt (maximum of about 30 %) in the low marshes, while in a levee marsh, silt (at about 50 %) was slightly more abundant than clay (about 40 %). Sand was not a significant component of the marshes sampled except on nearby Matagorda Peninsula where it was predominant (ranging up to almost 90 %). Sediment samples in a brackish marsh system on the Trinity River delta along the upper Texas coast were more variable with maximum concentrations of clay and silt at about 80% and 20 % in low (back) marshes; high marshes along levees had higher silt content ranging to a maximum of near 55 %, while maximum content for clay was near 40%. Marshes with sandier substrates were located on delta substrates undergoing active progradation near the bay margin. At one site near a distributary channel, the concentrations of sand, silt, and clay were 67, 18, and 15 %, respectively (White and Calnan, in preparation). The marsh at this latter site was only partly vegetated; sediment samples contained less than 0.5 % organic carbon.

Marsh Degradation

In deltaic areas where fluvial sediments are abundantly supplied to the marshes during flood events, the marshes may follow a geologic succession from low marsh to high marsh to uplands as levee deposits aggrade through successive depositional events. When active distributaries are abandoned and the delta progrades into new areas, the fluvial-sediment supply to the older delta lobe is cut off or is reduced. Marshes may continue to receive sediments through tidal inundation, but will not remain emergent unless the sediment supply is enough to offset destructive processes. Processes that tend to degrade a delta such as waves and currents and subsidence begin to dominate and the abandoned delta lobe undergoes deterioration through erosion and submergence. Upland levee deposits may revert to marshes as the delta lobe subsides (fig. 47).




Figure 47. Development and deterioration of deltaic marshes. (From Bernard and LeBlanc, 1965, modified from Fisk, 1960.)

Life Cycle of a Subdelta in a Delta System

The extensive loss of wetlands on the Mississippi River Delta (Gagliano and others, 1981) has made it the center of deltaic wetlands research along the Gulf coast. Wells and Coleman (1987) noted that although the causes of wetland loss are complex, there is agreement among most scientists that a combination of natural processes such as subsidence, sea-level rise, and changing depositional sites, as well as human-induced causes, such as artificial canals and levees, and sediment diversion are major contributors. Wells and Coleman (1987) concluded that the life history of a subdelta, which undergoes a constructional and destructional phase (Scruton, 1960), could be characterized as follows: (1) subdeltas follow a natural cycle of growth and deterioration that lasts for a period of about 115 to 175 years; (2) growth rate is regulated by depth of the receiving basin, sediment discharge through the crevassing channel, amount of sediment deposited in the receiving basin, and the efficiency of the channel network in the subdelta to deliver the sediment; (3) there is an initial period of slow infilling (10-30 years) during which a subaqueous channel system is being constructed; this period is followed by rapid subaerial growth as channels extend farther into the receiving basin by elongation and bifurcation, until a peak is reached as the channel network becomes too complex (about 25 channels) to efficiently deliver sediments; (4) rapid subsidence is characteristic of the deteriorating phase, which allows for continuous infilling during subaerial deterioration; reversion to open water initially occurs in the older parts of the subdelta, and expands from there. The Mississippi delta has a complex evolutionary history (fig. 48).

A significant process that leads to the destruction of abandoned delta environments is subsidence. Morgan (1967) noted that in addition to regional downwarping, subsidence can result from the effect of overloading and attendant compaction and water loss in underlying sediments. When sedimentation ceases




Figure 48. Evolutionary sequence of the Mississippi River Delta distributaries. (From Morgan, 1977, in Boesch and others, 1983.)

following abandonment of a delta lobe, subsidence continues. The effects are apparent, first, in interdistributary marshes, which begin to be replaced by open water (fig. 49). Because the distributary levees are more massive than other delta environments, they subside more rapidly and drag down adjacent marshes forming elongate ponds that parallel the levees (Morgan, 1967). As subsidence continues, the action of waves and currents in the expanding ponds accelerates the process through erosion, and the smaller ponds coalesce. The marshes ultimately are converted back to marine or estuarine open-water habitats.

The deltas in the bay-estuary-lagoon system along the Texas coast have developed in a significantly different setting than the much larger Mississippi River Delta, which has prograded over thick muds on the continental shelf. As noted by Winker (1979), deltas formed by the Guadalupe River (Donaldson 1970) and Trinity River (McEwen, 1969) are examples of relatively thin deltas that have prograded into shallow, protected estuarine systems. Still, these deltas have gone through stages similar to the Mississippi River Delta (fig. 48). For example, the growth and development of the Guadalupe River delta at the head of San Antonio Bay has had a history of delta lobe construction, abandonment, and subsidence (fig. 50). The reversion of marsh to open water occurs in the abandoned parts of the delta as subsidence outpaces sediment supply.

Marsh Maturation and Aggradation

After marsh vegetation begins to colonize deltaic deposits as described in the preceding discussions, the marsh substrate may continue to aggrade (accrete vertically) as sediments are delivered with each inundation. The rate of aggradation, or vertical accretion, depends on many factors such as (1) frequency, depth, and period of inundation, (2) amount of sediment transported into the marsh with each inundation, (3) type of inundation, whether from river flooding or estuarine tides, (4)




Figure 49. Effect of subsidence from sediment compaction on distributary and interdistributary areas. (From Morgan, 1967.)




Figure 50. Growth and development of the Guadalupe delta at the head of San Antonio Bay. The most recent development has occurred in Mission Lake at Traylor Cut subdelta located at E in illustration D. (From Donaldson and others, 1970.)

location and elevation of the marsh, whether streamside (levee) or backmarsh, and (5) biological factors such as density and type of vegetation.

In his classic study of Barnstable Marsh in Massachusetts, Redfield (1972, p. 224) reported that marsh aggradation rates were variable and depended "on the elevation which the intertidal marsh had reached, the availability of waterborne sediment, and the distance from open water." Rates of vertical accretion were highest in the initial stage of development of an intertidal marsh as exemplified by a rate of 5.1 cm/yr. The rate of accretion diminished as the surface rose, producing an average rate of about 1.8 cm/yr. In the high marsh at greater distances from open water, long-term (approximately 1,000 yrs) rates average about 0.21 to 0.15 cm/yr. Redfield concluded that the vertical accretion and the transformation of intertidal marshes to high marshes are dependent on availability of sediment. The accretion rate is presumably more rapid along the margins of the marsh where levees form, and less rapid in backmarsh areas. As the marsh surface becomes more extensive and reaches higher elevations, it becomes more inaccessible and the rate of accretion slows (Redfield, 1972).

In a study of tidal salt marshes in England, Pethick (1981) concluded that there is a positive statistical relationship between marsh elevation/age and vertical accretion rates. His model showed that young marshes (10-year-old marshes) had accretion rates of up to 1.7 cm/yr while older marshes (> 500 yrs old) had rates as low as 0.002 cm/yr. The concept of the model is that younger marshes are lower in elevation, which amplifies the depth and frequency of inundations compared to older marshes. Older marshes have reached higher elevations, which decreases the frequency and period of inundation resulting in less sediment deposition and lower rates of vertical accretion through time. Pethick reported that the age/height relationship defines an asymptotic curve where the asymptote is apparently controlled by the frequency of tidal maxima, which lies about 80 cm below the level of the highest

spring tides in the North Norfolk, England, marsh that he investigated.

Stages of development, or maturation, of marshes have been recognized by many researchers (Fisk, 1960; Gould and Morgan, 1962; Chapman, 1974; and Frey and Basan, 1985). As mentioned by Frey and Basan (1985), the maturation process is a reflection of floral and faunal succession, as well as size, position, and differentiation of the marsh in the coastal system. Geologically, succession occurs through progradation and the resulting lateral and vertical displacement of low marshes by high marshes (Frey and Basan, 1985). The elevation and age of a marsh may follow a chronology like that described by Pethick's model discussed in the preceding paragraph, but actual ages of low (young) and high (old) marshes may be different from those detailed by Pethick. Progradation and development of salt marshes at the mouth of the Colorado River along the central Texas coast, for instance, occurred primarily after 1929 when a log jam was removed from the river (the delta prograded approximately 6.4 km [4 mi] in about 6 years (Wadsworth, 1966). High levee marshes and low interdistributary marshes, although still being modified, have developed during this relatively brief period. Frey and Basan (1985, p. 238) pointed out that while a scale of relative maturation, from youthful to mature to old, can be established for marshes, the emphasis "clearly is upon physiographic "stage" and not "finite" age". The Colorado River delta marsh can be characterized as youthful by the fact that low-marsh environments make up most of the total area. A marsh with a large terrestrial sediment supply and rapid deposition may generally favor high-marsh development, whereas the stable phase of a marsh with a low terrigenous sediment supply, pronounced tidal range and tidal sedimentation, and growth restricted to bay or estuarine margins, may be submature to supermature (Frey and Basan, 1985).

Marsh sedimentation and rates of marsh aggradation have been investigated by many researchers including Van Straatan and Kuenen, 1958; Ranwell, 1964; Schou,

1967; Pestrong, 1972; Redfield, 1972; Armentano and Woodwell, 1975; Flessa and others, 1977; Harrison and Bloom, 1977; Richard, 1978; DeLaune and Patrick, 1980; Letzsch and Frey, 1980; Nixon, 1980; Pethick, 1981; Hatton and others, 1982; DeLaune and others, 1983; Hatton and others, 1983; Stumpf, 1983; Baumann and others, 1984; Boesch and others, 1984; Frey and Basan, 1985; Smith and Frey, 1985; Stevenson and others, 1985; Stevenson and others, 1986; Cahoon and others, 1987; Stevenson and others, 1988, and Oenema and DeLaune, 1988. Studies of accretion rates of Gulf Coast marshes have been conducted principally in Louisiana marshes where loss of wetlands has occurred at the alarming rate of more than $100 \text{ km}^2/\text{yr}$ (Gagliano and others, 1981). There is a lack of information on the rate of accretion in Texas marshes, although a loss in marsh area has been documented in the Galveston-Houston area (Johnston and Ader, 1985; White and others, 1985) and Beaumont-Port Arthur area (Gosselink and others, 1979; White and others, 1987).

Marsh aggradation is an important process in which inorganic and organic sediments that are deposited on the marsh surface provide nutrients for plant growth and help keep the marsh emergent (DeLaune and others, 1978; 1981; DeLaune and Patrick, 1980). Some of the highest rates of marsh aggradation have been documented along the Gulf Coast in Louisiana (DeLaune and others, 1978; Boesch and others, 1983). The rates are considerably higher than along the east coast (table 8), and presumably are related to the high rates of subsidence and relative sea-level rise in coastal Louisiana (DeLaune and others, 1978; Hatton and others, 1983). It has been proposed by Rusnak (1967) that the accumulation of sediments in estuaries must be equal to sea-level rise. Although the rate of aggradation in marshes may be decreased by compaction, it may be increased by subsidence and rise in sea level (Letzsch and Frey, 1980). The higher rates in areas of more rapid subsidence are probably due to more frequent inundations and sediment deposition; that is, there is a natural feedback loop (comparable to Pethick's, 1981, model

Table 8. Marsh aggradation (vertical accretion) rates measured in coastal Louisiana and along the U. S. Atlantic coast (from Boesch and others, 1983).

Location	Marsh type	Marsh accretion rate (mm/yr)	Mean sea-level rise (mm/yr)	Source
Louisiana Deltaic Plain	Freshwater streamside backmarsh	10.6 6.5	11.0	Hatton and others (1983)
	Intermediate (<u>Spartina patens</u>) streamside backmarsh	13.5 6.4		" "
	Brackish (<u>S. patens</u>) streamside backmarsh	14.0 5.9		" "
	Saline (<u>S. alterniflora</u>) streamside backmarsh	13.5 7.5	13.0	DeLaune and others (1978); Baumann (1980)
Chenier Plain	Salt-brackish (<u>S. patens</u>)	7.0	12.0	Baumann and DeLaune (1982)
Georgia	<u>S. alterniflora</u>	3-5		Summarized by Hatton and others (1983)
Delaware	<u>S. alterniflora</u>	5.0-6.3	3.8	" "
New York	<u>S. alterniflora</u>	2.5-6.3	2.9	" "
Conn.	<u>S. alterniflora</u> <u>S. patens</u>	8-10 2-5	2.5	" "
Mass.	<u>S. alterniflora</u>	2-18	3.4	Redfield (1972)

describing the relationship between marsh accretion and its height and age) in which a submerging (and thus topographically low) marsh attempts to retain an equilibrium condition by trapping sufficient sediments to remain emergent. If there is insufficient sediment, however, the rate of aggradation will not keep pace with the relative rise in sea level and the marsh will eventually be replaced by open water. Rates of marsh aggradation in many areas of coastal Louisiana have apparently not kept pace with rates of submergence (DeLaune and others, 1978; DeLaune and others, 1983; Baumann and others, 1984).

Average aggradation rates (determined from analysis of ^{137}Cs in cores) in Spartina alterniflora salt marshes in Barataria Basin in Louisiana vary from 1.35 cm/yr to 0.75 cm/yr in levee and backmarsh areas, respectively (DeLaune and others, 1978). Measurements of aggradation rates in fresh, intermediate, brackish, and salt marshes in Barataria Basin (also using ^{137}Cs) provided similar results of 1.35 cm/yr to 0.75 cm/yr in levee and backmarsh environments (Hatton and others, 1983). Rates of aggradation in brackish marshes on the Chenier Plain near Calcasieu Lake to the east of Sabine Lake averaged 0.8 cm/yr, as determined by ^{137}Cs and artificial marker horizons (DeLaune and others, 1983). The rate of relative sea-level rise (based on tide-gauge records) in these areas of Louisiana ranges from about 1.1 to 1.3 cm/yr (DeLaune and others, 1978; DeLaune and others, 1983; and Hatton and others, 1983). A comparison of marsh accretion rates with mean sea-level rise for the Louisiana Deltaic Plain and Chenier Plain, as well as for marshes along the Atlantic Coast was summarized by Boesch and others (1983) in table 8. While the accretion rates of levee marshes in the Deltaic Plain are near the rate of relative sea-level rise, accretion rates in inland, or backmarshes, are not keeping pace and the marsh sites are deteriorating into small open water areas (DeLaune and others, 1978; Hatton and others, 1983). In the Chenier Plain, DeLaune and others (1983), conclude that "Clearly, an accretion rate of 0.8 cm/yr is not sufficient to maintain the elevation of the marsh in an area that is submerging at 1.2 cm/yr".

Baumann and others (1984) investigated marshes in the Mississippi Deltaic Plain and found that although accretion rates in marshes of Barataria Bay ranged from 9 to 15 mm/yr and marshes in Fourleague Bay had accretion rates of from 6 to 13 mm/yr in inland and streamside marshes respectively, the marshes in Barataria Bay were deteriorating whereas marshes in Fourleague Bay were more stable. The authors attributed the closer balance between marsh aggradation and relative sea level rise in Fourleague Bay marshes to fluvial sediments delivered by the Atchafalaya River. Annual river flooding was the major depositional process in Fourleague Bay marshes. Annual flood cycles contributed 91 and 69 percent of the streamside and inland sediments, respectively, during 1981 and 1982 (Baumann and others, 1984). Basinal sedimentary processes at work in Barataria Bay, where deteriorating marshes received most of their sediment during storm events, were insufficient to counter the effects of relative sea-level rise. Baumann and others (1984) further concluded that major accretion in Barataria Bay effectively ended with the damming of Bayou Laforche in 1904, and with the construction of an artificial levee system along the Mississippi River after the 1927 flood. Sediments delivered to the Barataria Basin marshes were principally a result of hurricanes, tropical storms, and winter storms (fig. 51). Hurricanes and winter fronts, the latter occurring at an average frequency of 5.7 per winter month, are apparently major depositional and erosional agents in both bay areas. Baumann and others (1984, p. 224) stated that "The transformation from marsh to open bay cannot be halted unless there is a reintroduction of riverine sediment".

The delivery of sediments by river flooding is an important process, especially in areas characterized by low-tidal regimes. As pointed out by Frey and Basan (1985), the difference, in terms of hydraulics, between river and tidal flooding is one of duration; there is a longer slack-water period during river flooding, which allows more time for suspended sediments to settle out.




Figure 51. Seasonal variations in aggradation rates of salt-water marshes in Barataria Basin, Louisiana. Mean seasonal aggradation for 1975-1978 (A) indicate rates were highest in winter, compared with mean seasonal aggradation for 1975-1979 (B) when rates were elevated by a hurricane and tropical storm that made landfall during the summer of 1979. (From Baumann, 1980, modified by Gosselink, 1984.)

Van Heerden and others (1981), using cores and bathymetric charts to study the evolution of the Atchafalaya Delta, found sedimentation rates of levee environments to be associated with annual flood events. They identified deposits with high sedimentation rates (> 0.3 m per annual flood), medium sedimentation rates (0.15 to 0.30 m per annual flood), and low sedimentation rates (< 0.15 m per annual flood).

In an investigation of the emerging Atchafalaya Delta, Louisiana, DeLaune and others (1987) reported that marsh accretion rates were as great as 1.4 cm/yr. The higher rates were in bay bottoms and marshes closest to the active delta. Fluvial sediment input causes the delta to prograde and also provides the source of marsh nutrients supporting vegetation growth, which in turn provides organics for vertical marsh accretion. Marsh areas updrift, away from the prograding delta, had a smaller mineral sediment input as well as organic input, and were accreting at a slower rate; the authors concluded that these areas will likely continue to deteriorate.

In the Barataria Basin, which receives insignificant riverborne sediment, there is a landward decrease in inorganic sediments as one progresses from the Gulfward lying salt marshes to the landward lying fresh marshes (Hatton and others, 1982). The inorganic sediments, which increase in a seaward direction in response to hydraulic energy, are apparently locally derived from erosion of adjacent marshes. The organic matter accumulating in the marsh shows an inverse relationship to the inorganic sediment; that is, the accretionary role of organic matter increases in a landward direction, from saline- to fresh-water marshes (Hatton and others, 1982). These inorganic-organic relationships are reversed near active deltas where rivers and distributary channels are sources of inorganic minerals (fig. 52).

Stevenson and others (1985) reviewed 15 areas (Atlantic and Gulf coasts) in a comparison of rates of marsh aggradation with rates of relative sea level rise. They found that in at least four of the sites, three of which are along the Gulf coast, marsh aggradation rates were not keeping pace with relative sea level rise (fig. 53).




Figure 52. Percentages of inorganic (mineral) sediments in marsh soils in Mississippi delta basins arranged in order of increasing age. (From Gosselink, 1984, based on data from Chabreck, 1972.)




Figure 53. Relationship between rates of marsh aggradation and relative sea-level rise between 1940 and 1980. Aggradation rates were determined from ²¹⁰Pb (solid circles) and ¹³⁷Cs (open circles) measurements, and historical measurements (asterisk for Barnstable marsh). Abbreviations are: BB (Barataria Bay); BN (Barnstable marsh); BW (Blackwater marsh); FL (Fourleague Bay); FR (Farm River); FX (Flax Pond); FP (Fresh Pond); LC (Lake Calcasieu); LW (Lewes); PI (Prudence Is); NI (North Inlet); NN (Nanticoke River); NR (North River); SI (Sapelo Island); and SR (Savannah River). (From Stevenson and others, 1986.)

Rates of relative (apparent) sea-level rise at the Gulf coast sites, however, are more than two-times the rates at the other sites reviewed (fig. 53). The authors found a strong correlation between mean tidal range and accretionary balance (the four sites that were not keeping pace with relative sea level rise are in microtidal areas), but concluded that more data are needed from areas of low tidal energy to confirm the relationship. Nevertheless, Stevenson and others (1985) postulated that where tides are weak and irregular and sediment inputs are low, or reduced relative to past levels, marshes have difficulty in keeping pace with relative sea-level rise. They further suggest that hurricanes and storms, which can sporadically add sediments to marshes, may play a critical role in the sediment budget in areas with low tides, which is in agreement with Stumpf's (1983) conclusion that storms control sediment supply on microtidal marshes, and that sedimentation depends directly on storm frequency and sediment availability.

Nixon (1980) presented data on marsh accretion rates and sea-level rise to support his conclusion that salt marshes are sinks of suspended sediments "at least on the time scale of years to centuries". Using Chesapeake Bay, he concluded that over 15 percent of the annual sediment input from the Susquehanna River and other sources was deposited on the marshes. A review of the Chesapeake Bay sediment budget by Stevenson and others (1988) indicated Nixon's estimate was high and the marshes only trapped 5 to 11 percent of the total sediment input. Providing other evidence, Stevenson and others (1988) concluded that "although the concept that marshes act as major sediment sinks may be accurate over the last few millenia, tidal transport studies suggest considerable variability with most marshes presently exporting material on an annual basis". These authors suggest that estuarine sedimentation occurs principally in subtidal flats, which are below the limit of emergent marsh vegetation. Stevenson and others (1988) further concluded that (with respect to north Atlantic, south Atlantic, and Gulf coast marsh systems) southern marshes

are apparently more susceptible to erosion and export of materials than northern marshes because of (1) differences in tidal dynamics, (2) seasonal changes in sea levels, and (3) higher temperatures. Another critical factor in the erosion and export processes, they hypothesized, is the reduction of the terrigenous sediment supply from the rivers along the southern U.S., which has led to sediment starvation and undernourishment of the coastal wetland systems over the last half century (fig. 54).

Biodeposition

Biodeposition is an important part of sedimentation in marshes. Smith and Frey (1985) have proposed that suspension-feeding invertebrates, which actively filter suspended particles from flood waters and bind and deposit the particles on the marsh surface, may be responsible for much of the net sedimentation in the marsh. Feces and pseudofeces of these invertebrates are important in stabilizing the sediment surface by forming large aggregates of silt and clay-size particles (Haven and Morales-Alamo, 1968), providing the marsh with a means of retaining nitrogen, phosphorus and trace elements, and are capable of recycling organic materials within the detrital food chain (Kraeuter, 1976). The manipulation of sediment during ingestion and in the digestive tract of the organisms will change the apparent grain-size of the sediment, and while passing through the digestive tract, the processes by which organics are removed may alter the organic and inorganic chemistry of the sediments (Carney, 1981).

Many suspension-feeding invertebrates, such as oysters, barnacles, tunicates and copepods, ingest large quantities of small particles in the 1-5- micron range, and after passage through the digestive tract, the particles are voided into the water as fecal pellets which range in length from 50 to 3,000 microns (Haven and Morales-Alamo, 1972). The upper size limit of particles ingested by suspension feeders is generally smaller than that of deposit feeders (Rhoads, 1974). In many molluscan




Figure 54. Comparison of sediment delivery to near-shore areas along the eastern U.S. and Mississippi Delta coasts between the early-1900's and 1970. Along the Gulf Coast the scale is in 10^9 tons/yr. and along the Atlantic Coast 10^6 tons/yr. (From Stevenson and others, 1988.)

species, some particulate matter is rejected before ingestion and ejected from the shell cavity in a loosely compacted mass termed pseudofeces (Haven and Morales-Alamo, 1972). Fecal pellets settle to the bottom at a faster rate than that of the particles originally in suspension, and they may be resuspended or mixed into bottom deposits by deposit feeders. Pellets and their fine component particles may accumulate in bottom areas where they would not normally settle out because of differences in hydrodynamic properties between the pellets and particles (Haven and Morales-Alamo, 1968). Feces and pseudofeces that settle to the bottom are termed biodeposits. The entire complex process, involving many groups of animals and physical and chemical factors, is termed biodeposition (fig. 55) (Haven and Morales-Alamo, 1966).

Suspension feeders and deposit feeders may produce large quantities of biodeposits. For example, Rhoads (1974) reports that the top centimeter of mud in Buzzards Bay, Massachusetts is almost entirely pelletal because of reworking of the mud by the polychaete Clymenella torquata and the bivalves Yoldia limatula and Nucula annulata. Laboratory studies indicate that oysters on 0.405 hectare of an estuarine bottom may produce up to 981 kg of feces and pseudofeces (Haven and Morales-Alamo, 1966). Lund (1957) calculated that if oysters covered an acre of bottom, they would deposit 8.36 tons of fecal material (dry weight) or 6 tons of dry mineral matter in 11 days. Rates of biodeposition of mud by some suspension and deposit feeders found on the Texas coast are shown in table 9.

Suspension-feeding mussels, such as Geukensia demissa, may ultimately enhance the productivity of a salt marsh by filtering particulate nitrogen and depositing it as feces and pseudofeces in the marsh sediment. The major role of the mussels in nitrogen flow in a marsh is to increase retention of nitrogen within the marsh by filtering particulate nitrogen from suspension (Jordan and Valiela, 1982). Jordan and Valiela (1982) report that the entire population of Geukensia demissa in a New




Figure 55. Theoretical biodeposition cycle in an estuary (from Haven and Morales-Alamo, 1966).

Table 9. Rates of biodeposition of mud by some suspension and deposit feeders on the Texas coast. Modified from Rhoads, 1974; Kraeuter, 1976.)

<u>Species</u>	<u>Deposition</u> <u>(mg dry weight of feces per animal per day)</u>
<u>Littorina irrorata</u> (gastropod)	0.4
<u>Crassostrea virginica</u> (bivalve)	18.0
<u>Geukensia demissa</u> (bivalve)	13.0
<u>Balanus eburneus</u> (barnacle)	0.4
<u>Molgula manhattenensis</u> (tunicate)	64.0
<u>Clymenella torquata</u> (polychaete)	90.0
<u>Pectinaria gouldii</u> (polychaete)	164.0

England salt marsh is capable of filtering a volume of water in excess of the tidal volume of the marsh during each tidal cycle in the summer. Yearly, the mussels filter 1.8 times the particulate nitrogen exported from the marsh by tidal flushing (Jordan and Valiela, 1982). Half of this nitrogen is absorbed by the mussels and half is deposited as feces and pseudofeces. Since nitrogen limits productivity, increased retention of nitrogen may increase productivity of the marsh.

Deposit feeders probably play a more quantitatively significant role in 'pelletizing' a muddy sea floor than suspension feeders (Rhoads, 1974). The deposit-feeding polychaetes Clymenella torquata and Pectinaria gouldii can rework mud at rates of 96 to 400 ml wet mud/individual/yr (Rhoads, 1974). The rates become more impressive when the individual rates are multiplied by the standing crops found in some temperate and boreal bays which commonly range from 100 to 10,000 individuals/m² (Rhoads, 1974).

Biodepositional rates vary seasonally. Temperate estuaries are subjected to hot summers and cold winters requiring organisms to undergo seasonal metabolic changes which directly affect the organisms' feeding and filtration rates (Biggs and Howell, 1984). During periods of extreme heat or cold, organisms will reduce their filtering rates to a minimum or totally "shut down" (Biggs and Howell, 1984). Smith and Frey (1985) studied biodeposition by the bivalve Geukensia demissa in a salt marsh and found that the following variables influenced seasonal biodepositional rates: (1) mussel density; (2) length of tidal inundation (feeding time or time available for biodeposition), and (3) individual mussel biodepositional rates (g/mussel/hr). Individual mussel biodepositional rates are in turn affected by mussel size, water temperature, and amount of material suspended in water (Smith and Frey, 1985).

Bioturbation or sediment reworking in the form of burrowing, tube building, and the production of sediment-binding exudates, along with biodeposition, are known to influence the fate and transport of sediment (Schaffner and others, 1987). These




Figure 56. Comparison of size and shape of the Trinity, Guadalupe, and Colorado River deltas with the Atchafalaya River delta and crevasse-splay deposits of the modern Mississippi River. (From Cunningham, 1981.)

processes are, in turn, affected by sediment accumulation rates. Areas of rapid deposition (>3 cm/yr) exhibit little evidence of bioturbation, as do areas where erosion dominates (Schaffner and others, 1987). Areas with low sediment accumulation rates (0.5-3.0 cm/yr) exhibit the highest levels of mixing.

Texas Deltaic Marshes--Development and Current State

This literature synthesis deals principally with fluvial sediments and their role in marsh development and maintenance; thus, the following discussion focuses on deltaic areas, which are the loci of fluvial sediment deposition, and the sites of extensive marshes. As previously discussed, several rivers in Texas have constructed deltas in bay-estuary-lagoon areas (fig. 1). Deltaic areas studied most extensively are those at the mouths of the Colorado (Wadsworth, 1941, 1966; Kanes, 1965, 1970; Manka and Steinmetz, 1971), Guadalupe (Donaldson and others, 1970), and Trinity Rivers (McEwen, 1969). Comparisons of the size and shape of these deltas with others is shown in figure 56. Aggradation rates of marshes that have developed in deltaic areas have apparently not been previously investigated in Texas, but are currently under investigation in two areas by White and Calnan (in preparation).

Colorado River Delta

The Colorado River delta is the site of extensive salt-water marshes (McGowen and Brewton, 1975; McGowen and others, 1976a; Benton and others, 1977; Adams and Tingley, 1977; Ward and others 1980; van Beek and others, 1980; White and others, 1985a, 1988). In 1976, the delta was approximately 3,310 hectares (8,175 acres) in size, and about 80 percent was composed of marshes, shallow water bodies and channels, and intertidal flats; on the remaining part (20 percent) of the delta were spoil deposits, natural levees, man-made structures, and shell ridges (van Beek and others, 1980). The historical development of the Colorado River delta has been discussed by several authors including Wadsworth (1941; 1966), Kanes (1965; 1970),


Bouma and Bryant (1969), Manka and Steinmetz (1971), McGowen and Brewton (1975), USACE (1977; 1981), Ward and others (1980), and van Beek and others, (1980).

The Colorado River delta has had a relative brief history of development after removal of a log raft that had trapped large amounts of sediments along its lower reaches. Removal of the raft in 1929 led to rapid progradation of the delta across the eastern arm of Matagorda Bay (approximately 6 km) between 1929 and 1935 (Wadsworth, 1966). In 1936, a channel was dredged across Matagorda Peninsula allowing the river to discharge directly into the Gulf of Mexico. The successive stages of growth of the delta are depicted in figure 57. Between 1929 and 1941, the average rate of growth of the delta was approximately 500 acres a year (Wadsworth, 1966; fig. 58).


Kanes (1965; 1970) conducted a detailed geological study of different environments of deposition in the northeast lobe (Egret Island) of the Colorado River delta. His investigation, which was based on almost 40 cores, revealed a platform of deltaic sediments 2.4 to 3 m (8 to 10 ft) thick. Depositional environments he delineated are delta-plain clay and silt (0.3 to 1.5 m [1 to 5 ft] thick), delta-front sand (0.6 to 2.4 m [2 to 8 ft] thick), prodelta silty clays (0.3 to 1.5 m [1 to 5 ft] thick), and bay silty clays and clayey silts (10 to 14 ft thick). Depositional environments are similar to those of other deltas (fig. 45). The delta plain is subdivided into low and high marshes, beach ridge, and upper and lower distributary channel. The high marshes included natural levees.

Kanes (1970) found that the early prodelta deposits of the river are fine-grained laminated sediments that lacked sand and silt. He attributed the lack of a coarser fraction to a logjam along the river, which trapped the sand and silt upstream. Removal of the logjam led to the development of a delta-front sheet sand during an initial phase of deposition, followed by a second phase in which sand was deposited


(A)



(B)



(C)



(D)




Figure 57. Historical development of the Colorado River delta between 1908 and 1941. Bathymetry in (C) dated 1934. (From Manka and Steinmetz, 1971; growth of the delta after Wadsworth, 1966; bathymetry after Kanes, 1965; salinity after Galtsoff, 1930.)




Figure 58. Increase in area of the Colorado River delta for the period 1908-1953. Postulated flood years are marked by "P". (From Wadsworth, 1966.)

in prograding distributary-mouth bars forming bar fingers or a digitate outline (Kanes suggested that the pattern associated with this second phase of sand deposition may have been due to diversion of the river through a small manmade channel). Finer-grained sediments were deposited in interdistributary areas. High and low marshes of delta-plain environments formed on the subaerial delta lobe. Low marsh sediments were commonly finer grained and contained more laminations than the higher marsh sediment, which suggested deposition of fine particles from turbid bay waters in low marshes (Kanes, 1970).

Manka and Steinmetz (1971) investigated the depositional history of the delta's southeast lobe (gulfward of the lobe studied by Kanes). A three-dimensional reconstruction of the depositional history was based on 21 cores (the locations of 11 cores are shown in figure 59). Cross sections of this part of the Colorado River delta indicate thickness of approximately 2.1 m (7 ft) (fig. 60). Manka and Steinmetz agreed with Kanes in suggesting that the constructional phase of deltaic sedimentation (Scruton, 1960) ended in 1941 with the closing of all distributaries; they further suggested that the closing of the distributaries corresponded with the construction of the farm road along the eastern bank of the river channel. They concluded that the entire delta had entered a destructional phase (Scruton, 1960) in 1941. Manka and Steinmetz (1971, p. 322) "felt that if the Colorado River discharge channel had not been dredged into the Gulf of Mexico, and if flow and sediment load were not now regulated by dams upstream, most, if not all, of the eastern half of Matagorda Bay now would be filled with deltaic sediments."

The progradation of the Colorado River produced some exceptionally high rates of bay filling near the river's mouth. The delta lobe that formed between 1929 and 1930 prograded about halfway across the bay (fig. 57 A and 57 B). The exact depths of the bay in 1929 are not known; the bathymetry shown in figure 57 A is for 1859-1872 (Kanes, 1970). Kanes noted that bathymetric surveys taken in 1904




Figure 59. Deterioration or destruction of the southeast lobe of the Colorado River delta plain, as determined by comparing shoreline positions on aerial photographs taken in 1943, 1952, and 1965. Transgressive units indicate retreat of the shoreline as interdistributary bays replace marshes. Hatched lines indicate areas where interdistributary bays are developing. Solid circles are the locations of selected cores taken by Manka and Steinmetz. (Modified from Manka and Steinmetz, 1971.)




Figure 60. Cross sections of the southeast lobe of the Colorado River delta. Lines of sections are shown in figure 59 (From Manka and Steinmetz, 1971.)

and 1906 were questionable, but the surveys suggest that the bottom was shallower than shown by the 1800's survey. Assuming the depths were about 0.6 to 1 m (2 to 3 ft)¹, this indicates a sedimentation rate of about 30 to 50 cm/yr (deposition of 0.6 to 1 m of sediment in two years). This rate is similar to the sedimentation rate of the prograding Mississippi River of 300 m per 1000 yrs (Shepard, 1953; Scruton, 1960; Rusnak, 1967). However, estimates of rates of sedimentation of the Colorado River delta are considerably less if bathymetric data presented by van Beek and others (1980) are used. Changes in bathymetry in the area of the Colorado River delta between 1921 and 1935 reported by van Beek and others indicate a maximum of 1.2 m (4 ft) of sediment accumulation. This yields an aggradation rate of approximately 8 to 9 cm/yr for this 14 yr period. If, however, we assume that most of this change in depth occurred between 1929 and 1935, then the rate increases to 20 cm/yr (1.2m in 6 yrs). This assumption may be incorrect, however, because historical data indicates the growth of the delta had begun as early as 1921 (van Beek and others, 1970).

Sedimentation rates can also be roughly calculated from analysis of time-sediment intervals defined in the southeastern lobe of the Colorado River delta by Manka and Steinmetz (1971, fig. 60). Assuming that sequences A, B, and C (fig. 60) were deposited between 1933 and 1936 (the postulated year in which interval D began to accumulate), a sedimentation rate of about 40 cm/yr is derived (approximate total thickness of these units is about 1.2 m [4 ft] in core 5). This is similar to the 30

1 Galtsoff (1931), who conducted an oyster survey in this area in 1926, reported that depths decreased toward the head of the bay, and that channel depths were less than 0.9 m (3 ft)

to 50 cm/yr rate calculated for the northeastern lobe over a two-year period (see preceding paragraph). Manka and Steinmetz assumed that interval C was deposited during the 1935 flood in which river discharge reached 177,000 cfs (maximum recorded between 1916 and 1950; average discharge during this 34 yr period was 3,609 cfs). Interval C had a thickness of 0.5 m (1.7 ft) in core # 13 (Manka and Steinmetz, 1971, p. 314). This indicates sediment accumulation rates of more than 50 cm as a result of a major flood event. Although Manka and Steinmetz presented no accumulation rates, with the exception of 10 cm of sand that was postulated to have been deposited during Hurricane Carla in 1961, they do note that sedimentation rates decreased after 1936 when river discharge was diverted into the Gulf. Assuming that interval D (fig. 60) was deposited between 1936 and 1941, as implied by Manka and Steinmetz, will yield rates of about 10 to 15 cm/yr. Using the thin sheet sand that Manka and Steinmetz postulated was deposited during Hurricane Carla as an upper time line and the 1933 line as depicted in figure 60 as the lower time line, approximately 1.8 m of sediment was deposited during this 28-yr period, indicating "longer term" average sedimentation rates of about 6.4 cm/yr.

Rates of sediment accumulation near a prograding delta greatly exceed rates of accumulation in bay and estuary environments away from the delta (Shepard, 1953; Rusnak, 1967). Van Heerden and others (1981) found that as much as 1.5 m of sediment was deposited on levees of the Atchafalaya Delta during a single flood event. As stated by Scruton (1960, p. 82), "Relative rapid deposition is the fundamental characteristic of deltas".

The Colorado River delta is somewhat unique in that the rapid progradation was primarily a result of the large sediment volume carried by the river (after removal of the log raft) and the shallow basin into which it discharged (Kanes, 1970; McGowen and Brewton, 1975). Storms punctuated the transportation and deposition of the sediments (fig. 58). The rate of aggradation apparently diminished more and more

in deltaic environments that were approaching sea level and becoming subaerial. Assuming that delta deposits depicted in figure 57 D aggraded from 0.3 to 1 m (1 to 3.3 ft) (estimate based on selected bathymetric data shown in fig. 57 C) between 1936 and 1941, the respective rates would be in the range of 6 to 20 cm/yr. The higher value is equivalent to Rusnak's (1967) estimated long-term average rates of sediment accumulation of 200 m per 1000 years in river deltas. In areas where the Colorado River delta has become subaerial and marshes have developed, the rates are substantially less, perhaps in the range of 1 cm/yr or less (White and Calnan, in preparation).

Recent Changes in State of the Colorado River Delta

The end of the constructional phase of the Colorado River delta in 1941, apparently can be attributed to several factors including discharge of the river into the Gulf and closure of distributary channels, but also, as indicated by Ward and others (1980), the year 1940 corresponds to the period of reservoir regulation on the Colorado River. The Highland Lakes chain, which intercepts sediments from about 90 percent of the Colorado River drainage basin (Kanes, 1970; USACE, 1977) has decreased the sediment supply at the mouth of the river to a fraction of the previous load (figs. 25 and 36). Ward and others (1980) conservatively estimated that the mean sediment load of the Colorado River is an order of magnitude below that which was responsible for the relatively constant rate of delta growth of 550 acres per year from 1929 to about 1938.

Dredging of Tiger Island Cut in the 1950's led to the development of a subdelta that prograded into Matagorda Bay at a rate of about 8.4 m/yr (28 ft/yr) (McGowen and Brewton, 1975). Comparisons of aerial photographs taken in the mid 1950's with those taken in 1979 show extensive marsh development on the Tiger Island Cut subdelta (White and others, 1985; fig. 61). The growth of the delta has been attributed to both river and tidal flow (USACE, 1977). Sand that has




Figure 61. Changes in marsh distribution between 1956 and 1979 on the Colorado River delta at Tiger Island Cut. (From White and others, in press a.)

contributed to this delta is apparently derived from the Gulf and not the river (van Beek and others, 1980; White and others, 1985; White and Calnan, in preparation). Model studies by the Corps of Engineers in 1959 indicate that tidal currents through Tiger Island Cut reach 1.2 m/sec; the tidal flow is flood dominated because tidal levels in the Gulf are higher than in the bay approximately 90 percent of the time; this promotes salt water flow into Matagorda Bay except during high river discharges (USACE, 1977). A significant amount of the Colorado River flow enters Matagorda Bay through Tiger Island Cut (USACE, 1970; TDWR 1978). Another subdelta that has grown in area since 1940 is located at Culver Cut (near the Intracoastal Waterway northwest of Tiger Island Cut); van Beek and others (1980) suggest that this area is supplied by sediment that is transported down the Intracoastal Waterway and out Culver Cut.

Manka and Steinmetz (1971) traced the destruction of the southeast delta lobe using aerial photographs taken in 1943, 1952, and 1965. These investigators reported that not only was the margin of the delta receding, or eroding due to wave attack, but compaction of underlying interdistributary sediments was promoting submergence of the delta plain and the formation of water bodies (interdistributary bays) that were enlarging through time (fig. 59). Kanes (1970) estimated that 30 to 60 cm (1 to 2 ft) of subsidence, and locally as much as 1.2 m (4 ft), had occurred in the northeast lobe of the Colorado River delta as a result of compaction. Historical shoreline changes along the margin of the Colorado River delta determined by McGowen and Brewton (1975) indicated that the eastern margin of the delta (along East Matagorda Bay) was eroding at a rate of 1.8 to 2.4 m (6 to 8 ft) per year; the western delta (with the exception of progradation at the mouth of Tiger Island Cut) remained relatively unchanged during the period 1957-1972.

Van Beek and others (1980) investigated the potential of the Colorado River to construct a delta if diverted into Matagorda Bay (USACE, 1981). Their predictive

models indicate that with full diversion, a delta will prograde into the eastern arm of Matagorda Bay at a rate of about 37 acres/yr. Of their estimated 1,370 acre-feet of sediment volume supplied by the river annually, 7.7 percent would be retained in the delta (van Beek and others, 1980). Van Beek and Meyer-Arendt (1981) reported that for Atchafalaya Bay, a comparison of sediment supplied with that deposited over a 10 year period (1967-1977) shows a retention of less than 30 percent, which is considerably below the 70 percent average calculated for the Mississippi River subdeltas.

The textural composition of the sediment delivered by the Colorado River is primarily suspended load composed of silt and clay (USACE, 1977). The Corps of Engineers (USACE, 1977) estimated that of the 1,650 acre-feet of sediment transported annually by the Colorado River, about 300 acre-feet is coarser bed load. Most of the bed load, or sand, is trapped up stream in a shallow draft navigation channel and silting basin that is periodically dredged. Welborn and Andrews (1980) investigated sediment discharge of the Colorado River at the town of Matagorda by collecting data from two floods and found that only 2.1 percent of the total load was bed load. Because little if any sand makes it to the mouth of the river, the textural composition of facies of the subdelta that forms at the mouth of the diversion channel will be considerably different from that in facies in the existing delta. In other words, there will be a fining of those facies (delta front and distributary channels) that commonly are rich in sand (Kanes, 1970; Manka and Steinmetz, 1971). Gosselink (1984) noted that reservoir construction in the drainage basin of the Mississippi River has presumably depleted the river's supply of sand, which is the main foundation material for growth of the delta; he suggests that the river, therefore, cannot support as large a delta as it has historically. As mentioned above, the Colorado River delta is primarily in a destructional stage.

Guadalupe River Delta

The Guadalupe River delta, located at the head of San Antonio Bay, is characterized by extensive brackish- to fresh-water marshes (McGowen and others, 1976b; Benton and others, 1977; White and others, 1985). Donaldson and others (1970) conducted a thorough geological investigation of the delta. Similar to the Mississippi Delta, the Guadalupe River delta, on a much smaller scale, has had a history of delta lobe growth, abandonment, and deterioration (figs. 50 and 62). Donaldson and others (1970, p. 108) stated: "In a general sense, the Guadalupe delta represents a model of sedimentation characterized by a stream depositing its load in a shallow, relatively quiet body of water. The delta progrades into increasingly shallow water so that the distributary channels have become deeper than the bay floor."

It is estimated that the delta began forming about 2,000 yrs ago in water slightly deeper than present (Shepard and Moore, 1960). Based on this date, the average rate of progradation into the bay was approximately 12 m/yr (40 ft/yr) (Donaldson and others, 1970). Growth of the delta complex occurred in a counterclockwise direction (fig. 50) as the major distributaries apparently migrated in response to steeper gradients in that direction. An estimated 30 cm (1 ft) of subsidence on older abandoned subdeltas resulted in the transgression of marshes and barren tidal flats over natural levees (Donaldson and others, 1970).

A cross-sectional reconstruction of the Guadalupe River delta facies reveals six major environments of deposition that are composed of distinctive sediment textures: (1) distributary channel, (2) natural levee, (3) marsh, (4) interdistributary bay, (5) delta front, and (6) prodelta (fig. 63). Sediments of the delta front and basal part of distributary channel deposits are fine-grained sand, silty sand, or silt, while the remaining environments are composed principally of clay and silty clay (Donaldson and others, 1970). Abundant plant material occurs in the silty clay sediments of the




Figure 62. Map showing idealized spatial relationship of depositional environments of the Guadalupe River delta prior to formation of the Traylor Cut subdelta. (From Donaldson and others, 1970.)




Figure 63. Generalized cross section of the Guadalupe delta showing facies. (From Donaldson and others, 1970.)

marsh and natural levee. The configuration of the delta-front sand deposits is predominantly shoestring (barfingers), but locally, closely spaced distributary-mouth bars coalesce to form sheet sands. The average thickness of the delta (not including the channel-fill deposits which are as thick as 4.86 m [16.2 ft]) is 2.7 m (9 ft) (Donaldson and others, 1970). The delta plain environments, which include distributaries, natural levees, and marshes, make up the greatest volume of sediment in the delta. The thickness of the delta-plain deposits, which average 1.5 m (5 ft), decrease outward from the distributaries.

Sediment is delivered to the delta by both the Guadalupe and San Antonio Rivers, the confluence of which is less than 16 km (10 mi) upstream from the delta. Donaldson and others (1970) indicated that the average annual suspended load transported to the delta is approximately 1 million tons, with nearly equal amounts of sediment contributed by the Guadalupe River (461,214 tons annually for the period 1945-1954) and San Antonio River (568,218 tons annually for the period 1942-54) (fig. 20). Textural analysis of sediments collected from the channel bed of the Guadalupe River from Victoria to its mouth indicated a substantial decrease in grain size from granules and pebbles near Victoria with changes downstream to sand and sandy silt and then to dominantly clayey silt at the river mouth (Morton and Donaldson, 1978). Composition of suspended sediment of the Guadalupe is approximately 2 percent very fine sand, 27 percent silt, and 71 percent clay (Welborn, 1967). Morton (1972) reported that clay minerals in the Guadalupe delta as well as in San Antonio Bay are predominantly montmorillonite, illite, and kaolinite, with montmorillonite most abundant. In a comparison of the suspended load of the Guadalupe and San Antonio Rivers, Sorenson (1975) found that the suspended load of the Guadalupe is higher in organics than that of the San Antonio River. Although there is a rapid increase in inorganic sediments relative to organics during flood events, Sorenson concluded that measurements of suspended load during normal flow conditions, when organics are

high, can lead to overestimates of sediment-accumulation rates in the delta. This conclusion is based on the probability that organic debris will not be preserved in substantial quantities.

Recent Changes in State of the Guadalupe River Delta

A major modification occurred on the Guadalupe delta in 1935 with the dredging of Traylor Cut, which diverted about two-thirds of the discharge of the Guadalupe River into Mission Lake (Morton and Donaldson, 1978). This diversion resulted in the rapid progradation of a subdelta into the southwestern corner of the Lake. The historical development of the delta was depicted by White and Morton (1987) (fig. 64). Donaldson and others (1970) estimated the rate of delta growth (progradation) at about 22.5 m/yr (75 ft/yr), noting that Mission Lake, which was less than 1.5 m (5 ft) deep when growth of the delta began, was less than 0.6 m (2 ft) by 1970. White and Morton (1987) reported even higher rates of progradation of the Traylor Cut subdelta during selected periods, and noted that historical progradation rates have apparently varied, in part, because of a log raft that developed along the river. Progradation rates between 1958 and 1974 were slower than after the mid-1970's when the logjam was removed. Rates of aggradation of the Traylor Cut subdelta can be estimated from thickness of deltaic deposits presented by Donaldson and others (1970). The cumulative thicknesses of pro delta and overlying marsh deposits in the subdelta were approximately 1.56 m (5.2 ft) thick. This thickness would yield an average sedimentation rate of about 5 cm/yr (for the period 1935 to 1965). An isopach map of pro delta deposits at the Traylor Cut subdelta just offshore from delta plain deposits (Donaldson and others, 1970) shows a thickness of about 1.26 m (4.2 ft), which indicates an aggradation rate of approximately 4 cm/yr (0.13 ft/yr). These deltaic rates of aggradation are an order of magnitude higher than the average rate of sedimentation (shoaling) in San Antonio Bay (0.37 m/100 yrs [1.23 ft/100 yrs]) determined by Shepard (1953).




Figure 64. Shoreline changes along the Traylor Cut subdelta, based on aerial photographs, 1929 to 1982. (From White and Morton, 1987.)

Diversion of river discharge and sediment load through Traylor Cut apparently has hastened the retreat of the delta shoreline along Guadalupe Bay near the mouth of the North and South Guadalupe Rivers, which were the natural, active distributaries of the delta (Donaldson and others, 1970; White and Morton, 1987). Donaldson and others (1970, p.111-112) stated that "Relatively strong tidal flow through the narrow Guadalupe Bay and reduced discharge and load have transformed the North-South Guadalupe from a prograding to a deteriorating subdelta". The shoreline along most of the delta retreated (eroded) between 1930 and 1982, at an average rate of 1.5 m/yr (5 ft/yr); rates of erosion were locally as high as 2.7 m/yr (9 ft/yr) (White and Morton, 1987). Highest rates of erosion occurred on the gulfward half of the delta, where abandoned delta lobes do not receive sufficient fluvial sediment to counteract the effects of delta subsidence and wave and current action in San Antonio and Guadalupe Bays. A comparison of photographs taken in the mid-1950's with those taken in 1979 indicates an increase in water bodies and shallow subaqueous flats (fig. 65). It appears that under current conditions, the Traylor Cut subdelta will continue to prograde into Mission Lake, producing additional wetland acreage, while inactive parts of the delta will continue to deteriorate through erosion and submergence. (Additional data on sedimentation in this area is presented in a later section on San Antonio Bay.)

Trinity River Delta

The Trinity River, like the Colorado and Guadalupe Rivers, has constructed a delta at its mouth (figs. 1 and 66). The salinity regime that characterizes this deltaic area is more similar to that of the Guadalupe River than the Colorado River, and thus, the Trinity River delta is the site of extensive brackish- to fresh-water marshes (Fisher and others, 1972; Rice Center for Community Design and Research, 1974; Adams and Tingley, 1977; Benton and others, 1979; White and others, 1985). The Trinity and Colorado Rivers are similar in that they are the only two Texas rivers




Figure 65. Changes in the distribution of wetlands between 1957 and 1979 on the seaward half of the Guadalupe River delta. (From White and others, in press b.)




Figure 66. Trinity River delta. (From McEwen, 1969.)

(discharging into estuaries) that have appreciably extended their deltas since the mid-1800's (Shepard, 1953). Shepard reported that the Trinity delta had advanced a distance of about 0.48 km (0.3 mi) since 1855.

Sedimentary facies of the modern Trinity River delta have been investigated by McEwen (1969), from which most of the following discussion was derived. Similar to the other estuarine deltas discussed previously, the Trinity River delta has prograded into a relatively protected, shallow bay. McEwen (1969) defined the following depositional units in the Trinity River delta: (1) channel and associated deposits, (2) natural levee, (3) interdistributary bay and marsh, (4) delta-front churned sands, and (5) prodelta.

Composition of channel deposits, which include point bars, ranges from sand at the base of the channel to silty clay at the top. Sediment textures composing natural levee deposits vary from silty clay to clayey sand; plant debris and roots are common. Natural levees, which border the Trinity River and distributary channels, range from about 0.9 m (3 ft) in elevation in more landward parts of the delta, to sea level and below at the distributary mouths. According to McEwen (1969), the widths of natural levees range from a few feet along smaller distributaries to several hundred feet along the main river channel. Channel mouth bars are composed primarily of sand. Submarine levees are gradational with the channel mouth bars with sediment textures decreasing from channels toward the levee deposits. At the time of his study, McEwen reported that Jack's Pass (fig. 66) was the only active distributary that was in a relatively natural state. Other distributaries, for example, Blind Bayou, Southwest Pass, Triangle Pass, and Old River Pass (fig. 66), were abandoned and were being filled with sediments. Because the process of abandonment and filling is a gradual one, sediment characteristics are reflective of channel as well as lake and marsh deposits, and accordingly, range from sands and clays, to muds.

In terms of areal distribution in the subaerial part of the delta, the marsh facies is the most important. Marsh sediment composition is predominantly organic-rich clays and sandy clays. The organics are composed of very fine-grained material, as well as plant debris and rootlets. Marsh deposits with the highest sand content occur near distributaries and delta-front environments. McEwen examined sediments in Round Lake to define the characteristics of lake deposits, which he described to be mostly gray to black clay with local silty burrows and lenses. The sediments could be distinguished from marsh sediments because of a complete lack of visible plant remains. Interdistributary bays are shallow environments that are bordered by distributary levees and marshes. Composed principally of clayey sand in central areas, these deposits grade laterally into submarine levee, marsh, and delta-front deposits.

The delta-front deposits (delta-front churned sands) are several feet in depth and are composed of sandy sediments that extend from the front of the delta to a distance of about 2.4 to 3.2 km (1.5 to 2 mi). The sediments decrease in grain-size toward the bay, grading from well-sorted, fine sands to the almost pure clays that make up the open bay sediments. The bay or prodelta facies is predominantly black mud (McEwen, 1969).

The delta-front sands, which in most areas have been extensively churned by burrowing organisms, are volumetrically the most important depositional unit in the delta. In fact, McEwen (1969, p. 74) described the delta "as a mass of delta-front sands capped by a veneer of marsh deposits and transected by linear bodies of channel deposits". The base of the delta-front sands is about 4.5 to 4.8 m (15 to 16 ft) below sea level in the oldest part of the delta (northern part) where the sands reach a maximum thickness of between 4.2 and 4.5 m (14 and 15 ft). The unit thins to the east and southeast, apparently due to shoaling of the bay in that direction. Marsh deposits have a uniform thickness of 0.6 to 0.75 m (2 to 2.5 ft) in the older part of the delta. The base of marsh deposits, commonly about 0.3 m (1 ft)

below sea level, is as much as 0.6 m (2 ft) below sea level in the central part of the interdistributary bays. These clayey deposits may reach 0.6 m (2 ft) above sea level near the Trinity River.

The oldest radiocarbon dates measured in shells (Rangia flexuosa) from cores in the delta-front sands were 810 and 750 years B.P. (Before Present). Assuming that approximately 5.5 m (17.5 ft) (thickness of the delta-front sands + marsh deposits) accumulated over this period yields a long-term average accumulation rate of about 7 mm/yr. Assuming that a thickness of 4.5 m (15 ft) accumulated during a period of 500 yrs (Rice, 1969) will yield a rate of about 9 mm/yr. These rates are close to the average 8 mm/yr sea-level rise estimated by Emery and Uchupi (1972) for the past 5,000 yrs. It is presumed that sedimentation rates were high in the actively prograding parts of the delta and decreased substantially as delta surfaces became emergent and aggraded above sea level.

Rice (1967; 1969) calculated that sediment delivered by the Trinity River to the delta was made up of the following approximate textural percentages (in the suspended load): 25 percent sand, 30 percent silt, and 45 percent clay. In calculating the mass balance between the volume of sand delivered by the river and the volume composing the delta he found an imbalance. Considering the area of the modern delta to be about 6 mi² (about 3,800 acres or 1500 hectares), and assuming that the delta was composed wholly of delta-front deposits with a thickness of 4.5 m (15 ft) and with a composition of 75 percent sand, Rice (1969) calculated that the delta contains a volume of 43,200 acre-ft of sand. The assumption that over the past 500 years the Trinity River had a constant annual discharge of 3,000 acre-ft of sediment (based on suspended load records upstream at the Romayor station; fig. 20) composed of 25 percent sand allowed Rice to calculate a total volume of 375,000 acre-ft of sand. A comparison of these two volumes indicates almost ten-times as much sand was transported by the river as is found in the delta. Rice noted that

sand does not appear to be moving across the mouth of modern Trinity Bay and hypothesized that the lower portion of the Trinity River must be aggrading its valley through valley subsidence. In other words, the sand, assisted by the process of subsidence, was being deposited by the river upstream from the delta in floodplain and point-bar channel deposits.

Failing (1969), referring to the imbalance between sand delivered by the river and that found in the delta (Rice, 1969), noted that the Trinity River valley between Romayor and the delta had an area of about 500 mi^2 (320,000 acres; 129,600 hectares) and that sediment spread over this area to a thickness of 1 ft (30 cm) would have a volume of about 300,000 acre-ft (the approximate amount of the imbalance noted in the preceding paragraph). Failing (1969) concluded that because 30 cm (1 ft) of sediment equals the amount of subsidence that would have occurred over the past 500 yrs at the rate of 6 cm/100 yrs (0.2 ft/100 yrs) (Rehkemper, 1969), then subsidence and sedimentation must be occurring at the same rate in the lower Trinity River valley.

McEwen (1969) made the observations that marsh deposits in the delta were thin and that no marsh or levee deposits were found at depths greater than 0.6 m (2 ft) below sea level. This led McEwen to conclude that no measurable subsidence had occurred during the growth of the delta, and there had been no relative change in elevation of the delta and sea level since delta formation began. This is somewhat puzzling, especially because of the hypothesis that subsidence was occurring along the river valley (Rice, 1969; Failing, 1969). Failing (1969) suggested, however, that if the delta was formed between 500 and 1,000 yrs ago, as indicated by radiocarbon dates (McEwen, 1969), and the minimum subsidence rate is 6 cm/100 yrs (0.2 ft/100 yrs) (Rehkemper, 1969), then total subsidence of the delta would be on the order of 30 to 60 cm (1 to 2 ft), which may have gone undetected in McEwen's coring investigation.

For comparison, estimated compactional subsidence of parts of the Colorado River delta are on the order of 30 to 60 cm (1 to 2 ft), and locally as high as 1.2 m (4 ft) (Kanes, 1970). Donaldson and others (1970) estimated that at least 0.3 m (1 ft) of subsidence had occurred on the Guadalupe River delta as reflected in the transgression of marshes over relict levees.

Recent Changes in State of the Trinity River Delta

As mentioned previously, the Trinity River delta is one of only two Texas bay-head deltas identified by Shepard (1953) as having undergone significant progradation in recent historic times. Analysis of historical shoreline changes also indicates local delta shoreline progradation in which marshes have advanced bayward at rates of more than 4.3 ft/yr (with a high, locally, of 103.4 ft/yr) between 1930 and 1982 (Paine and Morton, 1986; fig. 67). Older inactive parts of the delta (stations T46 to T51 in fig. 67) retreated at rates of 3.8 ft/yr to 10.2 ft/yr between 1930 and 1974 or 1982. Shorelines at most stations on the active delta lobe (T45 to T51) prograded (accreted) into the bay during the period 1930 to 1982. The shoreline at delta station T42, however, eroded. In addition, although shoreline changes at station T43 indicated net accretion between 1930-1982, this shoreline began to retreat (erode) during the latter part of this period as indicated by a comparison of marsh areas mapped on 1956 photographs (Fisher and others, 1972) and 1979 photographs (White and others, 1985). Comparison of marsh changes in interior deltaic areas indicates some encroachment of water into areas formerly mapped as marshes (White and others, 1985). Trends toward erosion and marsh submergence may be indicative of subsidence and reductions in sediment supplied to the delta.

Subsidence in the vicinity of the delta between 1943 and 1978 was about 22.5 cm (0.75 ft) (Gabrysch, 1984). This rate, approximately 6 to 7 mm/yr, is not exceptionally high. However, there has apparently been a marked decline in the quantity of fluvial sediment delivered to the delta by the Trinity River since about




Figure 67. Rates of shoreline change along Trinity Bay and Trinity delta, 1930 to 1982. (From Paine and Morton, 1986.)



1969 when Lake Livingston was constructed (fig. 35). Declines in sediment load of the river are treated more thoroughly in a later section (Trends in Relative Sea-Level Rise and Fluvial-Estuarine Sedimentation), but in general terms, recent suspended-load measurements (1969-1984) show that the average annual load of the Trinity River at Romayor (located downstream from Lake Livingston) is only about 14 percent of that reported by Shepard (1953). As stated by the U. S. Corps of Engineers: "At present, only about 2 percent of the sediment entering Lake Livingston passed through the spillway, and most sediment contributions from tributaries below Livingston dam will be inhibited by Wallisville dam. It seems likely that, in the future, aggradation and progradation of the delta will occur only during high flows, and that the primary agents in the shaping of the delta will be wind-generated wave erosion and deposition" (USACE, undated, p. A-10).

Recent Changes in State of Wetlands in Other Fluvial-Deltaic Areas

The preceding discussion on the depositional framework, development, and current state of the Colorado, Guadalupe, and Trinity river deltas sets the stage for examining recent conditions and trends in other bay-head delta and fluvial areas along the Texas coast. Two fluvial-deltaic areas, San Jacinto River and Neches River, stand out because of relatively recent extensive losses in wetland acreage (White and others, 1985; 1987).

San Jacinto River

A comparison of the distribution of wetlands as interpreted from photographs taken in 1956 and 1979 of the lower San Jacinto River valley at the head of Galveston Bay indicates a displacement of more than 560 hectares (1,380 acres) of fluvial woodlands, swamps, and fresh- to brackish-water marshes by open water (fig. 68). The principal reason for this loss of wetlands is that the lower reach of the San Jacinto River is in an area of subsidence (fig. 9) caused primarily by ground-water



Wetland Map Unit	1956		1979		Net Change	
	Acres	Hectares	Acres	Hectares	Acres	Hectares
Water	875	355	2,295	930	+1,420	+ 575
Fluvial Woodlands & Swamps	3,480	1,410	2,090	845	-1,390	- 565
Fresh to Brackish Marsh	650	265	300	120	-350	-145

QA 937

Figure 68. Changes in the distribution of wetlands between 1956 and 1979 of a subsiding segment of the San Jacinto river near Houston, Texas. (After White and others, 1985.)

withdrawal (Gabrysch, 1984). Between 1943 and 1978, between 1.2 and 2.1 m (4 and 7 ft) of subsidence occurred in this part of the river valley (Gabrysch, 1984). The San Jacinto River lies within an entrenched valley similar to that of the Trinity River (fig. 3), and as subsidence occurs, submergence and resulting changes in wetland environments progress inland along the axis of the valley (fig. 69).

The change in wetlands along the San Jacinto River is pronounced because of the proximity of the valley to the center of maximum subsidence. Rates of subsidence, between 1943 and 1978, are as high as 60 mm/yr, which apparently greatly exceeds the rate of wetland aggradation in this area. The volume of sediments reaching the mouth of the San Jacinto River has diminished as a result of reservoir development in the drainage basin (fig. 70). Lake Houston, which is located only a few miles (< 10 mi) upstream from the mouth of the San Jacinto River, has an estimated trap efficiency of about 87 percent (USDA, 1959), which suggests that only small quantities of sediment are delivered to the area where wetlands are being submerged. Nevertheless, even without Lake Houston and other reservoirs in the San Jacinto River basin, land-surface subsidence is so pronounced that it is unlikely aggradation rates could keep pace with subsidence rates except farther upstream where the rates decline. It appears that submergence will continue up the axis of the valley in the future, but rates of change should diminish somewhat as bay waters move into areas with slightly higher elevations and lower rates of subsidence.

Neches River

The most extensive loss of wetlands in a modern estuarine fluvial-deltaic system on the Texas coast has occurred along the Lower Neches River (White and others, 1987). The Neches River discharges at the head of Sabine Lake; the lower part of the river valley is the site of relatively extensive marshes that have developed on fluvial-deltaic deposits (Fisher and others, 1973). Losses in wetlands in this area have been reported by Wiersema and others (1973), Gosselink and others (1979), and



Figure 69. Aerial photograph of the lower reaches of the San Jacinto River. Compare with figure 68.








Figure 70. Cumulative reservoir storage capacity for selected rivers along the Texas coast. Data from Texas Water Development Board.

White and others (1987). Between the mid-1950's and 1978, about 3,800 hectares (9,410 acres) of marshes were displaced primarily by open water along an approximately 16-km (10-mi) stretch of the lower Neches River valley (fig. 71). The displacement of marshes by open water and shallow subaqueous flats in this area is apparently related to several factors (Wiersema and others, 1973; Gosselink and others, 1979; White and others, 1987), including: (1) relative sea-level rise resulting primarily from subsidence possibly due to both natural and human-induced causes (Swanson and Thurlow, 1973; Ratzlaff, 1980), (2) a decline in sediments supplied to this alluvial area as a result of (a) reservoir development in the Neches River basin (TDWR, 1981c; fig. 70), (b) artificial levees (dredged spoil, Fisher and others, 1973) along the dredged portion of the river, and (c) changes in hydrology due to artificial channels (Wiersema and others, 1973), (3) active faulting, in which the downthrown side of one identified fault is subsiding at a more rapid rate than the upthrown side (White and others, 1987; fig. 13 and 71), and (4) dredged canals, which can cause direct and indirect losses in marshes (Wiersema and others, 1973; Scaife and others, 1983).

In a brackish-water marsh about 50 km (30 mi) east of Sabine Lake on the Chenier Plain (gulfward of Calcasieu Lake, Louisiana), DeLaune and others (1983) reported changes from marsh area to open water are occurring at ever increasing rates apparently because marsh aggradation rates (averaging 0.8 cm/yr) are not keeping pace with submergence rates (averaging 1.2 cm/yr). The change in marsh area to open water has been increasing by a factor of approximately 2 every 6 years since 1963. DeLaune and others (1983) predict that this marsh area will complete its transformation to open water in less than 40 years if the trends that have characterized the past 25 years continue. Among the human activities that may be contributing to the transformation to open water are (1) ship channel construction (promoting salt intrusion and possibly sediment diversion) and (2) oil, gas, and



QA3384

Map Unit	1956		1978		Net Change*	
	Acres	Hectares	Acres	Hectares	Acres	Hectares
Water	2,560	1,037	11,070	4,483	+8,510	+3,446
Marsh	15,740	6,375	6,330	2,564	-9,410	-3,811

* Loss of additional 900 acres (365 hectares) of marsh primarily due to spoil disposal.

Figure 71. Changes in the distribution of wetlands between 1956 and 1978 in the lower Neches River valley near the head of Sabine Lake. The fault crossing this area has apparently contributed to the changes. (From White and others, 1987.)

groundwater withdrawals (accelerating subsidence). However, DeLaune and others (1983) concluded that it is difficult to document the human component precisely because of its pervasiveness in this area and because some of the observed trends contradict expected effects.

Similar to those of the marsh system near Lake Calcasieu, the factors contributing to marsh loss listed in the preceding discussion of the Neches River valley are complex and impossible to quantify adequately with existing data. But the conversion of marsh to open water indicates marsh aggradation rates are not keeping pace with subsidence rates and relative sea-level rise. Subsidence rates in this area of the Neches River valley are not known, but the rate at Sabine Pass reported by Swanson and Thurlow (1973), for the period 1960 to 1969, is 1.25 cm/yr, a rate equal to that reported by DeLaune and others (1983). This rate is among the highest reported along the Texas coast (fig. 7). In the Neches River valley, the rate of subsidence may be similar or possibly even higher due to withdrawal of underground fluids (Ratzlaff, 1980) and faulting (White and others, 1987). Over a 22-year period (1956-1978) the marsh has been replaced by open water at an average rate of about 160 hectares/yr (400 acres/yr) (fig. 71). It is possible that the rate of change has increased in more recent years as reported by DeLaune and others (1983) for the marsh near Lake Calcasieu. There is little evidence to suggest that there will be a reversal in trends, so it is likely that transformation of marsh to open water in the Neches River valley will continue in the future.

Lavaca River





The Lavaca River fluvial-deltaic area is small compared to the Colorado, Guadalupe, and Trinity river deltaic areas. The delta extends only a short distance into Lavaca Bay; fluvial-deltaic deposits capped by marshes are principally restricted to the alluvial valley stretching inland from the head of the bay (fig. 1). An analysis of historical changes in deltaic shorelines and associated marsh areas in the Lavaca River

delta indicates that the configuration of the delta margin did not change significantly between the mid-1850's and mid-1950's (McGowen and Brewton, 1975). Only about 39 hectares (96 acres) of new wetlands formed as a result of deltation at the mouth of the Lavaca River; the active delta prograded about 4.3 km (2.7 mi). Inactive parts of the delta to the west eroded at rates of 0.1 to 1.5 m/yr (1 to 5 ft/yr) since 1934 (McGowen and Brewton, 1975). McGowen and Brewton reported that the Lavaca River delta was stable in terms of marsh surface-water level relationships, indicating a balance between sedimentation and subsidence rates.

Upstream from the delta along the Lavaca River valley, relative sea-level rise due principally to subsidence has apparently contributed to local wetlands submergence and the expansion of water bodies into areas formerly occupied by marshes and fluvial grassland (White and others, 1985; fig. 72). Between 1918 and 1973, subsidence of approximately 15 cm (0.5 ft) has occurred over a relatively broad area and locally has exceeded 30 cm (1 ft) (Ratzlaff, 1980). Aggradation rates do not appear to be keeping pace with subsidence rates in this area (fig. 72) of the Lavaca River valley. Not shown in figure 72 are extensive alterations of the natural hydrologic regime by artificial levees and canals. Reductions in fluvial sediments supplied to areas like this one, will probably accelerate the transformation of wetlands to open water. Sediment yields to the mouths of the Lavaca River and Navidad River drainage basins are reportedly about 273 and 784 acre-ft/yr, respectively (Greiner, 1983). Lake Texana, completed in 1980, effectively traps most of the sediment delivered through the Navidad River system. It seems likely that fluvial-deltaic areas in the Lavaca River valley will fall farther behind aggradation rates promoting more extensive submergence of valley wetlands in the future.

Nueces River

As pointed out by Morton and Paine (1984), Nueces Bay, into which the Nueces River discharges, is shallower, has a shorter wave fetch, and receives more fluvial sed-



045048

Figure 72. Changes in the distribution of wetlands between 1958 (A) and 1979 (B) at Menefee Flat in the Lavaca River valley. (From White and others, in press b.)

iment than nearby Corpus Christi Bay; these factors can promote shoreline accretion and reduce erosion. Monitoring of historical changes in shorelines from map surveys and photographs made in the late 1800's to 1982 indicates net accretion (advancement of the shoreline toward Nueces Bay) of marshes on the Nueces River fluvial-deltaic complex. Although net progradation was recorded for this period (1867 to 1982), aerial photographs taken in 1959, 1971, and 1974 indicate that marsh progradation ended sometime between 1930 and 1959 (Morton and Paine, 1984).

Comparison of interior marshes in the Nueces River fluvial-deltaic system on photographs taken in the mid 1950's and late 1970's suggests there has been an increase in shallow subaqueous flats and a corresponding decrease in marsh area (White and others, 1983). Subsidence reported in the Nueces fluvial-deltaic area is interpreted to be on the order of 6 to 30 cm (0.2 to 1.0 ft) for the period 1942-1951 (Brown and others, 1976). These amounts of subsidence translate into annual rates of about 0.7 to 3 cm. It is possible that marsh sedimentation rates are falling behind subsidence rates. Fluvial sediments along the Nueces River have been reduced by Lake Corpus Christi (fig. 32). In fact, more recent annual averages of suspended sediments measured on the Nueces River are only about 4 percent of those reported by Shepard (1953) (see Corpus Christi-Nueces Bay System presented in the section on Bay-Estuary-Lagoon Sedimentation). Sediment yield at the delta for the drainage area below Lake Corpus Christi is approximately 190 acre-ft/yr (Greiner, 1982). If all the suspended load measured below Lake Corpus Christi at Mathis were to reach the delta, this would add another 19 acre-ft/yr. These data indicate that an average of about 210 acre-ft of fluvial sediment reaches the delta each year. This volume is about 40 percent of pre-1952 sediment loads reported by Shepard (1953) for the Nueces at Three Rivers.

BAY-ESTUARY-LAGOON SEDIMENTATION

Sediment Composition

Benthic sediment textures in the bay-estuary-lagoon system along the Texas coast have been mapped, most extensively, by McGowen and Morton, (1979) and White and others (1983, 1985, 1986, 1987, in press a and b, and in preparation). Earlier maps depicting sediment distribution in selected Texas bays include those by Shepard and Moore (1955), Fagg (1957), Shenton (1957), Elliot (1958), Rusnak (1960), Kane (1959, 1966), Alaniz and Goodwin (1974), Shideler and others (1981), Hall (1976), and McGowen and others (1979).

Bay sediment types are principally sand, mud (silt and clay), and gravel (composed primarily of shell material). The distribution of these different textural components follows a somewhat systematic trend for the larger bay systems, where muds characterize the deeper bay centers and sand the shallower bay margins (figs. 73 to 79). The distribution of sediments is in part a reflection of the wave and current energy, which is related to water depth. Sands along the margins of larger bays reflect not only nearby sources of sand but also the relatively high energy of these shallow environments where breaking waves and littoral currents are common. Sand eroded from the bay margins remains in this environment because current energy decreases in deeper water. In a study of bay margin sands in the Galveston-Trinity Bay system, Love and others (1985) found a correlation between prevailing and dominant wind directions and bay-margin sands, indicating a relationship between the wind-generated wave and current energy and water depth of sand deposition. Sands are concentrated (1) at the mouths of rivers where channel mouth sand bars are deposited and reworked in near shore areas, (2) at the mouths of tidal inlets where current energy is sufficient enough to transport sand, and (3) along the margins of sand-rich barrier islands and peninsulas (figs. 75 and 76). Shelly




Figure 73. Index for sediment distribution maps shown in figures 74-79.






Figure 74. Distribution of sediments in Sabine Lake. (From White and others, 1987.)



QA-3597

Figure 75. Distribution of sediments in Galveston Bay. (From White, 1985; modified from White and others, 1985)








Figure 77. Distribution of sediments in San Antonio and selected bays to the southwest. (From Shepard and Moore, 1960.)



QA-3596

Figure 78. Distribution of sediments in Corpus Christi and Nueces Bays. (From White, 1985; modified from White and others, 1983.)



QA-3598

Figure 79. Distribution of sediments in southern Laguna Madre. (From White, 1985; modified from White and others, 1986.)

substrates are concentrated near oyster reefs and locally along bay margins where shell is deposited by waves and currents in beaches and storm berms.

Inorganic bottom sediments in the bay-estuary system are mostly composed of quartz, feldspar, and clay minerals. Oolites and coated grains are among the constituents of sediments in Baffin Bay along the south Texas coast (Behrens, 1963; Dalrymple, 1964; Frishman, 1969). In the suite of clay minerals, montmorillonite is more abundant than illite and kaolinite along the western coast of the Gulf of Mexico (Elliot, 1958; Morton, 1972; Sorenson, 1975; Byrne, 1976; Folger, 1972). Along the eastern Gulf and southeast Atlantic coast kaolinite is predominant, and along the northeast Atlantic coast, illite and chlorite are most abundant.

Sources of Sediment

Kranck (1984), in citing Judson (1968) and Milliman and Meade (1983), noted that about 90 percent of the 15×10^{15} g/yr of terrestrial sediment delivered by the world's rivers is deposited near the continents, primarily in estuaries. Most of the estuaries along the Atlantic Coast appear to be filling without much transfer of sediments to the shelf (Meade, 1982). Although landward sources of sediments are apparently the principal ones that supply sediments to estuaries (Meade, 1982), transport of sediments from the marine system into the estuaries, or from the mouth of the estuary landward, appears to be an important process (Stewart, 1958; Ritter, 1969; Windom and others, 1971; Pevear, 1972; Emery and Uchupi, 1972; Schubel and Carter, 1976; Summerhayes and others, 1985; Renwick and Ashley, 1984). Studies of sedimentation in Chesapeake Bay indicate that the importance of various sources of sediment changes from the head of the bay to its mouth (Schubel, 1968, 1972; Schubel and Carter, 1976, 1984; Biggs, 1970; Biggs and Howell, 1984; Officer and others, 1984). Approximately 70 percent or more of sediment from the Susquehanna River, which is the principal source of sediment to the main body of

Chesapeake bay, is trapped in the upper 30 km of the bay (Biggs, 1970; Schubel, 1972). Schubel (1968) estimated that in northern Chesapeake Bay the annual contribution of sediment from river runoff was about 5 times the amount from shoreline erosion. In the middle bay, bay shoreline erosion is the primary source, and near the mouth of the estuary, sediments are derived principally from a marine source. Yarbrow and others (1983), in a study of the sediment budget of a major estuary (Choptank River Estuary) on the eastern shore of Chesapeake Bay, found shoreline erosion to be the principal source of sediment, contributing seven times the volume supplied by upland runoff. Factors influencing the predominance of erosional sources over river sources were low relief, rural character of the drainage basin, and poorly consolidated shoreline material, which contributed to high erosion rates (Yarbrow and others, 1983).

Rusnak (1967) classified estuaries on the basis of the principal sediment source areas that were filling the estuary. Estuaries primarily filled from a land source (fluvial) were positive-filled estuaries, from a marine source, negative-filled, and from an internal source (redistribution of sediments), neutral-filled (or non-filling). Those Texas bays with relatively large fluvial sources, such as Sabine, Trinity, Lavaca, San Antonio, and Nueces would be classified as positive-filled, whereas Laguna Madre, with a principal source from the Gulf would be negative-filled. Some estuaries are being filled from both a marine and landward source (fig. 80). Although most of the Texas estuaries along the central and northern Gulf coast are dominated by sediment deposition from fluvial sources, and secondarily from shoreline erosion, deposition also occurs at the mouths of tidal inlets (flood tidal deltas) indicating a marine source. The sources and major depositional features, as well as some of the hydrodynamic interactions in the estuarine basins are depicted in figure 81.

Principal sources of sediments in the Texas bay-estuary-lagoon system include (1) suspended load and bed load of rivers and streams, (2) erosional products from





Figure 80. Conceptual model of sediment sources and associated deposition relative to the magnitude of the source. A system like the Mississippi in which fluvial sediments escape seaward is depicted in "A"; a system like Chesapeake Bay in which fluvial and marine sediments are trapped in the estuary is depicted in "B"; a system like Laguna Madre in Texas, which has relatively small fluvial input is depicted in "C". (From Biggs and Howell, 1984.)



DISPERSAL ZONES & ROUTES




Figure 81. Depositional features typical of Texas bay-estuary systems (A) (From McGowen and others, 1979), related to sediment sources and processes (B and C) (From Nichols and Biggs, 1985).

bay margins, (3) Gulf or marine sediments transported through tidal passes and storm washover channels, (4) sediments transported across barrier islands and peninsulas by eolian processes, (5) nonterrestrial biogenic materials, primarily oyster shells (serpulid reefs in Baffin Bay), and (6) dredged material placed on the bay floor or bay bottom. Along the central and upper Texas coast, rivers that discharge into the bay-estuary-lagoon system are the principal source of sediments. For example, in the Lavaca Bay system, Wilkinson and Byrne (1977) concluded that rivers discharging into the bay account for roughly 73 percent of the sediment deposited there. Sediment from shoreline erosion was a significant, although secondary source, supplying about 24 percent of the sediment supply. Estimates of sediment contributions from various sources in the Galveston-Trinity Bay system by the U.S. Army Corps of Engineers (USACE, 1942; cited in Rehkemper, 1969), indicate that fluviially derived sediments make up more than 90 percent of the total supply, with shoreline erosion contributing less than 5 percent. Along the southern Texas coast, where major rivers do not discharge into the bays and lagoons, fluvial sediment input is less important. The major sources of sediment deposited in Laguna Madre, for instance, are the Gulf and Padre Island, from which sand is transported by eolian and storm washover processes (Price and Gunter, 1943; Lohse, 1955; Price 1958; Fisk, 1959, Hunter and Dickinson, 1970).

Turbidity

The characteristically turbid nature of waters in an estuary is a product of: (1) particulate matter from the watershed, off-inlet shores and bottoms, reworking and scouring of estuarine bottoms by tidal currents and waves, loosening of bottom sediments by burrowing animals, and decomposition of pelagic and benthic organisms; (2) the net two-layered opposing estuarine circulation pattern; (3) the mixing of fresh and salt water and consequent flocculation of finer particles; and (4) the presence of

relatively quiet sedimentation areas provided by semi-enclosures and widening of the estuarine basin (Carriker, 1967). The release of large quantities of organic-laden silts and clays into the water column is a dominant feature of Texas bays and estuaries whether from beach erosion, river input, storm reagitation or dredging (Odum and Wilson, 1962).

Turbidity generally varies widely throughout the year and reaches a maximum during floods of the rainy season. At all times of the year, there is a gradual decrease in turbidity with distance from the river mouth (Emery and Stevenson, 1957). As a consequence of estuarine tidal mixing, sedimentation, and reworking processes, turbidity is higher at a given point in the estuary at low than at high tide (Carriker, 1967). Turbidity can also change considerably over short distances and from hour to hour. For example, Mackin and Hopkins (1961) studied oyster mortality in Barataria Bay, Louisiana, and reported turbidity readings ranging from 18 to 198 ppm over a distance of less than 200 ft (61m). Also, after a combination of high wind and rainfall, Mackin and Hopkins (1961) reported turbidities of up to 900 ppm in Barataria Bay. However, high turbidities were of short duration, normally lasting a matter of hours. Shideler (1980) observed that the response of Corpus Christi Bay's circulation system and associated turbidity patterns to changing wind conditions was rapid (less than a few hours), mainly because of the shallowness of the bay. Shideler (1980) concluded that for Corpus Christi and Nueces Bays wind was the dominant forcing agent influencing turbidity in the bayhead sector, where it both generates waves that resuspend bottom sediment and regulates river inflow. Turbidity in the baymouth sector was mainly influenced by tidal-forcing effects from Aransas Pass tidal inlet (Shideler, 1980).

Turbidity fluctuates diurnally in Texas bays and estuaries, especially in late spring and summer (Conover, 1964). Conover (1964) observed that on clear, summer mornings, winds were often light until noon but would increase to 18 knots

in the afternoon. Winds above 14 knots could raise fine sediments from the bottom and hold them in suspension. On a summer day, afternoon turbidities would reduce by more than one-third the amount of light penetrating one meter below the surface compared with measurements taken in the morning. Turbidity was of little consequence in the winter except during stormy weather (Conover, 1964).

Spatial and temporal data on turbidities in Texas bays are generally recorded at only a few stations in a bay system and at variable times during the year. One of the best long-term data sets on turbidity is from measurements taken by Texas Parks and Wildlife (Martinez, 1969 through 1975). Martinez (1969 through 1975) took monthly turbidity measurements in most Texas bays, estuaries, and lagoons, and even though spatial coverage in some bay systems was sometimes minimal, measurements were taken at most stations throughout the year. For example, turbidity data were taken from 1969 through 1975 at two to three stations in upper Trinity Bay and two to three stations in Matagorda Bay near the Colorado River delta (table 10). Mean turbidities were generally higher in upper Trinity Bay (66 to 169 ppm) than in Matagorda Bay near the Colorado River delta (35 to 73 ppm). Minimum turbidities in upper Trinity Bay generally occurred in the summer or fall and maximum turbidities occurred in most months of the year. In Matagorda Bay near the Colorado River delta, minimum turbidities occurred generally in winter months and maximum turbidities in the summer or fall.

Turbidity and subsequent siltation are important environmental variables for pelagic organisms, benthic species, and seagrasses, because they can affect organism and plant distribution by: (1) decreasing the depth of the euphotic zone and thus decreasing primary production and productivity in seagrass beds; (2) increasing oxygen demand; (3) limiting suspension feeding; and (4) smothering benthic forms. Turbidity plays a principal role in regulating primary productivity rates (Flint and others, 1982). Odum and Wilson (1962) conclude that in order to produce maximum total

Table 10. Minimum, maximum, and mean turbidity measurements (ppm) for upper Trinity Bay and Matagorda Bay near the Colorado River delta. Compiled from Martinez (1969 through 1975).

Upper Trinity Bay				Matagorda Bay near the Colorado River delta		
<u>Year</u>	<u>Minimum</u>	<u>Maximum</u>	<u>Mean</u>	<u>Minimum</u>	<u>Maximum</u>	<u>Mean</u>
1969	22 (Sept)	211 (Aug)	66	5 (July)	300 (May)	73
1970	12 (Nov)	258 (May)	91	5 (Dec)	120 (Aug)	36
1971	44 (Feb)	114 (July, Nov)	76	0 (Jan)	250 (Mar)	40
1972	16 (Aug)	395 (Oct, Dec)	102	20 (Feb)	100 (July)	35
1973	5 (Apr)	395 (Mar)	169	8 (Feb)	90 (June)	45
1974	51 (Aug)	380 (Feb)	125	0 (Mar)	200 (Sept)	48
1975	22 (July)	390 (June)	154	4 (Nov)	75 (Oct)	38

photosynthesis in all the waters of Texas, measures for management should include reducing turbidity. Phytoplankton production per unit area is greater in shallow, clear bays, such as Redfish Bay and upper Laguna Madre, than in deep, turbid bays, such as Copano, Aransas, and Corpus Christi Bays (Odum and Hoskin, 1958). Increases in turbidity can reduce light penetration in seagrass beds and reduce productivity (Odum, 1963). Phytoplankton and seagrasses are dependent upon sunlight as the energy source for photosynthesis, and as the suspended sediment content of the water increases, the depth of light penetration decreases, resulting in decreased abundance and productivity of the phytoplankton and seagrasses. In turbid bays, much of the available light is being absorbed by clay particles before reaching phytoplankton plant cells (Odum and Hoskin, 1958). Increased turbidities can also reduce dissolved oxygen values (Sherk, 1971). The reductions have been attributed to an oxygen demand exerted by increased suspended sediment and the reduction of light penetration, which reduced photosynthetic oxygen production (Sherk, 1971).

Additions of inorganic nutrients into the water column may indirectly stimulate photosynthesis (Odum and Wilson, 1962). Where respiration is in excess of photosynthesis, inorganic nutrients accumulate stimulating photosynthesis. Thus a turbid mixture of organic and inorganic matter both interferes with photosynthesis by shielding light and stimulates photosynthesis by indirectly raising inorganic nutrient levels (Odum and Wilson, 1962).

Benthic standing crop is dependent on turbidity levels. Flood events may resuspend and transport sediments, increase turbidity, and cause a rapid decrease in the standing crop of phytoplankton, zooplankton, benthos, and nekton (Texas Department of Water Resources [TDWR], 1980 b). The time necessary for the recovery of the estuarine system after such an event is governed by variables such as season of year and temperature (TDWR, 1980 b). Gilmore and others (1976) reported that

benthic standing crop and species diversity in the Lavaca Bay estuary were significantly related to turbidity--species diversity was negatively correlated with turbidity and benthic standing crop was positively correlated with turbidity. The positive correlation between benthic standing crop and turbidity may be related to an increase in nutrients brought into the estuarine system by increased river inflows (Gilmore and others, 1976; Harper and Hopkins, 1976).

Turbidity is clearly an important factor in determining some molluscan communities. Suspension-feeding bivalves feed most effectively in relatively clear water (Loosanoff, 1962). Heavy concentrations of turbidity-creating substances can be lethal to bivalves, because they are not able to respire or feed normally (Loosanoff, 1962). The oyster, Crassostrea virginica, is able to tolerate turbid conditions with the assistance of well-developed mantle margins that mimic siphonal tubes, with discriminatory palps, with additional cleaning currents, and with more quick muscle fibres in the adductor muscle (McLusky and Elliott, 1981). Mackin (1962) reported that turbidities of up to 700 ppm produced no significant mortalities in experiments on Crassostrea. Ostrea edulis cannot tolerate conditions as turbid as those accepted by Crassostrea (McLusky and Elliott, 1981). Stora and Arnoux (1983) reported that when the ratio of freshwater to sediment volume discharged (the inverse of suspended sediment concentration) was lower than 1,500, mollusks can die. Even when the freshwater to sediment volume is less than 10,000, high concentrations of suspended matter may prove stressful to suspension-feeding mollusks and lower their resistance to other environmental factors (Stora and Arnoux, 1983).

Deposit-feeding organisms, by reworking muddy sediments, may create an unstable bottom and turbid water conditions that are unsuitable for suspension feeders. This kind of exclusion of one trophic group by another is termed trophic-group amensalism (Rhoads and Young, 1970). The feeding activities of deposit feeders

may result in high biogenic reworking rates producing a fluid, fecal-rich surface easily resuspended by waves or by low-velocity tidal currents. Instability of the bottom, resulting in high turbidities, may inhibit the growth of most suspension feeders and reduce infaunal diversity (Rhoads and Young, 1970; Aller and Dodge, 1974). Unstable bottoms can also cause high larval mortality for settled, suspension-feeding larvae (Aller and Dodge, 1974).

Effects of Marine Grasses on Sedimentation

Marine grasses or seagrasses form the basis of many estuarine ecosystems. The grassflat environment exhibits high biologic productivity (Odum, 1963) and has long been recognized as an important source of food and shelter for benthic macroinvertebrates, attachment sites for epifauna and epiphytes, nursery grounds for fishes, and direct food sources for some animals, including migratory waterfowl, sea urchins, green sea turtles, manatees, and a variety of herbivorous and juvenile fish (Cornelius, 1975; Stoner, 1980; Ward and others, 1980). The infauna of seagrass communities is generally much more diverse and dense than that of surrounding unvegetated areas (O'Gower and Wacasey, 1967; Santos and Simon, 1974; Orth, 1977; see sections on Biomass/Density and Species Diversity and Species Richness in this report).

Seagrasses and their diverse faunal communities serve as major sources of detrital material, dissolved organic matter, and nitrogen (Tenore, 1977; Kenworthy and others, 1982). The plants have the ability to remove both nitrogen and phosphorus from the water surrounding their leaves and roots (Short and Short, 1984), and through decomposition and retention of organic matter from animal and plant detritus, they make substantial contributions to the sedimentary nitrogen pool (Kenworthy and others, 1982). Increasing the nitrogen supply increases the productivity of plants and animals in an estuary (Jordan and Valiela, 1982).

Five species of seagrasses occur on the Texas coast, including Halodule beaudettei (= H. wrightii) (shoalgrass), Ruppia maritima (widgeongrass), Halophila engelmannii, Thalassia testudinum (turtlegrass), and Cymodocea filiformis (manateeegrass). Seagrasses occur in many areas along the bay margins of barrier islands and near tidal inlets but are much less abundant in the upper bays and near river deltas. Marine grasses are most abundant on the Texas coast in both upper and lower Laguna Madre (White and others, 1983; White and others, 1986). All five species found on the Texas coast occur in lower Laguna Madre (White and others, 1986). On the middle and upper coast, marine grasses are most abundant in the shallow waters of Redfish and Port Bays in the Corpus Christi area (White and others, 1983), Espiritu Santo Bay in the Port Lavaca area (White and others, in press b), and Christmas Bay in the Galveston area (W. Pulich, personal communication, 1988).

Species distribution is primarily a function of temperature, salinity, tidal regimes, water depth, and turbidity. Other factors affecting distribution include compactional and human-induced subsidence and eustatic sea-level rise. Hurricanes also greatly affect seagrass distribution and density.

Salinity is probably the most significant environmental factor controlling distribution. McMillan and Moseley (1967) and McMahan (1968) determined the general salinity tolerance of the various species. Halodule beaudettei was successfully grown in salinities of 60 ppt and flourished best in a salinity of 44 ppt. The growth rate of Cymodocea indicated it had the least salinity tolerance of the five seagrasses: growth terminated at a salinity of 45 ppt under controlled conditions. Chin (unpublished manuscript, 1977) citing McMillan and Moseley (1967), ranked the various grasses from most to least salt tolerant: Halodule, Thalassia, Ruppia, and Cymodocea. Studies of Halophila were inconclusive, but its salinity tolerance is suspected to be between that of Halodule and Cymodocea. Halodule and Ruppia are most able to

tolerate low salinities (Armstrong and Gordon, 1979); however, few marine grasses grow near areas of fluvial input.

Turbidity is a dominant feature in Texas bays and estuaries with organic- laden muds released in large quantity from the rivers, from dredging, from resuspension of the bottom sediment during storms, and from beach erosion (see Turbidity section in this report). Increases in turbidity can impair light penetration thus limiting primary productivity and reducing photosynthetic oxygen production. During dredging activities in Redfish Bay, Texas, Odum (1963) noted that primary production in turtlegrass beds decreased, and there was an imbalance of respiration over photosynthesis. This may have been caused by silts that were resuspended during dredging. However, diminished productivity was not permanent, since Odum (1963) reported that growth was exceptional the year following dredging. Odum (1963) speculated that dredging may have stimulated growth the year following dredging by releasing nutrients.

Clay particles in suspension tend to filter out the shorter wave lengths of light (Conover, 1964). There is evidence that terrestrial plants fail to develop normal reproductive organs if grown in sunlight without the higher frequency portion of the visible spectrum (Conover, 1964). The reproductive sterility of many seagrasses on the Texas coast may be related to the filtering effect of clay particles that absorb the shorter wave lengths of light (Conover, 1964).

Most seagrasses on the Texas coast grow best at depths of less than 2 meters (McMillan, 1984). Conover (1964) reported that Halodule wrightii and Thalassia testudinum grew best in water no deeper than 1.5 m in the Aransas and Redfish Bay area. Cymodocea filiformis grew well to depths of 2 m in lower Laguna Madre from Port Mansfield to Port Isabel (Conover, 1964). Light appeared to be the only factor that diminished with depth and may have limited seasonal growth in the deeper water. Species that extend into shallow water must also be tolerant to high

temperatures. Of the five species that occur on the Texas coast, Halodule beaudettei has the greatest heat resistance and can extend into shallow, in-shore sites that may be exposed to air at low tides (McMillan, 1984). Thalassia occurs at intermediate depths, and Cymodocea occurs in areas that are most likely to remain submerged. Halophila may occur in shallow areas as well as in areas in the depth range of Thalassia (McMillan, 1984).

Recent aerial photos show that the areal extent of seagrasses is increasing in some areas along the Texas coast (for example, along the bay margin of Mustang Island and the flood-tidal delta near Pass Cavallo) probably as a result of relative sea-level rise caused by both natural compactional and human-induced subsidence and eustatic sea-level rise (White and others, 1983; White and others, in press b). These factors tend to raise water levels, thereby decreasing the width of wind-tidal flats and leading to the more extensive and frequent inundation that favors the establishment of seagrasses (White and others, in press b).

In contrast to factors causing a gradual increase in the areal extent of seagrasses, hurricanes have contributed to their sudden decline. For example, grassflats that were once present along Matagorda Peninsula just southwest of the Colorado River delta, have virtually disappeared since Hurricane Carla in 1961 (White and others, in press a). The loss of marine grasses may be related to hurricane washover processes and the resulting deposition of sediments along the bayward margin of Matagorda Peninsula (White and others, in press a).

Marine grasses alter the sedimentation processes by increasing sedimentation rates, by concentrating preferentially the finer particle sizes, and by stabilizing the deposited sediments (Burrell and Schubel, 1977). Studies by Ward and others (1984) and Almasi and others (1987) indicate that sedimentation rates are substantially higher in seagrass communities than in unvegetated areas. Leaf and root structures effectively attenuate waves and baffle tidal currents leading to increased deposition

and consolidation of sediments and reduced resuspension (Ginsburg and Lowenstam, 1958; Wayne, 1974; Ward and others, 1984; Kemp and others, 1984). The reduction in resuspension is a direct function of plant biomass (Kemp and others, 1984). The efficiency of seagrasses in baffling the current flow and removing fine suspended particles depends primarily upon the leaf structure of the species and upon plant density (Burrell and Schubel, 1977; Harlin and Thorne-Miller, 1982). Ward and others (1984) estimated that sediment accumulated in seagrass beds at 2-3 mm/mo over a six month growing season. However, the fate of trapped sediment during plant senescence in the fall and winter is uncertain, especially in view of intense winter storms. The trapped sediments are probably held in place for at least 6-8 months, thus increasing water clarity during the most productive part of the year (Kemp and others, 1984).

Almasi and others (1987) reported that during the summer in the Indian River Lagoon in southeastern Florida, the mean weight percentages of mud trapped in artificial and natural seagrass (Thalassia) areas were always significantly greater than the amount of mud trapped in grass-free areas. In the summer, the average depositional rate was 4.96 grams of mud trapped per week in the Thalassia bed and 3.04 grams of mud trapped per week in the sandy, grass-free area; during the winter, these rates were 1.60 and 1.50 g /week, respectively. The amount of mud retained in grass-free and grassy areas was always greater in summer than winter as a result of the effects of runoff, bioturbation, wind speed, and direction: (1) Rain, surface runoff, and river discharge decrease during the winter, causing a reduction in the amount of suspended particles. (2) Increased biological activity during the summer causes destabilization of the bottom sediment, reduces erosion velocities, and thereby enhances resuspension. (3) Even though winter winds are strong, their direction has a short fetch and resuspension and settlement of sediment decreases during this time. Almasi and others (1987) argue that reduction of sediment accumulation in

seagrass beds during the winter is not a result of winter die-off that subsequently reduces the baffling effect of the grass blades. Partial winter die-offs do occur but not in their artificial seagrass plot. Therefore, the decreased winter mud flow in artificial grass beds must be due to decreased particle supply and can't be from a decreased baffle effect (Almasi and others, 1987).

Damming of rivers may result in a decrease in streamflow, which possibly could lead to inadequate flushing of pollutants from seagrass beds. Since the harnessing of the Rhone River in France has been completed, floods are rare and weaker than before (Peres and Picard, 1975). The accumulating clay minerals adsorb some anionic detergents when suspended in sea water and later desorb them in interstitial water (Peres and Picard, 1975). Comparison of Posidonia beds submitted to high sedimentation rates of clay in unpolluted and polluted areas on the French Mediterranean coast reveals that the disappearance of the beds only occurs in areas that were polluted by domestic sewage containing detergents, and this does not happen in areas where bottom currents are sufficient to prevent high sedimentation rates.

Seagrass sediments generally contain a higher percentage of silt and clay and organic carbon and are more poorly sorted than nonseagrass sediments (Lynts, 1966; Orth, 1973; Orth, 1977; Grady, 1981; Hoskin, 1983). Lynts (1966) found a close correlation between sediment size and density of turtle grass (Thalassia testudinum). Finer-grained sediments were found in areas of densest growth, about equal ratios of sand and mud in regions of dense to moderate growth, while the coarsest sediments were found in regions of patchy growth (Lynts, 1966). Hoskin (1983) found more silt in Thalassia sediments than in nearby sediments with bare sand. Orth (1973; 1977) reported that particle size and degree of sorting decreased and organic carbon increased from the edges of an eelgrass (Zostera marina) bed to an area where the eelgrass was most dense. Organic carbon content in sediments with Halodule and

Thalassia within the intertidal zone was 1.9 times higher than that of adjacent sand flats but much less than that in the subtidal seagrass meadows (Grady, 1981).

Trap Efficiencies of Bays

Much has been written with regard to various sources of sediments and whether bays and estuaries are permanent sinks for most sediments. There is evidence that many estuaries and bays may be relatively efficient in trapping sediments delivered to them. For example, Conomos and Peterson (1976) concluded that only about 6 percent of the annual riverborne load delivered under normal discharge conditions to San Francisco Bay was lost to the ocean. Biggs and Howell (1984) examined various estuaries using a modified version of a ratio formulated by Brune (1953) to measure the trapping efficiency of reservoirs. Brune (1953) compared the volume of a reservoir with the volume of incoming river water to estimate the amount of sediment that would be trapped, and therefore, the expected life of the reservoir. A reservoir large enough to hold one-hundredth of the annual volume of inflowing water could trap about half of the incoming sediment, and one large enough to hold one-tenth of the annual inflow could trap 80 to 90 percent of the sediment influx (Brune, 1953; Meade 1982). By applying this concept to estuaries, Biggs and Howell (1984) found that the sediment trapping ability of estuaries could be grossly predicted (fig. 82). The ratio C/I is a comparison of the volume (capacity; C) of the estuary to the total potential freshwater inflow (I), which is the product of the watershed area and the annual precipitation. The C/I ratios of various estuaries presented by Biggs and Howell (1984), and some Texas bays and estuaries are shown in table 11. These data suggest that Sabine Lake, which has a ratio similar to that of Mobile Bay, may have the lowest trapping efficiency of any Texas Bay at around 60 to 70 percent. However, a complicating factor in Sabine Lake is the Port Arthur and Sabine-Neches Canal network through which as much as 80 percent of




Figure 82. Sediment trap efficiency curves of Brune (1953) (see figure 37) applied to estuaries. Measured trapping efficiencies for selected estuaries are plotted in the figure: (1) Chesapeake Bay; (2) Rappahannock River; (3) Choptank River; (4) James River; and (5) Mobile Bay. (From Biggs and Howell, 1984.)

Table 11. Comparisons of predicted trapping efficiencies of other estuaries with those in Texas. Data for estuaries outside of Texas from Biggs and Howell (1984); data on estuarine volume and inflows for Texas from Armstrong (1982), based on data from Texas Department of Water Resources (1980 a, b and 1981 a, b, c), and Currington and others (1966). Percent sediment trapped estimated from Brune (1953).

Estuary	Volume (km ³) (C)	Inflow (km ³ /yr) (I)	C/I	Predicted Trapping Efficiency (percent)
Narragansett Bay, RI	2.4	6.24	0.4	95 ± 5
Long Island Sound, NY	60.8	46.2	1.3	97 ± 3
New York Bay	2.3	41.8	0.05	77 ± 10
Delaware Bay	19.4	36.3	0.5	98 ± 3
Chesapeake Bay	80.5	121	0.7	98 ± 3
Patuxent River	6.1	2.42	2.5	> 99
Pamlico Sound, NC	0.9	14.3	0.06	80 ± 10
Apalachicola Bay, FL	0.4	61.6	0.006	30 ± 15
Mobile Bay, AL	3.2	150	0.02	61 ± 12
Barataria Bay, LA	0.3	6	0.05	76 ± 10
San Francisco Bay, CA	2.5	112	0.02	61 ± 12
<u>Texas Estuaries</u>				
Sabine-Neches	0.33	16.1	0.02	61 ± 12
Trinity-San Jacinto	2.9	12.1	0.24	95 ± 5
Lavaca-Tres Palacios	2.1	3.6	0.59	98 ± 3
Guadalupe (San Antonio)	0.75	2.8	0.27	95 ± 5
Nueces (Corpus Christi)	1.15	0.84	1.37	97 ± 3

the fresh-water discharges of the Sabine and Neches River flow to the Gulf of Mexico and thus bypass Sabine Lake (Ward, 1973, cited in Gosselink and others, 1979). Other Texas bays, based on C/I comparisons characterizing other estuaries, appear to have trapping efficiencies exceeding 90 percent. Maps of shoaling rates in Texas estuaries (Shepard, 1953) indicate that most of the river-derived suspended sediments may be deposited in the upper half of the bays. However, estimates by the Corps of Engineers (USACE, 1942) of the disposition of fluvial sediments in the Trinity-Galveston Bay system indicate that about 40 percent of the riverine sediments pass through the bay and into the Gulf of Mexico. Wilkinson and Byrne (1977) concluded that much of the fluvial sediment delivered into Lavaca Bay was transported through the bay and into Matagorda Bay. The USDA (1972) estimated that 60 to 70 percent of the total sediment delivered to Texas bays and estuaries is retained, while the remainder is transported into the Gulf.

In a study of the expected delta progradation at the mouth of the Colorado River when it is diverted into Matagorda Bay (USACE, 1981), van Beek and others (1981) concluded that about 7 to 8 percent of the incoming suspended load (there is little or no bed load because of a sediment trap upstream) would be retained in the area of the delta and the remainder would be carried out of the bay (eastern arm of Matagorda Bay).

Holmes (1982) postulated that during frontal passage a substantial amount of suspended sediment from the bays and estuaries along the central and upper Texas coast is transported out of the bays through tidal inlets and is deposited on the Continental Shelf (Holmes' model is discussed more fully below). White and others (1983; 1985) found some supporting evidence from trace metal concentrations in sediments on the shelf suggesting that the sediments were derived from adjacent bays. Major flood events that apparently transport the largest volumes of sediments may push substantial amounts of sediment out to sea, at least in Gulf coast estuaries.

Importance of Episodic Events on Fluvial-Estuarine Sedimentation

The importance of extreme events such as hurricanes and other storms in transporting fluvial sediments into estuaries has been reported by several investigators (for example, Schubel and Meade, 1977; Gross and others, 1978; Milliman and Meade, 1983).

Only a few river sediment stations have been in operation long enough to have documented the extreme events that are so important in the introduction of sediment: events such as the hurricane flood of August 1955 when the Delaware River carried more sediment past Trenton in two days than in all five years combined in the mid-1960's drought; or the three days in December 1964 when the Eel River in northern California transported more sediment than in the preceding eight years; or the week following Tropical Storm Agnes in June 1972 when the Susquehanna discharged 20-25 times as much sediment as during the previous year. Events of this magnitude occur only rarely--a few times a century at most--but their importance to estuarine sedimentation is so great that programs should be designed to record their effects when and where they do occur. (Schubel and Meade, 1977: p. 205)

In a study of sediment loads of the Susquehanna River delivered to Chesapeake Bay, Gross and others (1978) found that of 50 million tons of suspended sediment discharged by the river during a ten year period (1966-1976), approximately 80 percent was discharged in only 10 days. Two extreme events, Tropical Storm Agnes (1972) and Hurricane Eloise (1975) were responsible. Apparently the sources of

much of the contributed sediment were reservoirs along the Susquehanna River that were flushed during the events (Gross and others, 1978; Meade, 1982). The sediment from the reservoirs represented about 20 years of storage (Meade, 1982). Meade (1982) cautioned about extrapolating these findings to other areas due to the fact that the reservoirs are narrow, and an independent check needed to be made of the observations.

Van Heerden and Roberts (1980) and van Heerden and others (1981) reported evidence of the importance of annual flood events in the subaerial growth of the Atchafalaya Delta in Louisiana. Larger increases in area occurred following floods during the 1970's.

The importance of episodic flood events has also been reported in Texas coastal fluvial-estuarine areas. In a study of the suspended load of the Trinity River in 1965, Rice (1966) determined that approximately 80 percent of the year's sediment discharge occurred during two major floods; the larger flood, which carried 1,770 acre-feet of suspended sediment, accounted for 59 percent of the annual total. The sediment load carried by the Colorado River near Matagorda, Texas, during two floods in 1979 totalled 970 acre-ft, of which about 2 percent was bedload (Welborn and Andrews, 1979). Although the total sediment load of the river near Matagorda for 1979 is not known, estimates by the Corps of Engineers indicate an annual average of about 1,350 acre-ft (USACE, 1977), which is supported by van Beek and others (1981). The volume of sediment discharged by the two floods monitored by Welborn and Andrews (1979) represents about 70 percent of this annual average. The relationships between discharge and sediment load along the Colorado at Columbus and Matagorda were plotted by van Beek and others (1981) (fig. 83 A). In analyzing sediment transported by the Colorado River, van Beek and others (1981) found that sediment load frequency graphs indicated that 81 percent of the total




Figure 83. Sediment transport by the Colorado River. (A) Comparison of discharge and sediment load at Columbus (for 1957-1967) with that at Matagorda (May and June 1979); (B) Sediment load frequency of occurrence; (C) Tons of suspended load per class and percent of total suspended load per class by frequency of occurrence. (From van Beek and others, 1980.)

annual sediment load was transported by the five percentile classes representing the highest river discharge (figs. 83 B and 83 C).

Scott and others (1969) and McGowen (1970) reported on large contributions of sand to a fan delta on the margin of Nueces Bay that resulted from extensive rains accompanying Hurricane Beulah. The layer of sand deposited was more than a meter thick in many places and composed a volume of approximately $2.5 \times 10^5 \text{ m}^3$ (Scott and others, 1969). Apparently the sediment was deposited in less than a week (possibly a day) while 60 cm of rain fell within the 80 km^2 drainage basin of Gum Hollow Creek, the stream that supplies the fan delta with sediment.

The amount of suspended sediment that passes through an estuary and out onto the continental shelf during extreme flood events is not easily determined because of the difficulty of monitoring estuarine systems during such events. In examining the effects of Tropical Storm Agnes on the Rappahannock Estuary (in the Chesapeake Bay system), Nichols (1977) concluded that 90 percent of the total sediment influx was trapped within the estuary; deposition of sediment occurred mainly in the zone of the turbidity maximum (at the interface of the marine salt wedge and the inflowing fresh water). He further postulated that in large estuaries such as Chesapeake Bay, most flood events are not able to push the salt intrusion out to sea and that floods might actually enhance the ability of the hydraulic regime to trap sediments by intensifying the stratification and convergence between the fresh-water and salt-water zones. He further concluded that where water depths are increased by dredged channels, entrapment is compounded. Sediment deposition in the estuary as a result of the tropical storm decreased seaward from a thickness of 7.5 mm to 0.5 mm; deposition in a channel in the area was extreme, totalling about 40 percent of the average annual deposition (Nichols, 1977).

Texas bays are considerably different (shallower, lower tides, more homogeneous with respect to vertical salinities, more protected by barrier islands at their mouths)

from Chesapeake Bay (Nichols, 1977). The trapping efficiencies of bays and estuaries along the Gulf coast may be significantly less during storms. As mentioned briefly above, Holmes (1982) has proposed that bay systems along the northern Texas coast, including Matagorda Bay, are staging areas for the transport of fine-grained sediments out of the bays and southward onto the Outer Continental Shelf. The proposed delivery events are frontal systems that are common occurrences in the late fall, winter, and early spring. As noted by Holmes (backed by observations of other researchers) as a frontal system approaches the coast, southeasterly (onshore) winds increase in strength, pushing Gulf water into the bays and estuaries; associated waves and currents stir up sediments and increase bay turbidities. As the front passes, usually accompanied by significant amounts of precipitation, the winds shift to the north. The combination of high freshwater inflows and offshore winds acts to flush the swollen bay and estuarine waters loaded with sediments, out of the tidal passes and onto the continental shelf (fig. 84). The outflow may last for days as a result of the hydraulic head and flooding (Holmes, 1982).

Hurricanes and tropical storms can have even more impact with respect to flushing sediments from the bay-estuary-lagoon system. Isphording and others (1987) reported that more than 83 million tons of sediment, scoured from the bottom of the Apalachicola Bay system during the passage of 2 hurricanes in 1985, were carried into the Gulf of Mexico by high velocity currents. According to the authors, the removal of this large amount of sediment significantly extended the life of the bay, which is being filled with sediment. Hayes (1967) mapped a blanket of sediment more than 9 cm thick, locally, on the inner shelf that he attributed to deposition by Hurricane Carla surge waters discharging back to the Gulf (there is disagreement as to the source of this sediment, however; Robert A. Morton, personal communication, 1988). Hurricanes also dump considerable amounts of sediment (as manifested in coarse-grained washover-fan deposits at the termini of washover channels) into the bays and




Figure 84. Model depicting possible effect of frontal passage on the transportation and distribution of fine-sediment from the bay-estuary-lagoon system to the South Texas shelf. (From Holmes, 1982.)

estuaries. The net movement of fine-grained sediments is difficult to quantify because of the extreme conditions in which measurements have to be made. However, one would expect that extensive aftermath rainfall that often accompanies hurricanes (for example Hurricane Beulah; Scott and others, 1967) adds considerable amounts of fine-grained sediment to the estuarine system and contributes to the expulsion of sediment-laden water out of the bay. Again, the quantity of suspended sediment brought into the bays during storm surge flooding compared with how much is taken out during ebb flows has not been determined.

In summary, the sand (bed load) delivered into the bays and estuaries by rivers is deposited near the river mouths and trapped because of diminishing current velocities. Sand may be reworked along the margin of the bay and even transported along shore by littoral currents, but it is effectively trapped in the system unless the river fills the estuary with sediment and discharges directly into the Gulf. The fate of suspended sediments (silt and clay) is less clear. Apparently the largest amount of the fluvial sediment is deposited in the upper reaches of the estuary, however, the volume (or percentage of the total supplied) that is transported out of the estuary and into the Gulf cannot presently be confidently quantified.

Are Texas Bays Getting Shallower or Deeper?

Bay-estuary-lagoon systems have been characterized by many researchers as efficient sediment traps (Kennedy, 1984). As noted by Biggs and Howell (1984), Emery and Uchupi (1972) concluded on the basis of studies of suspended sediment discharged by rivers (excluding the Mississippi) along the Gulf and Atlantic coasts, that on average, if sea level remained unchanged, the bays and estuaries should be filled in approximately 9,500 years (or faster if other sources of sediments are included). Average rates of deposition in the bay-estuary-lagoon system, according to Emery and Uchupi (1972), are about 70 cm/100yrs, or 7 mm/yr. But this rate is

close to the average rate of sea-level rise (80 cm/100 yrs) over the past 5,000 years, which suggests a balance between shoaling and deepening (Emery and Uchupi, 1972).

As noted in the introductory section on the origin of bays along the Texas coast, the rate of global sea-level rise over the past few thousand years has varied. The average rate of rise of 8 mm/yr for the past 5,000 years (Emery and Uchupi, 1972) is more than the current global sea-level rise of between 1 and 1.5 mm/yr (Gornitz and others 1982). However, when subsidence (Swanson and Thurlow, 1973) is considered in conjunction with global rise, relative sea-level rise averages about 11 mm/yr along the central and upper Texas coast; the rates are considerably higher in areas of human-induced subsidence in the Houston area (fig. 9).

If the average rate of relative sea level rise is 11 mm/yr in the different bays, are bays along the Texas coast shoaling (becoming shallower from sediment deposition) or are they deepening (from subsidence and scouring at rates higher than sediment deposition)?

Using unpublished bathymetric data collected by the Coast and Geodetic Survey in the bay-estuary-lagoon system of the Texas coast, Shepard (1953) compared more than 20,500 points where soundings were made during the last century (1852-1875) with those made during the present century (1934-1935). Although there is much variation from bay to bay and within a bay (fig. 85), he detected, overall, a trend toward bay shoaling at an average rate of about 1.26 ft/century (3.8 mm/yr). Because the silt load of Texas streams is too high for this rate of shoaling, he concluded that the bays must be submerging at a rate of about 1.8 ft/century (5.4 mm/yr) based on the trend set by Texas coastal tide gauge records (Marmer, 1952). Thus, Shepard (1953) estimated the total rate of average sedimentation (submergence + shoaling) in the Texas bays to be 3 ft/century, or 9 mm/yr.

In a comparison of the silt load of Texas streams (from river stations as close to the coast as allowed by the sampling network), Shepard (1953) found a correlation

DEPTH CHANGES IN TEXAS BAYS

ACRE FEET ARE ESTIMATES OF TRANSPORT OF "SILT" PER 100 YEARS BY STREAMS AND OF SOLID FILL IN BAYS




Figure 85. Depth changes in Texas bays. Boxes for the different bays include shoaling (+) or deepening (-), shoaling rate per century in feet, approximate volume in acre-feet of sediment per century represented by shoaling rate, and depth of main parts of bay in feet. Acre-feet for rivers represents 100 years at rates for 1950-1951. (From Shepard, 1953.)

between stream sediment load and bay shoaling rates (or total bay fill) projected over a century. For example, the Trinity and San Jacinto Rivers, together, transport the largest amount of sediment and the Galveston-Trinity Bay system, into which they discharge, has the largest amount of bay fill (table 12). The total silt load transported by Texas streams, however, was too high for the amount of bay fill (table 12). This disparity was attributed in part to deposition of sediment in stream floodplains below the stream gauging stations, or behind dams, although Shepard noted that all the large dams were constructed near the end of period between bathymetric surveys. However, he postulated that other sources of sediments probably equalize the stream sediment load not reaching the bays, and suggested that the difference between the silt load carried by the streams and bay fill was an indication of submergence in all bays but Corpus Christi where bay shoaling and stream sediment load were more balanced. As noted previously, this theory of submergence along the Texas Gulf coast was supported later by Swanson and Thurlow (1973). The Corpus Christi Bay system is apparently undergoing subsidence also, as reflected by tide gauge records at Aransas Pass (Swanson and Thurlow, 1973).

Comparing the average bay depth of 7 ft (2.1 m) with his average shoaling rate of 1.26 ft/century (3.8 mm/yr), Shepard concluded that most of the Texas bays should be filled in less than 1000 years. However, evidence from cores and borings in bay floor sediments suggest that conditions similar to present have persisted for thousands of years and that sea-level rise and subsidence could maintain the bays far longer than present shoaling rates indicate. Shepard (1953) also recognized the high water content of recently deposited sediments and the fact that compaction after burial would produce a considerably thinner sedimentary sequence. Shoaling rates of Texas bays and estuaries are not significantly different from sedimentation rates reported in other bay-estuary systems; however, they are lower than rates reported for Apalachicola and Mobile Bays along the Gulf coast (table 13). lower

Table 12. Relation of bay fill to suspended sediment load of entering river (from Shepard 1953).

Bay	Solid Fill (Acre-ft per century) ¹	River and Station ²	"Silt" load (Acre-ft per century)	Ratio river load and bay fill ³
Corpus Christi	31,590	Nueces, Three Rivers	49,400	1.56
San Antonio	23,070	Guadalupe, Victoria & San Antonio, Goliad	77,500	3.36
Lavaca	4,155	Lavaca, Edna	10,800	2.61
East Matagorda	39,600	Colorado, San Saba	292,000	7.37
Galveston	86,400	Trinity, Romayor & San Jacinto, Huffmann	463,000	5.36

1 Reduced to 30 % of volume because of estimated water content.

2 From Texas Board of Water Engineers (1952).

3 The river "rafts" between 1865 and 1926 (Price and Gunter, 1942) probably led to the diversion of an unusually large amount of sediment to the flood plain, thus increasing this ratio.

Table 13. Bay-estuary sedimentation and net deposition rates.

<u>Bay-Estuary System</u>	<u>Rate (mm/yr)</u>	<u>Source</u>
Chesapeake Bay	3.7	Biggs (1970)
Chesapeake Bay (head of bay)	3-4	Schubel and Carter (1984)
Great Sound	1-5	Thorbjarnarson and others (1985)
Raritan	1.5-3.4	Renwick and Ashley (1984)
Apalachicola Bay	6.7	Isphording (1986)
Mobile Bay	5.6	Ryan and Goodell (1972)
<u>Texas Bays</u>		
Galveston-Trinity	4.3	Shepard (1953)
	3.6	Rehkemper (1969)
Lavaca	1.4	Shepard (1953)
Matagorda	-0.7	Shepard (1953)
San Antonio	3.7	Shepard (1953)
Corpus Christi	2.6	Shepard (1953)

Contribution of Fluvial Sediments to Bay Sedimentation

Rivers are the primary source of sediments deposited in the bay-estuary-lagoon system, at least from Corpus Christi Bay northward. The importance, in terms of volume contributed, of fluvial sediments is reflected (in addition to table 12) by Shepard's maps of depth changes for the various bay systems. The map of the Galveston-Trinity Bay system is a good example (fig. 86). The decrease in water depth in the upper half of Trinity Bay compared to other parts of Trinity Bay and Galveston Bay reflects sediment deposition from the Trinity River. Maps of other Texas Bays show similar trends, although of lesser relative magnitude (except for the Colorado River).

The Filling of Lavaca Bay--An Example

Byrne (1975) and Wilkinson and Byrne (1977) reconstructed the geologic depositional history of Lavaca Bay. As noted in the section on the origin of Texas Bays, valleys that were eroded during the most recent sea-level lowstand were flooded by sea-level rise during the present interglacial stage. The depth of the axis of the incised Lavaca Bay valley (near the mouth of Lavaca Bay) is about 30 m (90 ft) below current mean sea level (Wilkinson and Byrne, 1977; Wright, 1980) (fig. 87). Deposition of sediment in the valley has taken place over the past 10,000 years, and consists of (from the base of the valley upward) (1) fluvial sand, (2) deltaic sand and muddy sand, and (3) bay-estuarine mud (with local oyster shells) (fig. 88). At the head of the bay near the mouth of the Lavaca River, fluvial deltaic sand has prograded over the estuarine muds (fig. 88). This sequence is generally similar to that recorded by Rehkemper (1969) for the Trinity-Galveston Bay valley-fill sequence with a few exceptions (including local, thin peat beds at the top of the fluvial-deltaic sequence in the Trinity-Galveston Bay depositional sequence).

Despite a 25-m rise in sea level, Lavaca Bay has remained a relatively shallow estuary with rates of sedimentation only slightly exceeding flooding rates throughout





Figure 86. Galveston Bay depth changes determined by comparing bathymetric surveys ranging from 1852 to 1867 with surveys in 1934. Highest rate of fill is near the Trinity River delta. (From Shepard, 1953.)



its depositional history (Wilkinson and Byrne, 1977). Wilkinson and Byrne (1977) constructed a paleobathymetric curve (of the past 10,000 years) representing the depth of the axis of the Lavaca Bay valley and its changes through time (fig. 89). The curve indicates that initial sedimentation (aggradation) rates were high but gradually slowed. This change was attributed to an ever-widening valley, which required larger volumes of sediments to deposit a given vertical unit of sediment (Wilkinson and Byrne, 1977). Water depths were relatively stable, but gradually decreased until about 3,000 years ago when maximum depths were about 3 m. This gradual shoaling indicates that rates of sedimentation exceeded sea-level rise (Wilkinson and Byrne, 1977).

To determine an argillaceous-sediment (predominantly clay-sized material) budget for Lavaca Bay, Wilkinson and Byrne (1977) analyzed the contributions of the various sources of sediments. As mentioned previously, they found the rivers to be the most important source of sediment, supplying about 73 percent of the total. Erosion of bay shorelines, characterized by Pleistocene bluffs, was a secondary but significant source, supplying 24 percent. The remainder was attributed to sediment contributions from Matagorda Bay.

Comparison of geologic sedimentation rates with the historic rate estimated by Shepard (1953) for the Lavaca Bay system indicates historic rates are higher. Wilkinson and Byrne (1977) suggested the higher rates over the past several decades may be due to historic agricultural practices (rice farming) in the drainage basin. Price and Gunter (1943) also reported increases in rates of bay-lagoon siltation in some areas along the Texas coast as a result of human activities.

Trends in Relative Sea-Level Rise and Fluvial-Estuarine Sedimentation

Data from Shepard (1953) and Wilkinson and Byrne (1977) suggest that Texas bays are becoming shallower due to infilling, but what are the more recent trends




Figure 88. Cross section down Lavaca River valley showing sediments that have accumulated in Lavaca Bay above the Pleistocene surface. (From Wilkinson and Byrne, 1977.)




Figure 89. Paleobathymetric curve over the past 10,000 years for Lavaca Bay. Sea-level positions are maximum values taken from various authors. Positions of the bay bottom are based on radiocarbon dates. Past maximum water depths of Lavaca Bay are represented by the vertical distance between the two curves. Radiocarbon data from Byrne 1975. (From Wilkinson and Byrne, 1977.)

along the Texas coast with regard to subsidence rates and reductions in sediment supplied by major rivers?

Corpus Christi-Nueces Bay System

The bathymetric surveys used by Shepard (1953) generally encompass a period of time that preceded placement of dams along the major rivers. He did recognize that after construction of a dam along the lower Nueces River (Corpus Christi Lake), contribution of fluvial sediment to Corpus Christi Bay ceased to be important in 1930. Based on suspended sediment loads of the Nueces River (at Three Rivers, upstream from Lake Corpus Christi) Shepard determined that the river transported about 494 acre-feet a year. More recent measurements of suspended sediment loads along the Nueces River (from a station downstream from Lake Corpus Christi) show an annual rate of less than 19 acre-feet a year (Texas Water Development Board and Water Commission, published and unpublished records, 1962-1984), or about 4 percent of the amount (translated into an annual mean) reported by Shepard (1953) (table 12).

Shepard noted that of all the bays he investigated, the Corpus Christi Bay system had the closest balance between sediment supply and bay fill (table 12). With the large decrease in sediment supply from the Nueces River, it seems likely that bay sedimentation rates have fallen behind rates of relative sea-level rise, and Corpus Christi Bay is becoming deeper through time. The stream silt load value used by Shepard (1953) is based on data from the late 1920's to the early 1950's (fig. 31) and probably reflects larger than natural sediment loads due to human development in the drainage basin. On the other hand, Price and Gunter (1943) reported that increases in bay siltation in some areas of south Texas were probably due to a combination of natural (droughts) and human events (clearing of the land for agricultural purposes and overgrazing) beginning in the 1880's. Soil conservation measures implemented by the Soil Conservation Service before the 1950's may have caused the average annual stream silt-load used by Shepard (1953) to be less than that carried

by the Nueces River between 1868 and 1934 (the period between the bathymetric surveys) when land use practices and droughts probably caused higher sediment yields. If this is the case, then the balance between stream silt load and bay fill, under more natural conditions, may have been even closer than reported by Shepard (1953) (table 12).

Much of the sediment delivered by the Nueces River is apparently deposited in shallow Nueces Bay (Shideler, 1980). Bay bathymetry, however, has also been affected by excavation of oyster shell as an economic resource. According to Texas Parks and Wildlife Department Shell Dredger's Reports, more than 2.5 million yd³ of shell was dredged from Nueces Bay from 1969 to 1975 (Kier and White, 1978). This is equivalent to about 1,550 acre-ft of material, or about a 47-year supply of suspended sediment from the Nueces River (based on an average annual supply of 32.7 acre-ft/yr from 1969-1975, measured at the Mathis station below Lake Corpus Christi; Mirabal, 1974, and Dougherty, 1979). Dredging of shell material apparently increased the depth of Nueces Bay at least locally.

San Antonio-Espiritu Santo Bay System

Comparison of depth changes between 1873-74 and 1934-35 for San Antonio and Espiritu Santo Bays indicates shoaling rates of 36.9 cm/100 yrs (1.23 ft/100 yrs) and 12.3 cm/100 yrs (0.41 ft/100 yrs), respectively (fig. 85; Shepard, 1953). The considerably higher rate in San Antonio Bay is reflective of the fluvial sediment delivered by the Guadalupe and San Antonio Rivers. Shepard (1953) reported that, together, these rivers supplied about 775 acre-ft of suspended sediment per year based on measurements at Victoria and Goliad (fig. 85). More recent measurements indicate that the annual volume of suspended sediments has decreased slightly (table 14), but not at the magnitude documented for other Texas Rivers such as the Trinity, Brazos, Colorado, and Nueces Rivers (figs. 21-26 and 31-32).

Table 14 . Annual average suspended load of the San Antonio and Guadalupe Rivers combined (Goliad and Victoria measuring stations; the period of record for the Guadalupe River station at Victoria began in 1946). Data compiled from Stout and others (1961), Adey and Cook (1964), Cook (1967, 1970), Mirabal (1974), Dougherty (1979), and unpublished records available from the Texas Water Development Board (1976-1984).

PERIOD (water years)	AVERAGE ANNUAL LOAD (acre-feet)
1943-1951	751.9
1951-1961	657.5
1961-1971	523.5
1971-1984	669.4

In San Antonio Bay, Shepard (1953) noticed that the greatest shoaling had occurred in the upper bay. In the central bay, less shoaling had occurred on the eastern half compared to the western half, but this trend was reversed in the lower part of the bay (gulfward of the Intracoastal Waterway) where less shoaling had occurred on the western half relative to the eastern half.

In contrast to San Antonio Bay, Espiritu Santo Bay has no important fluvial source of sediments, and the depths remained relatively constant with the exception of shoals at each end of the bay (Shepard, 1953). Although, overall, there was a slight decrease in depth in Espiritu Santo Bay (0.41 ft/100 yrs, which is equivalent to a shoaling rate of 1.2 mm/yr--considerably lower than the 3.7 mm/yr for San Antonio Bay), Shepard (1953) reported that at about 28 percent of the 905 notations where depths were compared, a slight deepening had occurred. This percentage is high compared to most other Texas bays.

Since the analysis by Shepard, depths of San Antonio Bay have been affected locally by the excavation of oyster shell. Records of the Texas Parks and Wildlife Department show that more than 36 million yd³ of shell material was mined between 1969 and 1983 (White and Morton, 1987). This is equivalent to about 22,336 acre-ft of material, which is a volume equal to about 33 years of suspended load supplied by the Guadalupe and San Antonio Rivers at the average rate of 670 acre-ft/yr (based

on the period 1971-1984). This volume is also very close to the amount of bay fill per century (23,070 acre-ft) estimated by Shepard (1953) for San Antonio Bay. Depths in the bay have undoubtedly been increased locally by shell removal.

Comparison of bay shoaling to sediment supplied by the San Antonio and Guadalupe Rivers by Shepard indicates that the ratio of river load to bay fill is about 3.4 to 1 (table 12). Differences in these volumes are attributed to deposition of sediments up stream in channels and flood plains, and transportation of sediments out of the bays into the Gulf; however the principal reason for the difference is attributed to subsidence or submergence (Shepard, 1953).

Much of the fluvial sediment delivered by the rivers is apparently being deposited in Mission Lake (fig. 64); when the lake is filled, the Guadalupe River delta could resume its progradation down the axis of San Antonio Bay (it should be noted, however, that with the exception of progradation of the Traylor Cut subdelta, and shoreline erosion, little change has been noted in the configuration of the Guadalupe delta since the mid-1800's (Shepard, 1953; Donaldson and others, 1970; and White and Morton, 1987). As mentioned previously, Donaldson and others (1970) reported that when the Traylor Cut subdelta began to prograde into Mission Lake in 1935, the lake was about 1.5 m (5 ft) deep, and by 1965 it was only 0.6 m (2 ft) deep. Citing the shoaling data presented by Shepard and Moore (1960), Donaldson and others (1970, p. 130) state that "the lower part of San Antonio Bay most likely will fill up and become a tidal flat before the Guadalupe delta reaches the barrier island". These researchers predict that if the many variables controlling sedimentation continue without drastic change, the bay will eventually (probably with reference to a geologic time frame) be filled and the river will transgress the barrier island and discharge into the Gulf.

According to a special report by the Bureau of Reclamation (USDI, 1978) on the San Antonio-Guadalupe Rivers basins, Childress and Bradley (1975) hypothesized that

the amount of fluvial sediments being delivered to San Antonio Bay estuary is in equilibrium with the bay system and that changes in the historic load would create an imbalance, which would reduce bay productivity and nutrients, and increase erosion.

Shoaling rates determined by Shepard (1953) for San Antonio Bay (3.7 mm/yr) are similar to the average (3.8 mm/yr) for all the bays that he analyzed. Assuming the average subsidence rate was 5.4 mm/yr (Marmer, 1952), then the sedimentation rate was about 9 mm/yr, which is similar to the average bay sedimentation rates determined by Shepard (discussed at the first of this section). If the more recent average rate of relative sea level rise in San Antonio Bay falls between the rates presented by Swanson and Thurlow (1973) for Aransas Pass (12.8 mm/yr) and Freeport (11.2 mm/yr) (eustatic sea level rise would add another 1.2 mm to these values), then the rate of sedimentation may not be keeping pace with submergence, and the bay, overall, could be getting deeper at a possible rate of 2 to 4 mm/yr. As noted in a preceding paragraph, however, if Mission Lake fills with sediment, then fluvial sediment loads may be redirected into San Antonio Bay and thus, increase sedimentation rates.

Lavaca-Matagorda Bay System

Wilkinson and Byrne (1977) suggest that sedimentation rates in Lavaca Bay have possibly increased in historic time. This conclusion was reached by comparing Lavaca Bay sedimentation rates during geologic time (based on the total volume of argillaceous sediment in the bay and its accumulation over the past 8,000 to 10,000 years) with more recent historic sedimentation rates (net rate of 1.4 mm/yr) reported by Shepard (1953). The geologic rate is about half the historic rate (Wilkinson and Byrne, 1977). Although several reasons may account for the difference, including rounding of the historic sounding data to the nearest one-half and one-fourth of a foot, and the fact that the recent rates reported by Shepard are for uncompacted sediments, Wilkinson and Byrne concluded that compaction alone could not account

for the difference and that the historic rate was higher possibly because of agricultural development in the drainage basin.

Comparing the silt load volumes transported along the Lavaca River (Edna station) with the volume of fill in Lavaca Bay, Shepard (1953) found that the stream silt load exceeded bay fill by a factor of about 2.5 (table 12), which he attributed mostly to submergence or subsidence. Over the geologic time frame, Wilkinson and Byrne (1977) decided that much of the sediment supplied by the streams discharging into Lavaca Bay was transported into and deposited in Matagorda Bay. In fact, they estimated that the total amount of fill in Lavaca Bay represented only 30 percent of all the sediment available from different sources; the remainder was transported into Matagorda Bay. It is probable that a certain amount reached the Gulf through processes described by Holmes (1982). Shepard (1953) found that during historic time, Matagorda Bay, with the exception of its eastern arm, which was supplied with sediments from the Colorado River, actually became slightly deeper, overall, at an average rate of about 0.7 mm/yr (fig. 85). Recent stream silt-load volumes for the Lavaca River (Edna station) are similar to that reported by Shepard (108 acre-ft/yr based on records prior to 1952; more recent rates are approximately 100 acre-ft/yr-based on silt loads from published and unpublished records of the Texas Water Development Board and Texas Water Commission, for the period 1950-1984). The Navidad River and Garcitas Creek are additional sources of fluvial sediments entering Lavaca Bay. The Navidad River source (quantified below) has undoubtedly declined with the completion of Lake Texana in 1980.

Although the upper half of Lavaca Bay decreased in depth (indicating net sedimentation) (Shepard, 1953), the lower half and most of Matagorda Bay had no change (fig. 90), which suggests an equilibrium in rate of sedimentation and submergence-erosion. Average contribution of the Navidad River, based on data from 1963 to 1984 collected at the Hallettsville station, was about 26.4 acre-ft/yr ($3.26 \times$

$10^{10} \text{ cm}^3/\text{yr}$). This rate of sediment delivery is approximately 20 percent of the geologic rate of sedimentation (annually) in Lavaca Bay, and about 10 percent of the historic rate (Wilkinson and Byrne, 1977; Shepard, 1953). However, Greiner's (1982) data on sediment yield indicate the Navidad River basin is a much larger source of sediment than measurements at the Hallettsville station indicate (about 75 percent of the drainage basin apparently lies below this station). Calculations of sediment volumes based on Greiner's sediment yield factors show that the respective yields of the Lavaca River and Navidad River drainage basins are 272.6 acre-ft/yr ($3.36 \times 10^{11} \text{ cm}^3/\text{yr}$) and 783.7 acre-ft/yr ($9.60 \times 10^{11} \text{ cm}^3/\text{yr}$). These volumes suggest that the Navidad River basin produces about 3-times as much sediment as the Lavaca River basin. The Garcitas Creek area adds another 77.9 acre-ft/yr ($9.61 \times 10^{10} \text{ cm}^3/\text{yr}$) (Greiner, 1982).

It is not possible to predict the exact effect on future depth changes in the Lavaca-Matagorda Bay system as a result of loss in sediment supplied by the Navidad River, but the sedimentation rate will likely decline and possibly fall behind relative sea-level rise in parts of Lavaca Bay. Using data from Shepard (1953), Wilkinson and Byrne (1977), and Greiner (1982), rough approximations of possible changes in the sedimentation rate can be estimated.

The net sedimentation rate presented by Shepard (1953) for Lavaca Bay is 0.14 cm/yr. Considering that the area of Lavaca Bay is $2.3 \times 10^8 \text{ m}^2$, Wilkinson and Byrne (1977) translated the vertical rate (0.14 cm/yr) into a net rate of sedimentation of $3.2 \times 10^{11} \text{ cm}^3/\text{yr}$ for historic time. Combining rates of sediment yield for the Lavaca and Navidad Rivers and Garcitas Creek, as presented by Greiner, provides a total yield of $1.398 \times 10^{12} \text{ cm}^3/\text{yr}$. Assuming that 30 percent of the fluvial sediment delivered to the bay is trapped in Lavaca Bay (Wilkinson and Byrne), the total fluvial sediment deposited is $4.20 \times 10^{11} \text{ cm}^3/\text{yr}$. This volume is about 1.3-times the net rate of sedimentation noted above. If this volume is spread over the area of the bay,

it yields a vertical sedimentation rate of 0.18 cm/yr, which is slightly higher than Shepard's net rate of 0.14 cm/yr. Assuming that Lake Texana traps about 95 percent of the sediment from the Navidad River Basin, the total fluvial sediment load (including the Lavaca River and Garcitas Creek) delivered to the bay would be reduced to $4.82 \times 10^{11} \text{ cm}^3/\text{yr}$. Again, assuming a trapping rate of 30 percent, this translates into a sedimentation rate from fluvial sources of 0.063 cm/yr, or only about 35 percent of the estimated pre-Lake Texana rate.

Plans to divert the Colorado River into Matagorda Bay (U.S. Army Corps of Engineers, 1981) are expected to increase the sedimentation rate in that bay. Estimated sediment loads that will be contributed to the eastern arm of Matagorda Bay are approximately 1370 acre-ft/yr ($1.69 \times 10^{12} \text{ cm}^3/\text{yr}$) (van Beek and others, 1980), which is about 1.7-times larger than the estimated sediment load (based on Greiner's data noted in the preceding paragraph) previously contributed by the Navidad River to Lavaca Bay. The eventual contribution of sediment from the diverted Colorado River to the central part of Matagorda Bay and the mouth of Lavaca Bay, however, is difficult to assess.

Galveston-Trinity Bay System

Subsidence in the Galveston-Trinity Bay system has had a definite impact on bay bathymetry. Shepard (1953) indicated that overall the Galveston-Trinity Bay system (excluding East and West Bays) had undergone shoaling between 1854 and 1933 at an average rate of 1.44 ft/century (4.3 mm/yr). The rate of sedimentation in Galveston and Trinity Bays calculated by Rehkemper (1969) on the basis of Carbon-14 dates (without considering sediment compaction) was 1.2 ft/century (3.6 mm/yr), which is relatively close to that calculated by Shepard (1953). If the average rate of sea-level rise of about 4.2 mm/yr reported by Marmer (1954) for Galveston (for the period 1909-1937) is assumed to be primarily from subsidence for the bay, then

Shepard's rate of sedimentation becomes about 8.5 mm/yr. The rate of sedimentation near the delta at the mouth of the Trinity River is obviously much higher.

However, more recent rates of subsidence in the Galveston-Houston area are much higher than in the past (fig. 9). As an example, the rate of subsidence near Morgan's Point at the head of Galveston Bay was approximately 47 mm/yr between 1943 and 1978 (Gabrysch, 1984). Of course the rate decreases away from the area of maximum subsidence. Near the center of Trinity Bay the rate is closer to 15 mm/yr (for the period 1973 to 1978; Gabrysch, 1984). Still, this rate is higher than the average sedimentation rate of 8.5 mm/yr based on Shepard's (1953) data.

Morton and McGowen (1980) compared recent sounding data from the Submerged Lands of Texas project (McGowen and Morton, 1980; White and others, 1985) collected during 1977 with 1968 National Ocean Survey bathymetric data (fig. 91). This comparison indicates considerable deepening of the Galveston-Trinity Bay system. The actual magnitudes of the depth increases, which are as high as 5 ft (150 cm) in the upper reaches Galveston Bay, are not reliable because the soundings in 1977 were not adjusted to a mean sea level datum. Nevertheless, a comparison of the changes in depths with subsidence (fig. 9) indicates that the trends or directions of deepening of the bay floor correlates with trends of increasing subsidence. Near the center of the subsidence "bowl", there is no question that the bay is getting deeper; adjacent areas that were previously uplands are now under water. In addition, some spoil deposits previously shown as islands on navigation charts were no longer subaerial features in 1977 when benthic sediment samples were collected for the Submerged Lands of Texas project (White and others, 1985).

Of all the bay systems considered by Shepard (1953), the Galveston-Trinity bay system had one of the largest differences between stream silt load and bay fill (this bay system was surpassed only by East Matagorda Bay, which was the receiving basin for the Colorado River). The combined silt load for the Trinity and San




Figure 91. Bathymetric changes for the Trinity-Galveston bay system for the period 1968-1977. Bathymetry for 1968 is from published navigation charts. Bathymetry for 1977 is based on sounding taken at sampling stations, on 1-mi centers, in the fall and winter of 1977. Because the soundings in 1977 were not adjusted to mean sea level, the depth changes shown in this figure must not be taken as absolute. (From Morton and McGowen, 1980.)

Jacinto Rivers was more than 5 times the calculated bay fill (table 12). Again, factors such as deposition of sediment in stream floodplains, but principally subsidence, probably accounted for the discrepancy according to Shepard (1953). In the upper half of Trinity Bay near the mouth of the Trinity River, deposition rates were high as indicated by changes in depth shown in figure 86. Net sedimentation rates near the margin of the Trinity River delta, in fact, exceeded 20 mm/yr and were more than 8 mm/yr for much of the upper bay. The Trinity River prograded into Trinity Bay about 473 m (0.3 mi) between 1854 and 1933 (Shepard, 1953).

Sedimentation in East Bay was high enough to indicate that sediment from the Trinity River was transported around Smith Point and into the bay (Shepard, 1953). White and others (1985) suggested that trace metals concentrated in East Bay were tied to sediment movement from Trinity Bay (and Trinity River).

In contrast to sedimentation in East Bay, West Bay actually became deeper during the 67 year period represented by Shepard's data. The rate of deepening was about 1.7 mm/yr, the highest rate of all the bays measured. Shepard (1953) concluded that the deepening was the result of slow deposition (far removed from a fluvial sediment source) and subsidence.

Recent measurements (1969-1984) indicate that the silt load of the Trinity River (fig. 21) is only about 14 percent of that reported by Shepard (1953). The U.S. Army Corps of Engineers (USACE, 1942; cited in Rehkemper, 1969) estimated annual average sediment contributions by the Trinity River to be 7,260 acre-ft (which is higher than Shepard's estimate for the combined load of the Trinity and San Jacinto Rivers); the more recent measurements of Trinity River sediment load (528 acre-ft/yr) are only about 7 percent of this value. Contributions of sediment by the Trinity and San Jacinto Rivers (before 1940) were more than 90 percent of the total sediment contributions to the bay system (USACE, 1942; Rehkemper, 1969). Rehkemper (1969) noted that the USACE (1942) estimate for the Trinity River exceeded both the Texas

Board of Water Engineers and USGS (by a factor of approximately 2), but concluded that this higher value could be due to a much shorter sampling period used by the Corps, or to reservoir construction on the Trinity River after 1940, in which case the Corps data would be more representative of unaltered sediment loads.

USDA (1959) and Greiner (1982) estimated (based on suspended load measurements at Romayor) that the annual volume of sediment being deposited in Galveston Bay from the Trinity River was about 4,500 acre-ft, of which about 800 to 900 acre-ft, or approximately 20 percent, was estimated to be bed load (unmeasured). Greiner (1982) acknowledged that more recent measurements indicated this rate of sediment delivery had diminished and that with continued implementation of soil conservation measures and construction of reservoirs up stream, this decline is expected to continue. Recent measurements reflect this decline. Average annual suspended load of the Trinity River at Romayor from 1969 to 1984 was approximately 530 acre-ft, or about 15 percent of the suspended load reported by Greiner (1982). Assuming that about 20 percent of this amount (530 acre-ft) provides an estimate of bed load transported by the Trinity River to the head of the bay will yield a bed-load volume of about 105 acre-ft/yr.

Dredging records (US Army Corps of Engineers unpublished maintenance dredging records) for Anahuac Channel, along which the Trinity River discharges into Trinity Bay, indicate the amount of material dredged from the channel during this period (1969-1984) averaged about 90 acre/ft per year, which is a volume close to the 105 acre/ft/yr approximated above. It is acknowledged that the volume of material deposited in Anahuac Channel is not necessarily a reflection of the total bed load carried by the Trinity River for several reasons. Among them are that some bed load is deposited upstream, some is removed by maintenance dredging along the Channel to Liberty, and some is transported into Trinity Bay at the mouth of the channel. Nevertheless, a closer examination of the dredging records for Anahuac Channel

indicates some trends. For example, dredging records for selected periods show that the annual average volume of sediment removed from the channel during the period 1965 to 1975 exceeds earlier and later periods by a factor of approximately 1.5 (fig. 92). Comparisons of streamflow and river suspended load with volume of sediments dredged on a year by year basis provides some possible reasons for at least some variations in dredged material (fig. 93).

Contributions of sediment to Trinity Bay by shoreline erosion was estimated as less than 5 percent of the total sediment contribution to the bay (USACE, 1942). Although erosion rate of the bay shoreline may have increased in recent years (Paine and Morton, 1986), which could increase sediment supply from this source, it seems likely that the reduction in stream sediment supply plus subsidence will yield increasingly deeper bay waters, particularly in more rapidly subsiding areas, during future years.




Figure 92. Annual-average volume of sediments dredged from Anahuac Channel along the Trinity River. (Based on unpublished dredging records provided by the U.S. Army Corps of Engineers.)




Figure 93. Relationship of sediment volume dredged from Anahuac Channel with annual streamflow along the Trinity River. Streamflow is in acre-feet divided by 120,000, and volume of dredged material is in yds³/length of channel dredged/number of months between dredgings. Two events that possibly affected rates of sediment accumulation in the channel are also shown. (Based on unpublished dredging records provided by the U.S. Army Corps of Engineers, and streamflow data from sources listed in figure 21.)

MACROBENTHOS-SEDIMENT RELATIONSHIPS

Introduction

Bottom sediments contain epifaunal and infaunal invertebrates that live on or in the substrate (benthos) and that obtain their food from the water column or from the sediment. Benthic organisms are important in estuarine ecosystems for several reasons: (1) they provide support for primary producers by recycling nutrients, such as nitrogen, in the sediments (Flint and Kalke, 1985). This source of nutrients might be especially important in stabilizing ecosystems largely dominated by nutrient input from river runoff, which may be subject to climatic variations (Rowe and Smith, 1977); (2) they consume bacteria and meiobenthos and serve as both primary and secondary consumers in the detritus based food chain (Armstrong, 1987); (3) they are food for many bottom-feeding fish; (4) they have limited mobility compared to plankton or fishes, and their abundance and diversity have often been studied in order to demonstrate changes in the health or productivity of an estuary; (5) they accumulate trace metals above concentrations found in surrounding waters or sediment (Scudato and others, 1976); (6) they pellitize fine-grained sediment into agglomerated fecal pellets which have greater settling rates than their composite particles; and (7) they rework the sediment and influence the transport and fate of sediment (Schaffner and others, 1987).

Biologically mediated sedimentation processes may be as important as the mechanical or physical processes which lead to deposition of fine sediments (Biggs and Howell, 1984). The conceptual model (Armstrong, 1987) in figure 94 illustrates the role and importance the benthos plays in cycling of nutrients and in organic sedimentation and organic matter decomposition. The heavier lines show the flow of organic matter to the sediments and the return of nutrients to the water column to be cycled again by phytoplankton. This is the primary flow of organic matter and




Figure 94. Conceptual model illustrating the role of the estuarine benthos in food chain dynamics and nutrient recycling (from Armstrong, 1987).

nutrients in the estuary contrasted to the lesser influence of riverine nutrients and oceanic inwelling (Armstrong, 1987).

Many ecologic factors affect the distribution, diversity, and biomass of benthic invertebrates in estuaries, including substrate, salinity, temperature, organic content, seagrass distribution, interspecific competition, predation, vagility, and others. Early workers in the field of benthic marine ecology, such as Bader (1954), Thorson (1957), and Sanders (1958) and more recent workers, such as Boesch (1971), Johnson (1971), Bacescu (1972), Gray (1974), Rhoads (1974), Holland and others (1975), Holland and others (1977), Loi and Wilson (1979), and Flint and Kalke (1985) concluded that macrobenthic species are greatly influenced by sedimentary parameters and closely associated ecologic variables, such as organic content, depth, turbidity, or sediment stability. However, simultaneous responses to other environmental factors, such as temperature, salinity, currents, light, and many others, make it difficult to determine the ranges of tolerances of organisms to various sediment parameters. Biological interactions between macrofauna (Comitato, 1982) and predation by large, motile predators (Virnstein, 1977) are also important processes controlling community structure. Although this report will concentrate on sediment-faunal relationships and their importance in the estuarine ecosystem, the estuarine environment is complex and no single ecologic factor governs the population dynamics of the benthic community.

The close association between substrate and the benthos begins with their pelagic larvae. Many larval species are restricted to certain types of sediment and settlement is far from random (Thorson, 1957; Gray, 1974). Larvae of many polychaetes and echinoderms are able to test the substrate and those ready to metamorphose are able to delay metamorphosis until they find a suitable substrate (Thorson, 1957). Behavioral responses to light, pressure, gravity, salinity, and water currents play a significant part in restricting the range of substrates available to the larvae (Gray, 1974). Physical properties of the substrate that are of great importance to larval

settlement are structure and contours of the surface, grain size (Gray, 1974), and sediment stability (Orth, 1977). Chemical and biological factors of the substrate that help determine larval settlement are (1) inorganic or organic compounds, (2) the presence of live microorganisms on the surface of the substrate, (3) the presence of populations of the same species, and (4) other factors (Gray, 1974).

Trophic Structure

Another important aspect in the relation between the benthos and sediment is the manner in which an organism feeds. Most benthic species feed in one or more of the following ways: (1) by consuming plant tissue (herbivores); (2) by feeding on living or recently dead animal tissue (carnivores-scavengers); (3) by feeding on the fluids of living tissues (parasites); (4) by feeding on deposited detritus (deposit feeders); or (5) by feeding on organic particles and inorganic detritus in the water column (suspension feeders) (Rhoads, 1974). The latter two methods are the most common; however, the distinction between these feeding types is not always clear because some species of estuarine organisms have shown flexibility in their mode of feeding. For example, McLusky and Elliott (1981) have shown that the bivalve Macoma balthica can spend 10 to 40% of its life suspension feeding and 60 to 90% of its life deposit feeding.

Organisms that feed exclusively on deposited food would be expected to reach maximum diversity on muds containing an abundant food supply (Rhoads and Young, 1970). Weak currents favor the deposition of silt- and clay-sized particles and prevent rapid removal of organic detritus. Consequently the proportion of deposit feeders comprising the fauna will increase as organic content of the sediment increases. High organic carbon in the sediment is related to grain size. Particulate organic carbon may be absorbed to fine-grained sediment, particularly to clay-sized particles (McGowen and others, 1979). It is the organic content of the sediment

and not silt-clay content that is causally related to the proportion of deposit feeders in the sediment (Sanders and others, 1962; Purdy, 1964).

The density (numbers of organisms/m²) of deposit feeders is, in part, controlled by the abundance of microorganisms (Driscoll, 1975; Levinton, 1977). Bioturbation and fecal formation by deposit feeders result in increased surface area for colonization by microorganisms. Driscoll (1975) suggested that a feedback relationship exists, increasing microorganisms resulting in increasing deposit feeder abundance. The latter, through bioturbation and biodeposition, can produce an environment suitable for more microorganisms. The feedback rate is temperature dependent (Driscoll, 1975).

The optimal mean grain size for suspension feeders is in the fine to medium sand range (0.125 to 0.50mm) (Sanders, 1958; Bloom and others, 1972; Whitlatch, 1977), although many suspension feeders also occur on gravel/shell-sized sediments (>2.0mm) (Craig and Jones, 1966). In contrast to deposit feeders, organic content of the sediment is not important to suspension feeders, for by definition suspension feeders feed on organic detritus in the water column, and it is the amount of food in the water that is of primary importance. Other factors being equal, the higher the current velocity, the greater the amount of organic matter brought to the suspension feeders per unit time and consequently the larger the proportion of suspension feeders in the benthos (Purdy, 1964). However, high current velocities can cause considerable substrate mobility or instability and this can reduce the benthic population (Purdy, 1964). Whitlatch (1977) found that coarse sand stations at Barnstable Harbor, Mass. had reduced abundances of both deposit and suspension feeders. These stations were characterized by pronounced surficial ripple marks, evidence of sediment instability.

Sediment instability can also be produced by deposit feeders in fine-grained sediments. Deposit feeders can modify silts and clays by (1) high turnover due to resuspension in tidal currents, (2) high turbidity at the sediment- water interface, and

(3) production of textural and compositional grading (Rhoads and Young, 1970). Such instability inhibits suspension feeders and sessile epifauna by clogging filtering mechanisms, resuspending and burying larvae, and discouraging the settlement of larvae of suspension feeders and adults of sessile epifauna. This process, the exclusion of one trophic group in an area occupied by another, is termed trophic-group amensalism (Rhoads and Young, 1970). Conversely, polychaete and crustacean tubes can help stabilize the sediment and increase species diversity (Orth, 1977). Seagrasses are also known for their ability to stabilize sediments by baffling currents and damping wave action (Ginsburg and Lowenstam, 1958; Orth, 1977).

In Texas bays and estuaries, deposit feeders are abundant in the river-influenced assemblage (table 15). This assemblage generally occurs in the upper bays, near areas of fluvial input and where sediments are dominantly mud or sandy mud. River input brings large quantities of organic laden muds that provide food for deposit feeders and suspension-feeding species that are able to tolerate high turbidities. However, this assemblage is also subjected to greater natural saline fluctuations than are other bay assemblages and the benthic community living in it is probably highly stressed. Although total benthic standing crop or density may be high due to large numbers of estuarine endemics, diversities are low because most benthic species are not able to tolerate the wide range in salinity.

Suspension feeders, especially bivalves, are most abundant on oyster reefs and in marine grassflats (table 15). These habitats are structurally complex and substrates, food, and shelter are available for most feeding types.

Biomass and Density

The most important part of biomass production in all biotopes (niches) of an estuary is that produced by the primary consumers, which include the benthos (Armstrong, 1987). Benthic organisms are in an intermediate position in the estuarine food chain, serving as primary and secondary consumers and transferring energy

Table 15. Feeding type of benthic macroinvertebrate species characteristic of river-influenced, oyster-reef, and grassflat assemblages in the Galveston Bay system.

Location and assemblage	Feeding type
Galveston-Trinity-East Bays	
River-influenced assemblage	
<u>Mulinia lateralis</u>	Suspension feeder
<u>Macoma mitchelli</u>	Deposit feeder
<u>Rangia flexuosa</u>	Suspension feeder
<u>Texadina sphinctostoma</u>	Deposit feeder
<u>Texadina garretti</u>	Deposit feeder
<u>Parandalia fauveli</u>	Carnivore?
<u>Streblospio benedicti</u>	Surface deposit feeder
<u>Capitella capitata</u>	Surface deposit feeder
<u>Mediomastus californiensis</u>	Subsurface deposit feeder
<u>Polydora ligni</u>	Surface deposit feeder
<u>Corophium louisianum</u>	Suspension feeder
Oyster-reef assemblage	
<u>Boonea impressa</u>	Parasitic
<u>Texadina sphinctostoma</u>	Deposit feeder
<u>Crassostrea virginica</u>	Suspension feeder
<u>Ischadium recurvum</u>	Suspension feeder
<u>Brachidontes exustus</u>	Suspension feeder
<u>Mulinia lateralis</u>	Suspension feeder
<u>Nereis succinea</u>	Surface deposit feeder
<u>Polydora ligni</u>	Surface deposit feeder
<u>Mediomastus californiensis</u>	Subsurface deposit feeder
<u>Streblospio benedicti</u>	Surface deposit feeder
<u>Parandalia fauveli</u>	Carnivore or omnivore?
<u>Melita nitida</u>	Unknown
<u>Rithropanopeus harrisii</u>	Scavenger or omnivore?
<u>Cassidinidea lunifrons</u>	Scavenger or omnivore?

Table 15 (cont.)

Location and assemblage

West Bay

Grassflat assemblage

Amygdalum papyrium

Laevicardium mortoni

Chone duneri

Nereis succinea

Streblospio benedicti

Ampelisca abdita

Edotea montosa

Feeding type

Suspension feeder

Suspension feeder

Suspension feeder

Surface deposit feeder

Surface deposit feeder

Deposit feeder

Scavenger or omnivore?

to higher trophic levels such as demersal fish and other predatory organisms.

Biomass is defined as the amount of living substance or living weight of the organisms being studied. Biomass is usually expressed as weight per unit area or grams per square meter. It can also be expressed in dry ash-free weight (Crisp, 1984). Other alternatives for measuring biomass include the chemical analysis of tissues for nitrogen and for caloric content (biomass expressed as energy) (Crisp, 1984). The relative merits of numbers or density (numbers of organisms per square meter), biomass, and energy flow are discussed by Odum (1971). Odum (1971) stresses that numbers overemphasize the importance of small organisms, and biomass overemphasizes the importance of large organisms, but energy flow provides a more suitable index for comparing all populations in an ecosystem.

Infaunal biomass measurements have been used to calculate annual benthic production (Nichols, 1977; Flint and others, 1981). Production can be defined as the total amount of tissue in the population under study during a given time period and can rarely be measured directly (Allen, 1971). Its measurement calls for knowledge of the biomass of the population at the beginning and end of the period and of the mass of living components that have been lost by death or emigration during the period (Allen, 1971). Nichols (1977) discusses the use of the ratio of production to biomass (turnover ratio) for making approximate estimates of annual production from biomass measurements. Production is computed by multiplying the mean of four seasonal estimates of biomass for each species by 4.5, a turnover ratio that Nichols (1977) felt is reliable for a first estimate of the productivity of the common species populations. Total production is obtained by summing the products for each of the common species. Others have questioned the accuracy of the turnover ratio in estimating production. Allen (1951) has stated that the simple multiplication of mean biomass by the number of turnovers in a year would lead to an underestimate of production. Allen (1971) emphasized that it is only when growth and mortality

figures are known that a mathematical relationship between biomass and production can be predicted.

Flint and others (1981) used regression equations to establish a relatively high correlation between infaunal total density and total biomass in the Corpus Christi-Nueces Bay system. After calculating expected biomass from total density measurements, expected and observed biomass measurements were then compared with fishery harvest figures for Corpus Christi Bay from 1973-1982 (Flint and others, 1981; Flint, 1985). Flint (1985) found a strong correlation ($r^2=0.67$) between shrimp fishery yields and mean annual benthic biomass (wet weight). Fisheries production was high following a large benthic bloom in 1980. Flint concluded that either the shrimp responded to the same physical/chemical changes as the benthic infauna or the shrimp derived at least part of their nutrition from the infauna and responded to an increased food source from the large benthic bloom in 1980.

Biomass and density measurements of macrobenthic communities vary considerably (tables 16 through 20), and high values often depend on many ecologic factors that affect either large, heavy organisms (e.g., bivalves and large polychaetes) in the case of biomass values or small, numerous organisms (e.g., small polychaetes) in the case of density values. Sediment type and the presence of seagrasses (table 17) and marsh vegetation (table 20) are just a few of the many important ecologic variables affecting benthic biomass and density measurements.

High biomass and density measurements have been reported for sand, mud, shell/gravel, or mixed (sandy mud or muddy sand) sediments. Studies finding highest biomass and/or density values in sandy sediments include Sanders (1958) in Buzzards Bay, Mass. (table 16), Young and Rhoads (1971) in Cape Cod Bay (table 16), Maurer and others (1978) in Delaware Bay, and Hines and Comtois (1985) in central Chesapeake Bay (table 16). Studies with highest values in muds include Lie and Evans (1973) in Puget Sound (table 16) and Flint and others (1981, 1982) and Flint and Kalke (1985) in Corpus Christi and Nueces Bays. Studies with highest

Table 16. Biomass and density of benthic macroinvertebrates from bays-estuaries-lagoons on the Texas coast and other coastal areas.

<u>Location</u>	<u>Total* or mean biomass (g/m²)</u>	<u>Total** or mean density (organisms/m²)</u>	<u>Average percent sand or textural type</u>	<u>Sieve size (mm)</u>	<u>References</u>
<u>Texas bays-estuaries-lagoons</u>					
Nueces Bay	49		49	0.5	Flint and Kalke, 1985
Corpus Christi Bay	86	2,238	2	0.5	Flint and Kalke, 1985; Armstrong, 1987
Corpus Christi Bay	26	1,925	94		Flint and Kalke, 1985; Armstrong, 1987
Upper San Antonio Bay		2,250	36	0.5	Matthews and others, 1975
Mid-San Antonio Bay		1,890	37		Matthews and others, 1975
Lower San Antonio Bay		590	27		Matthews and others, 1975
<u>Other bays-estuaries-lagoons</u>					
Mouth of Rhode River,					
Maryland (Chesapeake Bay)	119*	34,000**	5	0.5	Hines and Comtois, 1985
Mouth of Rhode River,					
Maryland (Chesapeake Bay)	177*	22,000**	73		Hines and Comtois, 1985
Gulf of Maine	3.18		silt and clayey silt	1.0	Larsen, 1979
Cape Cod Bay, Mass.	1.6-177.9*	11,190-30,150	81-96	1.0	Young and Rhoads, 1971
Cape Cod Bay, Mass.	10.3-16.7*	7,870-14,230	6-33		Young and Rhoads, 1971
Buzzards Bay, Mass.		1,629-12,576**	87-99	0.5	Sanders, 1958
Buzzards Bay, Mass.		1,187-7,982	7-56		Sanders, 1958
Hanno Bay, Baltic	38-90	5,410-6,169	fine sand	1.0	Persson, 1983
Puget Sound, Wash.	16.5-18.8	389-1,122	mud	?	Lie and Evans, 1973
Puget Sound, Wash.	2.9-13.7	544-3,427	silty sand-sand		Lie and Evans, 1973

*Total biomass

**Total density

Table 17. Mean density of benthic macroinvertebrates in marine grasses.

<u>Location</u>	<u>Mean density</u> <u>(organisms/m²)</u>	<u>Sieve size</u> <u>(mm)</u>	<u>References</u>
<u>Texas bays-estuaries-lagoons</u>			
South Bay	532	1.0	White and others, 1986
Lower Laguna Madre	1,911	1.0	White and others, 1986
Baffin Bay	3,354	1.0	White and others, in preparation
Upper Laguna Madre	1,601	1.0	White and others, 1983
Redfish Bay	793	1.0	White and others, 1983
Espiritu Santo Bay	5,726	1.0	White and others, in press
Espiritu Santo Bay (seasonal)	9,153	1.0	
West Bay	4,167	1.0	White and others, 1985
<u>Other bays-estuaries-lagoons</u>			
Indian River, FL	17,479	0.5	Virnstein and others, 1983
Chesapeake Bay, VA	48,900	1.0	Orth, 1977
Apalachicola Bay, FL	38,780	0.5	Sheridan and Livingston, 1983
Pensacola Bay, FL	6,077	0.5	Stoner and others, 1983
Beaufort, NC	672	6.0	Williams and Thomas, 1967

Table 18. Mean density of benthic macroinvertebrates versus mean percent sand in Espiritu Santo, Matagorda, Galveston, and Trinity Bays.

<u>Location</u>	<u>Mean density (organisms/m²)</u>	<u>Mean percent sand per station</u>	<u>Number of stations</u>
Espiritu Santo Bay	86	2	1
	589	35	8
	402	70	15
	532	90	7
Matagorda Bay	744	8	17
	833	33	10
	877	61	3
	1,598	90	15
Galveston Bay	216	9	6
	211	39	3
	333	65	10
	287	91	7
Trinity Bay	111	6	7
	65	34	6
	86	68	1
	53	89	3

Table 19. Mean density versus mean percent sand for benthic macroinvertebrate assemblages in Texas bays.

<u>Assemblage</u>	<u>Mean density (organisms/m²)</u>	<u>Mean percent sand per station</u>
<u>Galveston/Trinity/East Bays</u>		
Oyster reef	604	44.1
River influenced	197	35.7
Open bay center	168	35.1
Bay margin	260	77.6
Inlet influenced	339	59.2
<u>Matagorda Bay</u>		
Open bay center	861	21.7
Inlet influenced	1,969	87.9
Oyster reef	776	60.0
Bay margin	993	92.7
<u>Lavaca Bay</u>		
River influenced	670	54.0
Open bay center	1,538	23.4
Oyster reef	5,417	94.8
<u>Corpus Christi Bay</u>		
Inlet influenced	585	61.1
Bay margin	703	53.6
Open bay center depauperate	111	10.0
Open bay center	285	6.5
Oyster reef	1,298	--
River influenced	388	70.1
<u>San Antonio Bay</u>		
River influenced	2,702	34.3
Oyster reef	2,124	95.1
Enclosed bay center	45	39.2
Bay margin	316	96.8

Table 20. Biomass and density of benthic macroinvertebrates in marshes, mudflats, and sandflats.

<u>Location</u>	<u>Mean biomass (g/m²)</u>	<u>Mean density (numbers/m²)</u>	<u>Sediment type (percent sand)</u>	<u>Marsh vegetation or habitat</u>	<u>Remarks (sieve size in mm)</u>	<u>References</u>
West Bay, Texas		3-4,000	sand and muddy sand	<u>Spartina</u> <u>alterniflora</u>	total fauna (0.4)	Gilmore and Trent, 1974
Newport River, North Carolina	3	7,600	sand	<u>Spartina</u> <u>alterniflora</u>	macroinfauna (0.6)	Cammen, 1979
North Florida	123	475	sand	<u>Juncus</u> <u>roemerianus</u>	total fauna (1.0)	Subrahmanyam and others, 1976
St. Louis Bay, Miss.		316	sandy mud (37)	<u>Juncus</u> <u>roemerianus</u>	mollusks only (1.0)	Bishop, 1981
St. Louis Bay, Miss.		396	sandy mud (44)	<u>Spartina</u> <u>cynosuroides</u>	mollusks only (1.0)	Bishop, 1981
San Francisco Bay, Calif.	13-24	53,000-155,000	mud (14-24)	mudflat	macroinfauna (1.0)	Nichols, 1977
Lynher estuary, England	13.2*		mud (1-3)	mudflat	total fauna (0.5)	Warwick and Price, 1975
Great Bay, New Hampshire	17.8	476	fine sand	sandflat	macroinfauna	Winston and Anderson, 1970
Great Bay, New Hampshire	5.2-27.9	135-987	silt and silty clay	mudflat	macroinfauna	Winston and Anderson, 1970
Grevelingen, Netherlands	0.3-33.9		very fine sand with some mud	sandflat	total fauna (1.0)	Wolff and deWolf, 1977

*Total benthic biomass

values in mixed sediments or shell/gravel include Gillard (1974) in upper Galveston Bay, Holland and others (1975) in Corpus Christi, Nueces, Aransas, and Copano Bays, Matthews and others (1974, 1975) in San Antonio Bay (table 16), Parker (1975) in Cape Cod Bay and Texas bays, and Reinhartz and O'Connell (1983) in upper and central Chesapeake Bay. This is not a complete list of studies that include discussions of biomass/density and sediment relationships, but it illustrates differences between various bay-estuarine systems. The comparison of the results from these diverse areas is confounded not only by the complexity of the environment in the different bay-estuarine systems but also by the use of different sieve mesh sizes (tables 16 through 18) and different sampling periods. Sheridan and Livingston (1983) discuss this problem and note that other studies have found 2 to 6 times the number of individuals in the smaller mesh sieve of 0.5 mm versus the larger 1.0 mm-size. Although both larger and smaller mesh sizes are used, 0.5- and 1.0 mm-sizes are probably the most common sizes used in benthic studies.

The only studies in Texas estuaries correlating benthic infaunal biomass with sediment were those of Flint and others (1981, 1982) and Flint and Kalke (1985). Armstrong (1987) reported on the results of their studies (Flint and others, 1981, 1982; Flint and Kalke, 1985) of benthic infaunal production in Corpus Christi and Nueces Bays and on the inner shelf near Mustang Island from 1981 to 1983. Standing stock biomass was greatest at the muddy, midbay site (station 7) in Corpus Christi Bay and least at the sandy, inner-shelf site (fig. 95 and table 16). Sediment at the midbay site was more than 70% clay. There was a great deal of variation in standing stock biomass at the midbay site, with biomass ranging from less than 4 g/m² after the first year of study to over 100 g/m² after the second year. Biomass at the mixed-sediment site in Nueces Bay (station 2) consistently showed minimum standing stocks in the summer and fall (0.4 - 20.1 g/m²) and maximum biomass in the winter and spring (49.4 - 151.8 g/m²). Biomass at the sandy, inner-




Figure 95. Average measures for sediment characteristics and benthic macroinfaunal biomass at three stations in Corpus Christi Bay from 1981 to 1983. Bars represent 95% confidence intervals around the means (modified from Armstrong, 1987).

shelf site (station 10) was much less variable than biomass at the bay sites, probably because of the stabilizing influence of oceanic waters (Armstrong, 1987).

Densities of organisms from four different bays on the Texas coast illustrate the variation in density with mean percent sand (table 18). In Matagorda Bay, densities were highest in sandy (90 percent sand) sediments, whereas in Trinity Bay, densities were highest in muds (6 percent sand). Densities in Galveston Bay and Espiritu Santo Bay were highest in mixed sediments of sandy mud or muddy sand.

Benthic Assemblages

High biomass and density values are dependent on many variables and not restricted to a particular sediment type; however, samples taken from oyster reefs and seagrasses generally always have high density values (tables 17 and 19). Seagrasses provide food, diverse habitat, and protection from predators, and it is not surprising that many studies (O'Gower and Wacasey, 1967; Santos and Simon, 1974; Orth, 1977; and Virnstein and others, 1983) have reported higher densities of macrobenthos in seagrass than in adjacent bare sand. Mean densities of macrobenthos from seagrass beds along the Texas coast range from 532 organisms/m² in South Bay to 9,153 in Espiritu Santo Bay (table 17). Oyster reefs provide a heterogenous substrate and suitable surfaces for sessile epifauna to attach. Suspension feeders are abundant on oyster reefs or other shell/gravel substrates. Mean densities of organisms on reefs in various Texas bays range from 604 to 5,417/m² (table 19).

Densities are also relatively high in areas with an inlet-influenced assemblage (table 19). (also see the macroinvertebrate assemblage section in White and others, 1983, 1986). Sediments in these areas range from 59 to 88 percent sand (muddy sand to sand), and macroinvertebrate densities range from 339 to 1,969 organisms/m². An inlet-influenced assemblage occurs in bay-estuary-lagoon systems near tidal inlets, and species composition is typical of both bay and nearshore-shelf areas. Densities and diversities in areas with an inlet-influenced assemblage are probably relatively high because of the stability of the environment--salinities, except

in extreme cases of high, localized rainfalls, are probably maintained near oceanic levels.

Densities in areas with a river-influenced assemblage are highly variable, ranging from 197 organisms/m² in the Galveston-Trinity-East Bay area to 2,702 in San Antonio Bay (table 19). River-influenced assemblages occur in the upper bays and tidally-influenced parts of rivers. Organisms in these areas are subjected to natural stress primarily from salinity fluctuations (Bechtal and Copeland, 1970; Holland and others, 1973). Sediment type ranges from 34 to 70 percent sand (sandy mud to muddy sand). Densities in upper San Antonio, Hynes, and Guadalupe Bays are especially high because of large numbers of mollusks, primarily the brackish-water gastropod Texadina sphinctostoma and the bivalves Rangia cuneata, R. flexuosa, and Mulinia lateralis. Matthews and others (1975) and Harper and Hopkins (1976) also found high populations of brackish-water species, including T. sphinctostoma, in upper San Antonio Bay. In their seasonal study, Harper and Hopkins (1976) noted that benthic populations were highly variable and that some species in the upper bay increased dramatically after a spring flood. They concluded that the bloom was probably in response to increased nutrients brought in by the Guadalupe and San Antonio Rivers and only indirectly related to decreased salinities.

Macrobenthic communities in estuaries are dominated by populations that show large seasonal and year-to-year fluctuations in abundance (Holland, 1985). Many species respond to salinity changes, especially those of an extreme nature, with large population increases (Harper and Hopkins, 1976; Flint and others, 1981) or decreases (Stone and Reish, 1965; Johnson, 1980). Another source of variation for most species is that associated with seasonal recruitment cycles as illustrated by a seasonal study of macrobenthic populations in Espiritu Santo and Lavaca Bays (fig. 96). The large increase in density during March at the sandy (89 percent sand) station in Espiritu Santo Bay was primarily due to an increase in numbers of the suspension-feeding bivalve, Lyonsia hyalina floridana. This may be the time of year for the





Figure 96. Density (numbers of individuals/m²) by month of benthic macroinvertebrates at sandy and sandy mud stations in Espiritu Santo and Lavaca Bays.

annual recruitment pulse for Lyonsia, as Flint and others (1981) also reported large numbers of Lyonsia in Corpus Christi Bay during February and March. Benthic populations increased in the winter and early spring at both stations in Lavaca Bay (fig. 96), although the fluctuation was not as large as at the sandy station in Espiritu Santo Bay. Peaks in benthic populations in the estuaries on the Texas coast generally occur in the winter and early spring (Harper and Hopkins, 1976). Benthic populations at the sandy-mud (29 percent sand) station in Espiritu Santo Bay were small and fluctuated very little during the eight months of study (fig. 96).

Changes in sediment parameters, even on a small scale, can result in population fluctuations on a seasonal basis (Holland, 1985). Holland (1985) found that consistent but small-scale changes in silt-clay content of stations in a muddy-sand habitat accounted for seasonal variations in populations within that habitat. Other environmental factors, such as salinity, dissolved oxygen, and temperature, did not vary within the muddy-sand habitat over ranges sufficient to influence macrobenthic abundance.

Marshes

There is an extensive literature on salt-marsh animal distributions, life histories, and ecology (Daiber, 1982), and even many studies of faunal groups and their relationship to sediment in the marsh. For example, Whiting and Mashiri (1974), Daiber (1982), Barnwell and Thurman (1984), and Thurman (1984) have studied the close correlation between the distribution of the fiddler crabs, Uca and Sesarma, and sediment type and organic content of the sediment. However, relatively few studies have examined the relationship between sediment and biomass or density of the total benthic community in the marsh.

Gilmore and Trent (1974) compared the abundance of benthic macroinvertebrates in West Bay, Texas between a natural marsh, an adjacent marsh altered by channelization, bulkheading and filling, and an open-bay area, and they related abundance to substrate in the three areas. Gilmore and Trent (1974) found that

densities for all organisms combined were highest in the marsh (table 20) and least abundant in the open bay. Crustaceans showed a preference for sandy substrates in the marsh, whereas polychaetes were most abundant in canals and at stations with low to intermediate amounts of silt and clay.

Cammen (1979) took monthly benthic samples from a Spartina marsh in North Carolina. Infaunal abundance was greatest in late winter and early spring and least in summer and early autumn. Numbers of individuals ranged from 2,200 to 15,500/m² (mean density of 7,600) and biomass ranged from 1.3 to 6.1 grams ash-free dry weight (AFDW)/m² (mean biomass of 3 grams AFDW). Cammen (1979) estimated that annual production of the infauna was 5.9 grams AFDW/m². Production for the polychaete Nereis succinea, which accounted for most of the biomass, was estimated to be 4.1 grams AFDW/m². Sediment at the marsh sites was mainly medium to fine sand.

Subrahmanyam and others (1976) studied the infauna and epifauna in a Juncus marsh in north Florida. The gastropod Littorina irrorata accounted for 81% of a total biomass of 123 grams/m² (table 20). Peaks of invertebrate abundance occurred in the winter and fall. Subrahmanyam and others (1976) reported that organism densities and diversities are generally higher in the more flooded or low marsh zones than in the mid or high marsh zones, because estuarine species tend to invade lower marshes more easily and there is a greater availability of organic detritus in the low marsh. Subrahmanyam and others (1976) compared marsh densities of their study and other studies with densities of macroinvertebrates from estuaries and, contrary to the results of Gilmore and Trent, they found that marsh densities appeared to be lower. They speculated that this was because marsh sediments are covered with plant rhizomes and living space is reduced (Subrahmanyam and others, 1976).

Mudflats and Sandflats

The mudflat invertebrate community is an important link in the cycling of organic matter to the estuary. Several investigators working in mudflat environments

(e.g., Boyden and Little, 1973; Nichols, 1977) have proposed high productivity for this habitat despite small numbers of species.

Nichols (1977) found that infaunal densities and biomass were very high in the mudflats of San Francisco Bay (table 20). Densities ranged from 53,000 to 155,000 organisms/m², depending on the station; however, actual densities should be much higher as Nichols (1977) speculated he was losing more than one-half of all the individuals of the bivalve Gemma gemma through the 0.5 mm mesh sieve. Total biomass varied from 13 to 24 g AFDW/m² without large seasonal variations. Three species of bivalves made up most of the biomass. Nichols (1977) observed that the large densities of infaunal species indicate the mudflats act as sinks for organic matter produced on the marsh, in the water column, and on the mud itself. The high secondary productivity supported a large shorebird community.

Winston and Anderson (1970) studied the amount of bioturbation in sediments at a sandflat station and at five mudflat stations in Great Bay, New Hampshire. Densities and biomass measurements at the sandflat station were about half-way between the high and low measurements for the mudflat stations (table 20).

Wolff and de Wolf (1977) took monthly biomass measurements at five stations in the sandflats of the Grevelingen estuary, the Netherlands. The highest biomass values were found low in the intertidal zone, with a gradual decrease up to the high water mark. Below the low water mark, biomass was generally low. Mollusks and polychaetes, particularly Arenicola marina, comprised most of the biomass at each station.

Species Diversity and Species Richness

The oldest and most fundamental concept of diversity is species richness or the number of species in a community. Direct species counts provide one of the simplest, most practical, and most objective measures of species richness (Peet,

1974). However, direct counts do not provide enough information about the underlying community, as they don't show how individuals are distributed among the constituent species. As a result, various diversity indices have been devised that are influenced by both species richness and species dominance or how evenly the species are distributed. Probably the most widely used index is the Shannon index or Shannon-Weaver diversity index (H') (see White and others, 1983, for formula). The Shannon index will increase with the number of species and as the proportions of individuals among the species become more equal. Diversity index interpretations should be made cautiously (McIntosh, 1967), because it is very easy to read into the numbers meanings that are not there. This problem is inherent in the Shannon-Weaver formula, because it is affected by both species number and species dominance. Therefore, a single diversity number may be misleading.

Factors potentially increasing species diversity may be summarized as follows: (1) With time, all communities tend to increase in diversity. (2) With increased structural complexity, diversity may be expected to increase. (3) With a predictable environment having a constancy of climate, diversity will increase. (4) With increased competition, diversity will increase. (5) With increased number of predators, competition among prey species will be reduced and diversity will be increased (Gray, 1974). Of these factors, spatial heterogeneity or the structural complexity of the environment is the only one which relates directly to the influence of sediment on community structure (Gray, 1974). Coarse and heterogeneous sediments are more structurally complex and have higher diversities than fine and homogeneous sediments (Gray, 1974); therefore, gravel/shell and muddy sands or sandy muds are generally more diverse than muds or sands.

Sanders (1968) combined the predictable environment, time, and competition hypotheses to form the stability-time hypothesis. This hypothesis states that physical instability in an environment prevents the establishment of diverse communities, and conversely, in a stable environment, with time, speciation and immigration will lead

to high diversity. Physical changes such as rapid and severe changes in salinity and temperature, erosion, and rapid deposition may produce a physically controlled community which is characterized by low diversity (Johnson, 1970). Ecological systems in estuaries that are subject to high levels of natural stress include high energy beaches that are stressed from breaking waves and deltas that receive high rates of sedimentation (Copeland, 1970). Wohlschlag and Copeland (1970) emphasize that estuaries are fragile and that even slight stresses on estuarine systems over long periods of time tend gradually to reduce species diversity, even though yields of common species to exploitation tend to be maintained at reasonable levels while the stresses themselves may tend to remain unrecognized. Shallow bays and estuaries on the Texas coast experience large and unpredictable changes in the environment, and environmental instability may override structural complexity as the major factor controlling diversity.

River-influenced areas are especially susceptible to drastic and sometimes long-term changes in salinity, and benthic assemblages of these areas are characterized by having low diversities and numbers of species (table 21). Sediments are fairly homogeneous and mostly muds except for sands or muddy sands near bay margins. River-influenced assemblages may be found in all or most parts of a bay, such as upper Galveston and Trinity Bays (White and others, 1985), or they may be restricted to creeks or rivers, such as the Colorado River near Matagorda (White and others, in press a). Benthic species in these oligohaline to mesohaline zones (salinities of 0 to 15 parts per thousand) are generally estuarine endemics, such as Rangia cuneata, or euryhaline opportunists, such as Streblospio benedicti (Boesch, 1977; Schaffner and others, 1987).

Bay habitats where structural complexity is important and diversities and species richness are generally very high are oyster reefs and other shelly sediments and marine grassflats (tables 21 and 22). These environments provide surfaces for the attachment of epifauna and many potential niches for infauna. Oyster reefs and

Table 21. Mean percent sand, species diversity, and mean number of species of benthic macroinvertebrate assemblages.

<u>Assemblage</u>	Mean percent sand <u>per station</u>	Range in diversity (H')	Mean number of species
<u>Galveston/Trinity/East Bays</u>			
Oyster reef	44.1	1.15-2.23	10.3
River influenced	35.7	0.00-2.15	3.5
Open bay center	35.1	0.00-2.00	5.4
Bay margin	77.6	0.56-2.49	6.6
Inlet influenced	59.2	0.58-2.56	9.9
<u>Matagorda Bay</u>			
Open bay center	21.7	0.00-2.88	14.4
Inlet influenced	87.9	1.09-2.85	18.9
Oyster reef	60.0	2.87	34.0
Bay margin	92.7	0.00-2.67	13.6
<u>Lavaca Bay</u>			
River influenced	54.0	0.00-2.37	7.8
Open bay center	23.4	0.56-2.67	14.0
Oyster reef	94.8	2.79	42.0
<u>Corpus Christi Bay</u>			
Inlet influenced	61.1	0.64-2.87	15.3
Bay margin	53.6	0.00-2.17	7.4
Open bay center depauperate	10.0	0.00-1.89	3.8
Open bay center	6.5	0.89-2.46	8.0
Oyster reef	--	0.98-2.33	10.0
River influenced	70.1	1.08	4.0
<u>San Antonio Bay</u>			
River influenced	34.3	0.00-1.72	5.1
Oyster reef	95.1	1.05-1.86	8.4
Enclosed bay center	39.2	0.00-1.55	1.8
Bay margin	96.8	0.00-1.77	4.6

Table 22. Mean percent sand, species diversity, and mean number of benthic macroinvertebrates in marine grassflat assemblages.

<u>Location</u>	Mean percent <u>sand</u>	Range in <u>diversity (H')</u>	Mean number <u>of species</u>
Lower Laguna Madre	69	0.56-3.06	22.0
Upper Laguna Madre	79	0.00-2.70	14.8
Redfish Bay	50	0.00-2.38	12.2
Corpus Christi Bay	95	1.97-2.65	13.5
Espiritu Santo Bay	70	1.07-2.44	28.0
East Matagorda Bay	89	2.03-2.31	13.5
Christmas Bay	30	2.39	24.0

marine grassflats may provide many more refuges for prey species than might structurally simple localities, and more refuges should result in higher diversity of prey (Menge and Sutherland, 1976). Also, these habitats might decrease the foraging efficiency of the predator (Menge and Sutherland, 1976). Craig and Bright (1986) found high populations of the bivalve Mercenaria mercenaria texana in shelly sediments in Christmas Bay, and they speculated that shell fragments may provide young bivalves protection from predators, especially blue crabs, and a favorable substrate to which the bivalve larvae can attach in the byssal stage.

White and others (1985; in press b) found that stations in the Galveston and Matagorda Bay systems containing more than 10 percent shell (gravel) generally had more species (figs. 97 and 98). Of the eight stations in the Galveston Bay system having more than 10 percent shell, six occur in lower Galveston Bay, one in West Bay, and one in East Bay. Of the five stations having more than 10 percent shell in the Matagorda Bay system, three occur in Matagorda Bay and two in Carancahua Bay. Total numbers of species in Carancahua Bay are highest at the shelly stations. Of 10 stations sampled monthly for eight months in Espiritu Santo Bay, species diversity (H') was highest at the only station with sediment containing a high percentage of shell (38 percent shell). Holland and others (1975) also reported high numbers of species at stations with shelly sediments in Corpus Christi Bay.

Diversities (H') of benthic macroinvertebrates in marine grassflat assemblages range from 0.00 in upper Laguna Madre and Redfish Bay to 3.06 in lower Laguna Madre (table 22). Most diversities (H') at grassflat stations, especially in lower Laguna Madre, are high (H' above 2.0). Species richness (mean number of species per station) is also high, ranging from 12.2 in Redfish Bay to 28.0 in Espiritu Santo Bay (table 22). Except for the oyster-reef and grassflat assemblages, the mean number of species per station for most other bay assemblages is less than 10.0 (tables 21 and 22).




Figure 97. Scattergram of total species and percent sand in the bays of the Galveston-Houston area (from White and others, 1985).




Figure 98. Scattergram of total species and percent sand in the bays of the Port Lavaca area (from White and others, in press b).

Other investigators (O'Gower and Wacasey, 1967; Heck and Wetstone, 1977; Virnstein and others, 1983; Lewis and Stoner, 1983; Orth and others, 1984) have also found highly diverse faunal communities in seagrass beds, particularly in comparison with nonvegetated areas. Orth and others (1984) suggested that the abundance of many species, both epifauna and infauna, was positively correlated with two distinct aspects of plant morphology: (1) the root-rhizome mat, and (2) the plant canopy. Heck and Wetstone (1977) found that invertebrate species number was not significantly correlated with plant species number, but it was significantly correlated with plant biomass.

Other studies that show physical instability affecting species diversity in the bays include Woodward-Clyde Consultants (1977) and Holland and others (1973). Woodward-Clyde Consultants (1977) found that the benthic community was depauperate in species where the physical instability of the substrate was the greatest. For example, there were relatively few species on the shallow, western shoreline of Matagorda Bay where wave action disturbs the sandy sediment. Also, intensive shrimp trawling activity in the Matagorda Ship Channel created an unstable substrate and was responsible for low numbers of species and individuals at the channel station. On the other hand, physical disturbance was relatively low in the deep, level bottom habitat in Matagorda Bay. Substrates were predominantly silt and shell, and numbers of species were relatively high at all stations.

Holland and others (1973) studied the structure of the benthic community in the Galveston Bay system to ascertain water quality. They applied various diversity indices, including the Shannon index (H'), to data collected during four sampling periods in 1971 to 1972 at five stations in the Galveston Bay system. Two stations are located in the upper bay, one near the Houston Ship Channel - Clear Lake region and one in Trinity Bay. The three lower bay stations are in the middle of West Bay, near the Texas City Ship Channel, and in East Bay. Holland and others (1973) found that three of the stations were areas of "normal estuarine stress," or

stations having macrobenthic H' values above 2.0. The stations with normal stress were in upper Galveston Bay near Clear Lake, in East Bay, and in central West Bay. The Trinity Bay and Texas City Ship Channel stations showed evidence of great stress. Holland and others (1973) concluded that the Trinity Bay station was probably stressed naturally, primarily by salinity fluctuations. The Texas City Ship Channel site showed intermittent stress, possibly owing to manmade pollution.

Gilmore and others (1976) found that taxa diversity (numbers of taxa/m²) declined from the high salinity lower Lavaca Bay to the low salinity upper bay and river area. Taxa diversity was higher during late winter and early spring when sustained fresh-water inflow was generally low. Gilmore and others (1976) associated high diversity with high salinity, low turbidity, and low nutrient concentrations.

Summary

Benthic organisms are important in estuarine ecosystems because they provide an essential link in the estuarine food web, and they influence the fate of sediment and contaminants in the sediment or water column. A knowledge of animal-sediment relations is important because the benthos are closely associated with sediments, and any change in quantity, suspension, distribution, or deposition of sediments may affect the trophic structure, density, diversity, biomass, and ultimately the productivity of the benthos. Also, the loss of marsh habitat from the combination of a loss in fluvial sediment, subsidence, and sea level rise would affect the total productivity of the benthos. Any natural or man-induced changes that affect the benthos may upset the fragile ecological balance that is present and ultimately affect fisheries production.

The following findings resulted from this literature synthesis on macrobenthos-sediment relationships:

(1) Trophic structure

(a) Deposit feeders are most abundant in fine-grained sediment with high organic content. In Texas estuaries, deposit feeders are most abundant in the river-influenced assemblage of upper bays.

(b) Suspension feeders are most abundant on sediments of fine to medium sand and on shelly substrates. In Texas estuaries, suspension feeders and other trophic groups are most abundant on oyster reefs and other shelly substrates and in marine grassflats.

(2) Biomass and density

(a) High biomass and density measurements have been reported for sand, mud, shell/gravel, or mixed sediments.

(b) The only studies of Texas estuaries correlating benthic infaunal biomass with sediment were for the Corpus Christi-Nueces Bay system. In those studies, benthic biomass was highest at a muddy, midbay station.

(c) Benthic densities on oyster reefs and other shelly substrates, in marine grassflats, and near tidal inlets are generally high.

(d) Densities in areas with a river-influenced assemblage are highly variable because these areas are stressed from natural disturbance, primarily salinity fluctuations.

(e) Benthic densities are variable seasonally and are probably most dependent on seasonal recruitment cycles, although changes in sediment parameters, even on a small scale, can result in density fluctuations.

(f) There have been no studies of benthic biomass in Texas marshes, mudflats, or sandflats.

(g) Benthic biomass and densities are generally high in marshes, mudflats, and sandflats.

(h) The only study of Texas marshes that correlated benthic densities and sediment was in West Galveston Bay.

(3) Species diversity and species richness

(a) Coarse and heterogenous sediments, such as gravel/shell and sandy muds or muddy sands, are more structurally complex and thus more diverse than sands or muds.

(b) In Texas estuaries, environmental instability may override structural complexity as the major factor controlling species diversity, especially in areas with a river-influenced assemblage.

(c) Estuarine habitats where structural complexity is important and diversities and species richness are high are oyster reefs and other shelly sediments and marine grassflats.

(d) In areas where physical instability of the substrate is high, such as areas with intensive shrimp trawling activity, benthic diversities and species richness are low.

SUMMARY AND PRELIMINARY CONCLUSIONS

Sedimentation along the Texas Gulf Coast is affected by various interactive processes including riverine discharge, astronomical and wind-generated tides, waves and currents, episodic events such as fresh-water flooding and tropical cyclones, biodeposition, subsidence and sea-level rise, and human activities. The primary sources of sediments delivered to the estuaries are the rivers that cross the coastal plain. Deposition of the fluvial sediments along the alluvial river valleys and at the river mouths has produced extensive fluvial-deltaic deposits on which marshes and other wetlands -- essential components of a healthy estuarine ecosystem -- have developed. Sediments delivered to the marshlands not only provide a source of nutrients for sustained plant growth, but they also provide an inorganic foundation necessary to maintain the substrate above a rising sea level. The submergence of more than 4,000 hectares (10,000 acres) of fluvial-deltaic wetlands between the mid-1950's and the late-1970's in two areas along the Texas coast signifies that sediments in these areas are not accumulating at rates sufficient to counter the effects of relative sea-level rise.

Wetlands are being lost at a dramatic rate on the Mississippi River delta. In fact, land-loss rates have accelerated geometrically during the 20th century, largely as a result of natural processes, of harnessing the Mississippi River deltaic-sedimentation processes, and of accelerated subsidence (natural and possibly human induced) (Gagliano and others, 1981; Boesch and others, 1983; Wells and Coleman, 1987). Results of several investigations on Louisiana marsh sedimentation indicate that marsh aggradation (vertical accretion) rates are not keeping pace with relative sea-level rise.

There have been few studies of sedimentation in Texas marshes. Investigations have principally focused on shoreline changes to document retreat (erosion) and advancement (accretion) of the shoreline. The loss of interior marshes in fluvial-deltaic

areas has not been systematically investigated to determine the historical rates of marsh transformation to open water, a process that has been documented only in selective areas (White and others, 1985; 1987).

The delivery of fluvial sediments to the bay-estuary-lagoon systems has been a process operating through much of the Holocene Epoch, and of course continues today. The most extensive look at the accumulation of sediments in the bays and estuaries of Texas was done by Shepard (1953); the most thorough investigation of a single bay system (Lavaca Bay) was accomplished by Wilkinson and Byrne (1977). In general, Shepard (1953) concluded, on the basis of bathymetric surveys made in the latter half of 1800's and mid-1930's, that Texas bays and estuaries were shoaling (becoming shallower) at an average rate of 3.8 mm/yr. The highest rates of shoaling occurred at the heads of bays where deposition of fluvial sediments was at a maximum. In bays located away from fluvial input shoaling rates were much lower and in some areas deepening of the bay floors had occurred. Wilkinson and Byrne (1977) concluded that historical rates of sedimentation in Lavaca Bay are higher than rates over a geologic time frame (past 8 to 10 thousand years) and suggested the higher historic rate may be related to land-use practices (cropland) in the drainage basin.

Only two rivers discharging into Texas estuaries have significantly extended their deltas since the mid-1800's: the Colorado and Trinity rivers. The Colorado River delta has a unique history of very rapid progradation across the eastern arm of Matagorda Bay following the removal of a log raft upstream that had blocked sediment along the lower reaches of the river channel. Shepard (1953) estimated that the Trinity River had extended its delta about 0.5 km since the mid-1800's. On the Guadalupe River delta a small subdelta has prograded into Mission Lake as a result of the artificial diversion of river discharge and sediment load into the shallow lake.

A major process countering the trend toward net sedimentation or shoaling of bay floors and net aggradation of marsh areas is relative sea-level rise. Generally

composed of two components, a lesser component of eustatic (global) sea-level rise and a more significant component of land-surface subsidence, relative sea-level rise along the Texas coast ranges up to more than 12 mm/yr along portions of the Gulf coast (Swanson and Thurlow, 1973). Rates are considerably higher than this in areas undergoing human-induced subsidence due to underground fluid withdrawal, such as in the Houston area. More recent comparisons of bathymetric data in the Galveston-Trinity bay system indicate that water depths have increased as a result of subsidence (Morton and McGowen, 1979).

Over the past 3 to 4 decades, there has been a marked decline in fluvial sediments delivered by many coastal rivers. Among the rivers are the Trinity, San Jacinto, Brazos, Colorado, and Nueces. The sediment loads of other rivers have also possibly diminished, but in many instances sediment-load measuring stations are not located close enough to the coast to adequately reflect the decline. Sediment load in several rivers, for which there is data, is less than half the previous load measured before the 1950's, and in some cases the load is less than 15 percent of previous amounts. Decreases in stream sediment load are related to different factors including implementation of soil conservation measures. But comparisons of reservoir development in the drainage basins with reductions in stream sediment load indicate reservoirs are probably the major factor. Large reservoirs can trap from 95 to 100 percent of the sediment delivered to them, and reduction in peak flows below the dams decreases the ability of the stream to transport sediments accumulating downstream at the mouths of tributaries. The largest quantities of sediment are delivered to the estuaries during major flood events, which are controlled along streams with large reservoirs.

Channel degradation downstream from reservoirs can contribute sediment to estuarine areas, but the amounts are hard to quantify because of numerous variables involved. Isphording (1986) reported an increase in sand and clay deposition in

Apalachicola Bay after reservoir development, but he also reported a striking reduction in silt. Silt, which had been previously supplied under natural conditions, was trapped along with sand by the reservoir. He suggested that clay was washed over the spillways and continued downstream to the bay, and he attributed increases in sand to channel erosion downstream from the reservoirs. Studies by Williams and Wolman (1983) of effects downstream from reservoirs generally indicate substantial decreases in average annual peak discharges with marked reductions in suspended sediment load for hundreds of kilometers downstream. In some major rivers, annual sediment loads did not equal pre-dam values for hundreds or thousands of kilometers. Degradation of channel beds generally occurred during the first decade or two after dam completion.

The environments at the mouths of many rivers along the Texas coast have been significantly modified through canal dredging and sediment disposal activities. These modifications have altered the hydrologic regime and sediment dispersal pathways in some areas, which can hinder natural sedimentary processes and promote erosion.

With relative sea-level rise along the Texas coast matching rates reported along much of the Gulf coast of Louisiana, and with marked declines in fluvial sediments delivered to many Texas coastal fluvial-deltaic and estuarine systems, it is probable that marsh aggradation rates and bay floor sedimentation rates are no longer keeping pace with rates of relative sea-level rise in many areas. This conclusion is supported by observations in selected areas, but a systematic investigation of historical changes in interior marshes in fluvial-deltaic areas, and measurements of marsh aggradation rates have not been conducted along the Texas coast (except in two areas as part of this study, White and Calnan, in preparation). More recent comparisons of bathymetric data in Texas bay-estuary-lagoon systems would provide up-to-date information on shoaling or submergence/erosion rates to compare with Shepard's earlier work. Also, establishment and operation of stream-discharge and sediment-load (including

bed load) measuring stations on many streams at locations closer to the coast would allow a better estimation of the quantities of fluvial sediments delivered to the bay-estuary-lagoon systems of Texas.

REFERENCES

(Includes those cited in text and others)

- Adams, D. B., and Tingley, C. A., 1977, A study of the hydrological and nutrient exchange processes in the Trinity River delta, Texas, part II, marsh habitats and primary production in the Trinity River delta: report prepared for Texas Department of Water Resources by Espey, Huston and Associates, Inc., Document no. 77-103, Job no. 0216, 20 p.
- Adey, E. A., and Cook, H. M., 1964, Suspended-sediment load of Texas streams: compilation report, October 1959 - September 1961: Texas Water Commission, Bulletin 6410, 49 p.
- Aksoy, S., 1967, Riverbed degradation downstream of large dams: Sedimentation, annotated bibliography of foreign literature, 1967-1968, Survey No: 5, U. S. Department of Agriculture and National Science Foundation, Washington D.C., # 3787, p. 5.
- Alaniz, R. T., and Goodwin, R. H., 1974, Recent sediments of a hypersaline estuary bay: Gulf Coast Association Geological Societies Transactions, v. 24, p. 308-313.
- Allen, K. R., 1971, Relation between production and biomass: Journal of the Fisheries Research Board of Canada, vol. 28, p.1573-1581.
- Allen, K. R., 1951, The Horokiwi Stream, a study of a trout population: New Zealand Marine Department, Fisheries Bulletin No. 10, 238 p.
- Aller, R. C., and Dodge, R. E., 1974, Animal-sediment relations in a tropical lagoon, Discovery Bay, Jamaica: Journal of Marine Research, v. 32, no. 2, p. 209-232.
- Almasi, M. N., Hoskin, C. M., Reed, J. K., and Milo, J., 1987, Effects of natural and artificial *Thalassia* on rates of sedimentation: Journal of Sedimentary Petrology, v. 57, no. 5, p. 901-906.
- Anderson, R. Y., 1977, Short term sedimentation response in lakes in western United States as measured by automated sampling: Limnology and Oceanography, v. 22, p. 423-433.
- Armentano, T. V., and Woodwell, G. M., 1975, Sedimentation rates in a Long Island marsh determined by 210-Pb dating: Limnology and Oceanography, v. 20, no. 3, p. 452-456.
- Armstrong, N. E., 1982, Responses of Texas estuaries to freshwater inflows, in Kennedy, V. S., ed., Estuarine Comparisons: New York, Academic Press, p. 103-120.
- Armstrong, N. E., 1987, The ecology of open-bay bottoms of Texas: a community profile: Washington, D.C., U.S. Department of the Interior, Fish and Wildlife Service, 104 p.

- Armstrong, N. E., Harris, S. E., and Gordon, V. N., 1977, Exchange rates for carbon, nitrogen, and phosphorus in the Trinity River delta marshes: The University of Texas at Austin, Center for Research in Water Resources, CRWR-154, EHE 77-07, 63 p.
- Armstrong, N. E., and Brown, B. A., 1976, Exchange of carbon, nitrogen, and phosphorus in Lavaca Bay, Texas marshes. Volume 1, The role of sediments in nutrient exchange in the Lavaca Bay brackish marsh system: The University of Texas, Center for Research in Water Resources, final report prepared for Texas Water Development Board under IAC (74-75)-1217, CRWR-147.
- Armstrong, N. E., and Gordon, V. N., 1979, Nutrient exchange studies on the seagrasses of Texas: final report: Texas Department of Water Resources, IAC(78-79)-0914, 85 p.
- Armstrong, N. E., and Ward, G. H., 1981, Effects of alterations of freshwater inflows into Matagorda Bay, Texas, in Cross, R. D., and Williams, D. L., Proceedings of the National Symposium on Freshwater Inflow to Estuaries, vol. 2, p. 179-196.
- Aronow, Saul, 1971, Nueces River delta plain of Pleistocene Beaumont Formation, Corpus Christi region, Texas: American Association of Petroleum Geologists Bulletin, v. 55, p. 1231-1248.
- Aten, L. E., 1983, Indians of the upper Texas Coast: Academic Press, New York, 370 p.
- Bader, R. G., 1954, The role of organic matter in determining the distribution of pelecypods in marine sediments: Journal of Marine Research, v.13, p. 32-47.
- Barnwell, F. H., and Thurman, C. L., II, 1984, Taxonomy and biogeography of the fiddler crabs (Ocypodidae: Genus Uca) of the Atlantic and Gulf coasts of eastern North America: Zoological Journal of the Linnean Society, v. 81, p. 23-87.
- Bartberger, C. E., 1976, Sediment sources and sedimentation rates, Chincoteague Bay, Maryland and Virginia: Journal of Sedimentary Petrology, v. 46, no. 2, p. 326-336.
- Barth, M. C., and Titus, J. G., 1984, Greenhouse effect and sea level rise - a challenge for this generation: New York, Van Nostrand Reinhold Company, 325 p.
- Bascescu, M. C., 1972, Substratum - Animals, in Kinne, O., ed., Marine ecology, v. 1, p. 1291-1322.
- Baumann, R. H., Day, J. W., Jr., and Miller, C. A., 1984, Mississippi deltaic wetland survival: sedimentation versus coastal submergence: Science, v. 224, no. 4653, p. 1093-1095.
- Beard, J. H., Sangree, J. B., and Smith, L. A., 1982, Quaternary chronology, paleoclimate, depositional sequences and eustatic cycles: American Association of Petroleum Geologists Bulletin, v. 66, no. 2, p. 158-169.

- Bechtal, T. J., and Copeland, B. J., 1970, Fish species diversity indices as indicators of pollution in Galveston Bay, Texas: *Contributions in Marine Science*, v. 15, p. 103-132.
- Behrens, E. W., 1963, Buried Pleistocene river valleys in Aransas and Baffin Bays, Texas: *The University of Texas Marine Science Institute Publication*, v. 9, p. 7-18.
- Behrens, E. W., 1966, Surface salinities for Baffin Bay and Laguna Madre, Texas, April 1964-March, 1966: *University of Texas, Marine Science Institute Publication*, v. 11, p. 168-173.
- Behrens, E. W., and Watson, R. L., 1974, Corpus Christi Water Exchange Pass: a case history of sedimentation and hydraulics during its first year: unpublished report, U. S. Army Corps of Engineers, Coastal Engineering Research Center, Contract DACW 72-72-C-0027, 150 p.
- Bell, S. B., 1980, Meiofauna-macrofauna interactions in a high salt marsh habitat: *Ecological Monographs*, v. 50, no. 4 p. 487-505.
- Bennett, H. H., 1951, Relation of soil erosion to coastal erosion: *The Texas Journal of Science*, v. 3, no. 2, p. 147-161.
- Benton, A. R., Jr., Hatch, S. L., Kirk, W. L., Newnam, R. M., Snell, W. W., and Williams, J. G., 1977, Monitoring of Texas coastal wetlands: College Station, Texas A&M University, Remote Sensing Center, Technical Report RSC-88, 124 p.
- Benton, A. R., Jr., Snell, W. W., and Clark, C. A., 1979, Monitoring and mapping of Texas coastal wetlands, Galveston Bay and Sabine Lake areas, 1978 growing season: Texas A&M University, Remote Sensing Center, Technical Report RSC-102, 123p.
- Bernard, H. A., Major, C. F., Jr., Parrott, B. S., and LeBlanc, R. J., 1970, Recent sediments of southeast Texas--A field guide to the Brazos alluvial and deltaic plains and the Galveston barrier island complex: *The University of Texas at Austin, Bureau of Economic Geology, Guidebook 11*, 132 p.
- Bernard, H. A., Major, C. F., Jr., and Parrott, B. S., 1959, The Galveston barrier island and environs: a model for predicting reservoir occurrence and trend: *Gulf Coast Association of Geological Societies Transactions*, v. 9, p. 221-224.
- Bernard, H. A., and LeBlanc, R. J., 1965, Resume of Quaternary geology of the northwestern Gulf of Mexico province, in Wright, H. E., Jr., and Frey, D. G., eds., *The Quaternary of the United States*: Princeton, N. J., Princeton University Press, p. 137-185.
- Beschta, R. L., 1987, Conceptual models of sediment transport in streams, in Thorne, C. R., Bathurst, J. C., and Hey, R. D., *Sediment transport in gravel-bed rivers*: John Wiley & Sons Ltd, New York, p. 387-408.
- Biggs, R. B., 1970, Sources and distribution of suspended sediment in northern Chesapeake Bay: *Marine Geology*, v. 9, p. 187-201.

- Biggs, R. B., and Howell, B. A., 1984, The estuary as a sediment trap: alternate approaches to estimating its filtering efficiency, in Kennedy, V. S., ed., The estuary as a filter: New York, Academic Press, p. 107-129.
- Biggs, Robert, B., 1967, The sediments of Chesapeake Bay, in Lauff, G. H., ed., Estuaries, Washington, D.C., American Association for the Advancement of Science, Publication No. 83, p. 239-260.
- Bishop, T. D., 1981, Seasonal abundance of the molluscan fauna of a *Juncus roemerianus* and a *Spartina cynosuroides* brackish tidal marsh in Mississippi: Master's thesis, University of Southwestern Louisiana, 79 p.
- Bloodgood, D. W., and Mortensen, J. E., 1953, Fourteenth annual report of the silt load of Texas streams, 1951-1952 and a summary of silt studies made in Texas: Board of Water Engineers and U. S. Department of Agriculture, Soil Conservation Service, Austin, Texas, 53 p., 3 appendices.
- Bloom, S. A., Simon, J. L., and Hunter, V. D., 1972, Animal-sediment relations and community analysis of a Florida estuary: Marine Biology, v. 13, p. 43-56.
- Bodine, B. R., 1969, Hurricane surge frequency estimated for the Gulf Coast of Texas: U. S. Army Corps of Engineers, Coastal Engineering Research Center Technical Memorandum 26, 32 p.
- Boesch, D. F., 1977, A new look at the zonation of benthos along the estuarine gradient, in Coull, B. C., ed., Ecology of marine benthos: Columbia, SC, University of South Carolina Press, p. 245-266.
- Boesch, D. F., 1973, Classification and community structure of macrobenthos in the Hampton Roads area, Virginia: Marine Biology, v. 21, p. 226-244.
- Boesch, D. F., 1972, Species diversity of marine macrobenthos in the Virginia area: Chesapeake Science, v. 13, no.3, p. 206-211.
- Boesch, D. F., Day, J. W., Jr., and Turner, R. E., 1984, Deterioration of coastal environments in the Mississippi Deltaic Plain: options for management, in Kennedy, V. S., ed., The estuary as a filter: Orlando, Academic Press, p. 447-466.
- Boesch, D. F., Levin, D., Nummedal, D., and Bowles, K., 1983, Subsidence in coastal Louisiana: causes, rates, and effects on wetlands: U. S. Fish and Wildlife Service, Division of Biological Services, Washington, D. C., FWS/OBS-83/26, 30 p.
- Boesch, D. F., and Turner, R. E., 1984, Dependence of fishery species on salt marshes: the role of food and refuge: Estuaries, v. 7, no. 4a, p. 460-468.
- Borey, R. B., Harcombe, P. A., and Fisher, F. M., 1983, Water and organic carbon fluxes from an irregularly flooded brackish marsh in the upper Texas coast, USA: Estuarine Coastal and Shelf Science, v. 16, no. 4, p. 379-402.

- Boucot, A. J., 1981, Principles of benthic marine paleoecology: Academic Press, New York, 463 p.
- Bouma, A. H., and Bryant, W. R., 1969, Rapid delta growth in Matagorda Bay, Texas, Coastal lagoons, a symposium: University National Autonomo de Mexico, p. 171-190.
- Boyden, C. R., and Little, Colin, 1973, Faunal distributions in soft sediments of the Severn estuary: Estuarine and Coastal Marine Science, v. 1, p. 203-223.
- Brehmer, M. L., 1965, Turbidity and siltation as forms of pollution: Journal of Soil and Water Conservation, v. 20, no. 4, p.132-133.
- Brown, L. F. Jr., Morton, R. A., McGowen, J. H., Kreitler, C. W., and Fisher, W. L., 1974, Natural hazards of the Texas Coastal Zone: The University of Texas at Austin, Bureau of Economic Geology, 13 p., 7 maps.
- Brown, L. F., Jr., Brewton, J. L., Evans, T. J., McGowen, J. H., White, W. A., Groat, C. G., and Fisher, W. L., 1980, Environmental geologic atlas of the Texas Coastal Zone--Brownsville-Harlingen area: The University of Texas at Austin, Bureau of Economic Geology, 140 p., 9 maps.
- Brown, L. F., Jr., Brewton, J. L., McGowen, J. H., Evans, T. J., Fisher, W. L., and Groat, C. G., 1976, Environmental geologic atlas of the Texas Coastal Zone--Corpus Christi area: The University of Texas at Austin, Bureau of Economic Geology, 123 p., 9 maps.
- Brown, L. F., Jr., McGowen, J. H., Evans, T. J., Groat, C. G., and Fisher, W. L., 1977, Environmental geologic atlas of the Texas Coastal Zone--Kingsville area: The University of Texas at Austin, Bureau of Economic Geology, 131 p., 9 maps.
- Brune, G. M., 1953, Trap efficiency of reservoirs: American Geophysical Union Transactions, v. 34, p. 407-418.
- Burrell, D. C., and Schubel, J. R., 1977, Seagrass ecosystem oceanography, in McRoy, C. P., and Helfferich, C., eds., Seagrass ecosystems: New York, Marcel Dekker Inc. p. 195-232.
- Byrne, J. R., 1975, Holocene depositional history of Lavaca Bay, central Texas Gulf Coast: The University of Texas at Austin, Ph.D. dissertation, 149 p.
- Cahoon, D. R., DeLaune, R. D., Knaus, R. M., and R. E. Turner, 1987, Marsh sediment accretion rates in vicinity of man-made canals and natural waterways: U.S. Department of Interior Minerals Management Service, Proceedings, Seventh Annual Gulf of Mexico Information Transfer Meeting, Nov. 1986, OCS Study MMS 87-0058.
- Cammen, L. M., 1976, Macroinvertebrate colonization of *Spartina* marshes artificially established on dredge spoil: Estuarine and Coastal Marine Science, v. 4, p. 357-372.

- Cammen, L. M., 1979, The macro-infauna of a North Carolina salt marsh: The American Midland Naturalist, v. 102, no. 2, p. 244-253.
- Carney, R. S., 1981, Bioturbation and biodeposition, in Boucot, A. J., ed., Principles of Benthic Marine Paleoecology: New York, Academic Press, p. 357-399.
- Carr, J. T., Jr., 1967, The climate and physiography of Texas: Texas Water Development Board Report 53, 27 p.
- Carriker, M. R., 1967, Ecology of estuarine benthic invertebrates: a perspective, in Lauff, G. H., ed., Estuaries, Publication 83, American Association for the Advancement of Science, p. 442-487.
- Cary, R. S., 1981, Bioturbation and biodeposition, in Boucot, A. J., ed., Principles of Benthic Marine Paleoecology: New York, Academic Press, p. 357-399.
- Chapman, V. J., 1960, Salt marshes and salt deserts of the world: Leonard Hill, New York, 392 p.
- Chapman, V. J., 1974, Salt marshes and salt deserts of the world: J. Cramer, Lehre, 392 p.
- Chebreck, R. H., 1972, Vegetation, water and soil characteristics of the Louisiana coastal region: Louisiana State University and Agricultural and Mechanical College, Bulletin no. 664, 72 p.
- Chin, J. L., unpublished manuscript, 1977, Distribution of marine grasses in southern Laguna Madre, Texas: The University of Texas at Austin, Bureau of Economic Geology, 45p.
- Christman, S. P., Kochman, H. I., and Lippincott, 1978, An annotated bibliography of the fish and wildlife resources of Galveston Bay, Texas: U. S. Fish and Wildlife Service, National Coastal Ecosystems Teams, National Space Technology Laboratories, Mississippi, FWS/OBS-78/71, 2 volumes, volume 1, 647 p.; volume 2, 342 p.
- Church, T. M., Biggs, R. B., and Sharma, P., 1987, The birth and death of salt marshes: Geochemical evidence for sediment accumulation and erosion: EOS Transactions, American Geophysical Union, v. 68, p. 305 (abstract).
- Churchill, M. A., 1948, discussion of "Analysis and Use of Reservoir Sedimentation Data," by L. C. Gottschalk: Proceedings of the Federal Interagency Sedimentation Conference, Denver, U. S. Bureau of Reclamation, p. 139-140.
- Clanton, U. S., 1968, Sorption and release of strontium-89 and cesium-137 by recent sediments of the Guadalupe River of Texas, The University of Texas at Austin, Ph.D. Dissertation, 87 p.
- Coastal Ecosystems Management Inc., 1972, Environmental assessment of the Trinity River discharge on productivity in Trinity Bay: Final report prepared for Fort Worth District, Army Corps of Engineers, contract no. DACW63-72-C-0142, 73 p.

- Coleman, J., and Gagliano, S. M., 1964, Cyclic sedimentation in the Mississippi River Deltaic Plain: Gulf Coast Association of Geological Societies Transactions, v. 14, p. 67-80.
- Collier, W. J., and Hedgpeth, J., 1950, An introduction to the hydrography of tidal waters of Texas: University of Texas, Institute of Marine Science Publications, v. 1, no. 2, p. 125-194.
- Commuto, J. A., 1982, Importance of predation by infaunal polychaetes in controlling the structure of a soft-bottom community in Maine, USA: Marine Biology, v. 68, p. 77-81.
- Conner, J. V., and Truesdale, F. M., 1973, Ecological implications of a fresh water impoundment in a low salinity marsh, in, Chabreck, R. H., ed., Proceedings of the Coastal Marsh and Estuary Management Symposium: Louisiana State University, Baton Rouge, Louisiana, p. 259-276.
- Conover, J. T., 1964, The ecology, seasonal periodicity, and distribution of benthic plants in some Texas lagoons: Botanica Marina, v. 7, fascicle 1-4, p. 4-41.
- Cook, H. M., 1967, Suspended-sediment load of Texas streams: compilation report, October 1961 - September 1963: Texas Water Development Board, Report 45, 61 p.
- Cook, H. M., 1970, Suspended-sediment load of Texas streams: compilation report, October 1963 - September 1965: Texas Water Development Board, Report 106, 61 p.
- Copeland, B. J., 1970, Estuarine classification and responses to disturbances: Transactions, American Fisheries Society, v. 4, p. 826-835.
- Cornelius, S. E., 1975, Food choice of wintering redhead ducks (*Aythya americana*) and utilization of available resources in lower Laguna Madre, Texas: Texas A&M University, Master's thesis, 121 p.
- Craig, G. Y., and Jones, N. S., 1966, Marine benthos, substrate and paleoecology: Palaeontology, v. 9, p. 30-38.
- Craig, M. A., and Bright, T. J., 1986, Abundance, age distributions and growth of the Texas hard clam, *Mercenaria mercenaria texana* in Texas bays: Contributions in Marine Science, v. 29, p. 59-72.
- Crisp, D. J., 1984, Energy flow measurements, in Holme, N. A., and McIntyre, A. D., eds., Methods for the study of marine benthos: London, Blackwell Scientific Publications, p. 284-372.
- Cunningham, Robert, 1981, Atchafalaya Delta: Subaerial development, environmental implications, and resource potential, in Cross, R. D., and Williams, D. L., Proceedings of the National Symposium on Freshwater Inflow to Estuaries, Volume I: U. S. Fish and Wildlife Service Office of Biological Services, FWS/OBS-81/04, p. 349-364.

- Curry, J. R., 1960, Sediments and history of Holocene transgression, continental shelf, northwest Gulf of Mexico, in: Shepard, F. P., Phleger, F. B., and van Andel, T. H., eds., Recent Sediments, northwest Gulf of Mexico: American Association of Petroleum Geologists, p. 221-226.
- Currington, H. W., Wells, D. M., Masch, R. D., Copeland, B. J., and Gloyna, E. F., 1966, Return flows - impact on the Texas bay systems: Report prepared for the Texas Water Development Board by Bryant-Currington, Engineers, Austin, Texas, 419 p.
- Curtis, W. F., Culbertson, J. K., and Chase, E. B., 1973, Fluvial-sediment discharge to the oceans from the conterminous United States: U. S. Geological Survey Circular 670, 17 p.
- Daiber, F. C., 1982, Animals of the tidal marsh: New York, Van Nostrand Reinhold Company, 422 p.
- Dalrymple, D. W., 1964, Recent sedimentary facies of Baffin Bay, Texas: Rice University, Ph.D. dissertation, 192 p.
- Davies, D. K., and Moore, R. W., 1970, Dispersal of Mississippi sediment in the Gulf of Mexico: Journal of Sedimentary Petrology, v. 40, p. 339-353.
- Davis, R. A., and Fox, W. T., 1972, Coastal dynamics along Mustang Island, Texas: Western Michigan University Technical Report No. 9, ONR Contract No. 388-092, 68 p.
- DeLaune R. D., Patrick, W. H., Jr., and Buresh, R. J., 1978, Sedimentation rates determined by ¹³⁷Cs dating in a rapidly accreting salt marsh: Nature, v. 275, p. 532-533.
- DeLaune, R. D., Baumann, R. H., and Gosselink, J. G., 1983, Relationships among vertical accretion, coastal submergence, and erosion in a Louisiana Gulf coast marsh: Journal of Sedimentary Petrology, v. 53, no. 1, p. 147-157.
- DeLaune, R. D., Reddy, C. N., and Patrick, W. H., Jr., 1981, Accumulation of plant nutrients and heavy metals through sedimentation processes and accretion in a Louisiana salt marsh: Estuaries, v. 4, no. 4, p. 328-334.
- DeLaune, R. D., Smith, C. J., Patrick, W. H., Jr., and Roberts, H. H., 1987, Rejuvenated marsh and bay-bottom accretion on the rapidly subsiding coastal plain of U.S. Gulf Coast: a second-order effect of the emerging Atchafalaya Delta: Estuarine, Coastal and Shelf Science, v. 25, p. 381-389.
- DeLaune, R. D., and Patrick, W. H., Jr., 1980, Rate of sedimentation and its role in nutrient cycling in a Louisiana salt marsh, in Hamilton, Peter, and Macdonald, K. B., eds., Estuarine and wetland processes: Plenum Press, New York, p. 401-412.
- Deegan, L. A., Day, J. W., Jr., Gosselink, J. G., Yanez-Arancibia, A., Chavez, G. S., and Sanchez-Gil, P., 1986, Relationships among physical characteristics, vegetation distribution and fisheries yield in Gulf of Mexico estuaries, in Wolfe, D. A., ed., Estuarine variability: Academic Press, Inc., Orlando, p. 83-100.

- Diener, R. A., 1975, Cooperative Gulf of Mexico estuarine inventory and study-Texas: area description: National Oceanic and Atmospheric Administration, Technical Report, National Marine Fisheries Service Circular 393, 129 p.
- Donaldson, A. C., Martin, R. H., and Kanes, W. H., 1970, Holocene Guadalupe delta of Texas Gulf Coast, in Morgan, J. P., ed., Deltaic sedimentation: modern and ancient: Society of Economic Paleontologists and Mineralogists Special Publication 15, p. 107-137.
- Dougherty, J. P., 1979, Suspended-sediment load of Texas streams, compilation report October 1971-September 1975: Texas Department of Water Resources Report 233, 82 p.
- Driscoll, E. G., 1967, Attached epifauna-substrate relations: Limnology and Oceanography, v.12, p. 633-641.
- Driscoll, E. G., 1975, Sediment-animal-water interaction, Buzzards Bay, Massachusetts: Journal of Marine Research, v. 33, no. 3, p. 275-302.
- Edwards, J. M., and Frey, R. W., 1977, Substrate characteristics within a Holocene salt marsh, Sapelo Island, Georgia: Senckenberg. Marit., v. 9, p. 215-259.
- Elliot, A. B., 1958, Recent sediments of Corpus Christi and Nueces Bays, Nueces County, Texas: The University of Texas at Austin, Masters Thesis, 169 p.
- Emery, K. O., and Stevenson, R. E., 1957, Estuaries and lagoons: Geological Society of America Memoir 67, v. 1, p. 673-750.
- Espey, Huston and Associates, Inc., 1981, Environmental report, proposed deepwater channel and multipurpose terminal construction and operation near Brownsville, Texas: Austin, Texas, Espey, Huston and Associates, Inc., v. 6, p. K1-K140, v. 7, p.11-146.
- Espey, Huston and Associates, Inc., 1978, Hydrological and biological studies of the Colorado River delta, Texas: Texas Department of Water Resources, LP-79, 165 p.
- Etkins, R., and Epstein, E. S., 1982, The rise of global mean sea level as an indication of climate change: Science, v. 215, no. 4530, p. 287-289.
- Everett, J. R., and Reid, W. M., 1981, Active faults in the Houston, Texas area as observed on Landsat imagery, in Etter, E. M., ed., Houston area environmental geology: surface faulting, ground subsidence, hazard liability: Houston Geological Society, p. 13-27.
- Fagg, D. B., 1957, The recent marine sediments and Pleistocene surface of Matagorda Bay, Texas: Gulf Coast Association of Geological Societies, Transactions, v. 7, p. 119-133.
- Failing, M. S., 1969, Comparison of Trinity River terraces and gradients with other Texas Gulf Coast rivers, in Lankford, R. R., and Rogers, J. J. W., eds., Holocene geology of the Galveston Bay area: Houston Geological Society, Delta Study Group, p. 85-92.

- Faris, O. A., 1933, The silt load of Texas streams: U.S. Department of Agriculture Technical Bulletin No. 382, Washington, D.C., 71 p.
- Fisher, W. L., 1969, Facies characterization of Gulf Coast basin delta systems, with some Holocene analogues: Gulf Coast Association of Geological Societies Transactions, v. 19, p. 239-261.
- Fisher, W. L., Brown, L. F., Jr., McGowen, J. H., and Groat, C. G., 1973, Environmental geologic atlas of the Texas Coastal Zone-Beaumont-Port area: The University of Texas at Austin, Bureau of Economic Geology, 93 p., 9 maps.
- Fisher, W. L., Brown, L. F., Scott, A. J., McGowen, J. H., 1969, Delta systems in the exploration of oil and gas--A research colloquium: The University of Texas at Austin, Bureau of Economic Geology, 78 p., 168 figs., references.
- Fisher, W. L., McGowen, J. H., Brown, L. F., Jr., and Groat, C. G., 1972, Environmental geologic atlas of the Texas Coastal Zone-Galveston-Houston area: The University of Texas at Austin, Bureau of Economic Geology, 91 p., 9 maps.
- Fisher, W. L., and Brown, L. F., Jr., 1972, Clastic depositional systems--a genetic approach to facies analysis: Annotated outline and bibliography: The University of Texas at Austin, Bureau of Economic Geology, 211 p.
- Fisk, H. N., 1959, Padre Island and the Laguna Madre flats, coastal south Texas: Louisiana State University, 2d Coastal Geography Conference, April 1959, p. 103-151.
- Fisk, H. N., 1960, Recent Mississippi River sedimentation and peat accumulation: Cong. Av. Etudes de Stratigraphie et de Geologie du Carbonifere, 4th, Sept. 15-20, 1958, Heerlen, Compte rendu, p. 187-199, Maastricht, Netherlands.
- Flessa K. F., Constantine, K. J., and Cushman, M. K., 1977, Sedimentation rates in a coastal marsh determined from historical records: Chesapeake Science, v. 18, no. 2, p. 172-176.
- Flint, R. W., 1985, Long-term estuarine variability and associated biological response: Estuaries, v. 8, no. 2A, p. 158-169.
- Flint, R. W., 1984, Phytoplankton production in Corpus Christi Bay: Contributions in Marine Science, v. 27, p. 65-83.
- Flint, R. W., Rabalais, S. C., and Kalke, R. D., 1981, Quantification of extensive freshwater input to estuarine benthos: The University of Texas Marine Science Institute, Port Aransas, report to Texas Department of Water Resources, contract no. IAC(80-81), 55 p.
- Flint, R. W., Rabalais, S., and Kalke, R., 1982, Estuarine benthos and ecosystem functioning, in Davis, J. R., ed., Proceedings of the symposium on recent benthological investigations in Texas and adjacent states: Texas Academy of Science 85th annual meeting, San Angelo, Texas, p. 185-201.

- Flint, R. W., and Kalke, R. D., 1985, Benthos structure and function in a South Texas estuary: *Contributions in Marine Science*, v. 28, p. 185-201.
- Frey, R. W., and Basan, P. B., 1985, Coastal salt marshes, in Davis, R. A., Jr., ed., *Coastal sedimentary environments*, second revised, expanded edition: New York, Springer-Verlag, p. 225-301.
- Frishman, S. A., 1969, *Geochemistry of oolites, Baffin Bay, Texas*: The University of Texas at Austin, Master's thesis, 82 p.
- Fulton, K. J., 1975, *Subsurface stratigraphy depositional environments, and aspects of reservoir continuity--Rio Grande delta, Texas*: University of Cincinnati, Ph. D. Dissertation, 314 p.
- Funicelli, N. A., 1984, Assessing and managing effects of reduced freshwater inflow to two Texas estuaries, in Kenedy, V. S., ed., *The estuary as a filter*: Orlando, Academic Press, p. 435-446.
- Funicelli, N. A., and Benson, N. G., 1983, Choke Canyon Reservoir: Estuarine impacts and management, in, Magoon, O. T., and Converse, Hugh, *Coastal Zone '83, Volume III*: American Society of Civil Engineers, New York, p. 2866-2876.
- Gabrysch, R. K., 1984, Ground-water withdrawals and land-surface subsidence in the Houston-Galveston region, Texas, 1906-1980: Texas Department of Water Resources Report 287, 64 p.
- Gabrysch, R. K., 1969, Land-surface subsidence in the Houston-Galveston region, Texas: United Nations Educational, Scientific and Cultural Organization (UNESCO), *Studies and Reports in Hydrology, Land Subsidence Symposium*, v. 1, p. 43-54.
- Gabrysch, R. K., and Bonnet, C. W., 1976, Land-surface subsidence in the area of Moses Lake near Texas City, Texas: U. S. Geological Survey, Water Resources Division, Austin, Texas, Rep. No. USGS/WRD/WRI-77/007, USGS/WRI-76-32, 90 p.
- Gagliano, S. M., Kwon, H. J., and van Beek, J. L., 1970, Deterioration and restoration of coastal wetlands: *Proceedings of Twelfth Conference on Coastal Engineering*, Washington D.C., ASCE, v. 3, p. 1767-1781.
- Gagliano, S. M., Meyer-Arendt, K. J., and Wicker, K. M., 1981, Land loss in the Mississippi River deltaic plain: *Gulf Coast Association of Geological Societies, Transactions*, v. 31, p. 271-273.
- Galtsoff, P. S., 1931, Survey of oyster bottoms in Texas: U. S. Bureau of Fisheries, investigational report 6, 30 p.
- Gillard, R. M., 1974, Distribution, abundance, and species diversity of macrobenthic and meiobenthic invertebrates in relation to Houston Ship Channel pollution in upper Galveston Bay and Tabbs Bay, Texas: Texas A&M University, Ph.D. Dissertation, 186 p.

- Gilmore, G., Dailey, J., Garcia, M., Hannebaum, N., and Means, J., 1976, A study of the effects of fresh water on the plankton, benthos, and nekton assemblages of the Lavaca Bay System, Texas: Texas Parks and Wildlife Department, Technical report to the Texas Water Development Board, January, 1973-June, 1975, 113 p.
- Gilmore, G., and Trent, L., 1974, Abundance of benthic macroinvertebrates in natural and altered estuarine areas: Seattle, Washington, U.S. Department of Commerce, National Oceanic and Atmospheric Administration Technical Report NMFS SSRF-677, 13 p.
- Ginsburg, R. N., and Lowenstam, H. A., 1958, The influence of marine bottom communities on the depositional environment of sediments: *Journal of Geology*, vol. 62, p. 249-265.
- Gleason, M. L., Elmer, D. H., Pien, N. C., and Fisher, J. S., 1979, Effects of stem density upon sediment retention by salt marsh cord grass, *Spartina alterniflora* Loisel: *Estuaries*, v. 2, no. 4, p. 271-273.
- Goldman, C. R., 1971, Biological implications of reduced freshwater flows on the San Francisco Bay-Delta system, in Seckler, D., ed., *California water, a study in resource management*: University of California Press, Berkeley, p. 109-124.
- Gornitz, V., Lebedeff, S., and Hansen, J., 1982, Global sea level trend in the past century: *Science*, v. 215, p. 1611-1614.
- Gosselink, J. G., 1984, The ecology of delta marshes of coastal Louisiana: a community profile: U. S. Fish and Wildlife Service, FWS/OBS-84/09, 134 p.
- Gosselink, J. G., Cordes, C. L., and Parsons, J. W., 1979, An ecological characterization study of the Chenier Plain coastal ecosystem of Louisiana and Texas, v. 1, narrative report: Washington, D. C., U.S. Department of the Interior, Fish and Wildlife Service, Office of Biological Services, FWS/OBS-78/9 through 78/11, 302 p.
- Gould, H. R., and Morgan, J. P., 1962, Coastal Louisiana swamps and marshlands, in Rainwater, E. H., and Zingula, R. P., eds., *Geology of the Gulf coast and central Texas, and guidebook of excursions*: Houston Geological Society, Houston, Texas, p. 286-341.
- Govin, C. T., Jr., 1973, Sedimentation survey, Lake Buchanan, Texas, 1973: The University of Texas at Austin, Master's Thesis, 105 p. 1 map.
- Grady, J. R., 1981, Properties of sea grass and sand flat sediments from the intertidal zone of St. Andrew Bay, Florida: *Estuaries*, v. 4, no. 4, p. 335-344.
- Gray, J. S., 1974, Animal-sediment relationships, in Barnes, H., ed., *Oceanography and Marine Biology-an Annual Review*, v. 12, p. 335-344.
- Greiner, J. H., 1982, Erosion and sedimentation by water in Texas, average annual rates estimated in 1979: Texas Department of Water Resources, Report 268, 145 p.

- Gulf Universities Research Corporation. Case studies of estuarine sedimentation and its relation to pollution of the estuarine environment: Report to the Federal Water Pollution Control Administration.
- Gustavson, T. C., and Kreidler, C. W., 1976. Geothermal resources of the Texas Gulf coast--environmental concerns arising from the production and disposal of geothermal waters: The University of Texas at Austin, Bureau of Economic Geology Geological Circular 76-7, 35 p.
- Hackney, C., 1978. Summary of information: relationship of freshwater inflow to estuarine productivity along the Texas coast: U. S. Fish and Wildlife Service, FWS/OBS-78/73, Albuquerque
- Hall, G. L., 1976. Subenvironments of deposition in San Antonio Bay, in Bouma, A. H., ed., Shell dredging and its influence on Gulf Coast environments: Houston, Texas, Gulf Publishing Company Book Division, p. 100-131.
- Hann, R. W., Jr., 1970. Mathematic modeling of Gulf Coast estuaries: Texas A&M University, College Station, Department of Civil Engineering, Water Resources Bulletin, v. 6, no. 3, p. 323-338.
- Hansen, J., Johnson, D., Lacis, A., Lebedeff, S., Lee, P., Rind, D., and Russell, G., 1981. Climate impact of increasing atmospheric carbon dioxide: Science, v. 213, no. 4511, p. 957-966.
- Harlin, M. M., and Thorne-Miller, B., 1982. Seagrass-sediment dynamics of a flood-tidal delta in Rhode Island: Aquatic Botany, vol. 14, p. 127-138.
- Harper, D. H., and Hopkins, S. H., 1976. The effects of oyster shell dredging on macrobenthic and nektonic organisms in San Antonio Bay, in Bouma, A. H., ed., Shell dredging and its influence on Gulf Coast environments: Houston, Texas, Gulf Publishing Company, p. 232-279.
- Harrison, E. Z., and Bloom, A. L., 1977. Sedimentation rates on tidal salt marshes in Connecticut: Journal of Sedimentary Petrology, v. 47, no. 4, p. 1484-1490.
- Hatton, R. S., DeLaune, R. D., and Patrick, W. H., Jr., 1983. Sedimentation, accretion, and subsidence in marshes of the Barataria Basin, Louisiana: Limnology and Oceanography, v. 28, no. 3, p. 494-502.
- Hatton, R. S., Patrick, W. H., Jr., and DeLaune, R. D., 1982. Sedimentation, nutrient accumulation, and early diagenesis in Louisiana Barataria basin coastal marshes, in Kennedy, V. S., ed., Estuarine comparisons: Academic Press, New York, p. 255-267.
- Hauck, L. M., and Ward, G. H., 1980. Hydrodynamic-mass transfer model of deltaic systems, in Hamilton, Peter, and Macdonald, K. B., eds., Estuarine and Wetland Processes, with emphasis on modeling: Plenum, New York, p. 247-268.
- Haven, D. S., and Morales-Alamo, R., 1966. Aspects of biodeposition by oysters and other invertebrate filter feeders: Limnology and Oceanography, v. 11, p. 487-498.

- Haven, D. S., and Morales-Alamo, R., 1968, Occurrence and transport of faecal pellets in suspension in a tidal estuary: *Sedimentary Geology*, v. 2, p. 125-140.
- Haven, D., and Morales-Alamo, R., 1972, Biodeposition as a factor in sedimentation of fine suspended solids in estuaries, in Nelson, B., ed., *Environmental framework of coastal plain estuaries: Geological Society of America Memoir No.133*, p. 121-130.
- Hayes, M. O., 1967, Hurricanes as geological agents: case studies of Hurricane Carla, 1961, and Cindy, 1963: The University of Texas at Austin, Bureau of Economic Geology Report of Investigations 61, 56 p.
- Hayes, M. O., 1978, Impact of hurricanes on sedimentation in estuaries, bays, and lagoons, in Wiley, M. L., ed., *Estuarine interactions: Academic Press, New York*, p. 323-346.
- Hayes, M. O., 1965, Sedimentation on a semiarid, wave-dominated coast (South Texas) with emphasis on hurricane effects: The University of Texas at Austin, Ph.D. Dissertation, 350 p.
- Heck, K. L., Jr., and Wetstone, G. S., 1977, Habitat complexity and invertebrate species richness and abundance in tropical seagrass meadows: *Journal of Biogeography*, v. 4, p. 135-142.
- Henry, V. J., 1956, Investigation of shoreline-like features in the Galveston Bay region, Texas: Texas A&M University, Department of Oceanography, Technical Report, Project 24, 76 p.
- Hicks, S. D., 1978, An average geopotential sea-level series for the United States: *Journal of Geophysical Research*, v. 83, p. 1377-1379.
- Hildebrand, H., and King, D., 1978, A biological study of the Cayo del Oso and the Pita Island area of the Laguna Madre, final report, 1972-1978, volume I: Corpus Christi, Texas, Central Power and Light Company, 253 p.
- Hines, A. H., and Comtois, K. L., 1985, Vertical distribution of infauna in sediments of a subestuary of central Chesapeake Bay: *Estuaries*, v. 8, no. 3, p. 296-304.
- Hobbie, J. E., and Copeland, B. J., 1977, Effects and control of nutrients in estuarine ecosystems, in *Estuarine pollution and control and assesement. Proceedings of a conference, volume 1: U. S. Environmental Protection Agency, Washington, D.C., U.S. Government Printing Office*, p. 257-274.
- Hobbs, R. S., 1987, Clastic deltas, analysis of bed load sedimentation patterns, and river dynamics of the Brazos River, central Texas: Baylor University, Master's thesis, 114 p.
- Holdahl, S. R., and Morrison, N. L., 1974, Regional investigations of vertical crustal movements in the U.S., using precise relevelings and mareograph data: *Tectonophysics*, v. 23, p. 373-390.

- Holland, A. F., 1985, Long-term variation of macrobenthos in a mesohaline region of Chesapeake Bay: *Estuaries*, v. 8, no. 2A, p. 93-113.
- Holland, A. F., Mountford, N. K., and Mihursky, J. A., 1977, Temporal variation in upper bay mesohaline benthic communities: mesohaline benthic communities: 1. the 9-m mud habitat: *Chesapeake Science*, v. 18, no. 4, p. 370-378.
- Holland, J. S., Maciolek, N. J., Kalke, R. D., and Oppenheimer, C. H., 1975, A benthos and plankton study of the Corpus Christi, Copano and Aransas Bay systems: report on data collected during the period July 1974-May 1975 and summary of the three-year project: The University of Texas at Port Aransas Marine Science Institute, Final report to the Texas Water Development Board, 171 p.
- Holland, J. S., Maciolek, N. J., and Oppenheimer, C. H., 1973, Galveston Bay benthic community structure as an indicator of water quality: *Contributions in Marine Science*, v. 17, p. 169-188.
- Holliday, Barry, 1973, Bay circulation, in Ahr, W. M., project coordinator, Resource evaluation studies on the Matagorda Bay area, Texas: Texas A&M University Sea Grant publication, TAMU-SG-74-204, p. 5-20.
- Holmes, C. W., 1982, Geochemical indices of fine sediment transport, northwest Gulf of Mexico: *Journal of Sedimentary Petrology*, v. 52, no. 1, p. 370-421.
- Hoskin, C. M., 1983, Sediment in seagrasses near Link Port, Indian River, Florida: *Florida Scientist*, v. 46, no. 3/4, p. 153-161.
- Hunter, R. E., Watson, R. L., Hill, G. W., and Dickinson, K. A., 1972, Modern depositional environments and processes, northern and central Padre Island, Texas, in Padre Island National Seashore field guide: Gulf Coast Association of Geological Societies, convention field trip, p. 1-17.
- Isphording, W. C., Imsand, D. and Flowers, G. C., 1987, Storm-related rejuvenation of a northern Gulf of Mexico estuary: *Gulf Coast Association of Geological Societies, Transactions*, v. 37, p. 357-370.
- Isphording, W. C., 1986, Apalachicola Bay: dynamic sedimentation in a Gulf Coast estuary: *Transactions of the Gulf Coast Association of Geological Societies*, v. 36, p. 471-488.
- Isphording, W. C., Imsand, F. D., and Isphording, G. W., 1984, Identification of short-term changes in sediment depositional rates: importance in environmental analysis and impact investigations: *Gulf Coast Association of Geological Societies Transactions*, v. 34, p. 69-84.
- Jacobs, J., 1985, Fluvial responses to hydrologic changes on the Red River in northeast Texas: *Gulf Coast Association of Geological Societies Transactions*, v. 35, p. 409-414.
- Johnson, P. J., 1980, Seasonal variation in benthic community structure in Mobile Bay, Alabama: University of Alabama, Birmingham, Master's Thesis, 96 p.

- Johnson, R. G., 1971, Animal-sediment relations in shallow water benthic communities: *Marine Geology*, v.11, p.93-104.
- Johnson, R. G., 1970, Variations in diversity within benthic communities: *American Naturalist*, vol. 104, p. 285-300.
- Johnston, J. B., and Ader, R. A., 1983, The use of a GIS for Gulf of Mexico wetland change: in Magoon, O. T., and Converse, H., eds., *Coastal Zone '83*, Volume I: American Society of Civil Engineers, New York, p. 362-371.
- Jones, R. S., principal investigator, and others, 1986, Studies of freshwater inflow effects on the Lavaca River delta and Lavaca Bay, Texas: Final report to Texas Water Development Board, contract no. IAC (86-87) 0757, 387 p.
- Jordan, T. E., and Valiela, I., 1982, A nitrogen budget of the ribbed mussel, *Geukensia demissa*, and its significance in nitrogen flow in a New England salt marsh: *Limnology and Oceanography*, v. 27, no. 1, p. 75-90.
- Kalke, R. D., 1986, Benthos, in Jones, R. S., and others, Studies of freshwater inflow effects on the Lavaca River Delta and Lavaca Bay, Texas: Final report to Texas Water Development Board, contract no. IAC(86-87)0757, p. 5.1-5.41.
- Kane, H. E., 1959, Late Quaternary geology of Sabine Lake and vicinity, Texas and Louisiana: *Gulf Coast Association of Geological Societies Transactions*, v. 9, p. 225-235.
- Kane, H. E., 1966, Sediments of Sabine Lake, the Gulf of Mexico and adjacent water bodies, Texas-Louisiana: *Journal of Sedimentary Petrology*, v. 36, p. 608-619.
- Kanes, W. H., 1970, Facies and development of the Colorado River delta in Texas, in Morgan, J. P., and Shaver R. H., eds., *Deltaic sedimentation, modern and ancient*: Society of Economic Paleontologists and Mineralogists Special Publication No. 15, p. 78-106.
- Kanes, W. H., 1970, Facies and development of the Colorado River delta in Texas: Society of Economic Paleontologists and Mineralogists, Special Publication No. 15, p. 78-106.
- Kanes, W. H., 1965, Facies and development of the delta of the Colorado River, Texas: West Virginia University, Morgantown, West Virginia, Ph.D. Dissertation, 106 p.
- Kemp, W. M., Boynton, W. R., Twilley, R. R., Stevenson, J. C., and Ward, L. G., 1984, Influences of submersed vascular plants on ecological processes in upper Chesapeake Bay, in Kennedy, V. S., ed., *The estuary as a filter*: New York, Academic Press, p. 367-394.
- Kenworthy, W. J., Zieman, J. C., and Thayer, G. W., 1982, Evidence for the influence of seagrasses on the benthic nitrogen cycle in a coastal plain estuary near Beaufort, North Carolina (USA): *Oecologia*, v. 54, p. 152-158.

- Kier, R. S., and White, W. A., 1978, Land and water resources of the Corpus Christi area, Texas: The University of Texas at Austin, Bureau of Economic Geology Report of Investigations No. 95, 22 p.
- Kraeuter, J. N., 1976, Biodeposition by salt marsh invertebrates: *Marine Biology*, v. 35, p. 215-223.
- Kraeuter, J. N., and Wolf, P. L., 1974, The relationship of marine macroinvertebrates to salt marsh plants, in Reimold, R. J., and Queen W. H., eds., *Ecology of Halophytes*: New York, Academic Press, Inc., p. 449-462.
- Kranck, Kate, 1984, The role of flocculation in the filtering of particulate matter, in Kennedy, V. S., ed., *The estuary as a filter*: New York, Academic Press, p. 159-175.
- Kreitler, C. W., 1977, Faulting and land subsidence from ground-water and hydrocarbon production, Houston-Galveston, Texas: The University of Texas at Austin, Bureau of Economic Geology Research Note 8, 22 p.
- Lane, E. W., and Borland, W. M., 1951, Estimating bed load: *Transactions American Geophysical Union*, 32 (1), p. 121-123.
- Larkin, T. J., and Bomar, G. W., 1983, Climatic atlas of Texas: Texas Department of Water Resources, LP 87-04, 74 p.
- Larsen, P. F., 1979, The shallow-water macrobenthos of a northern New England estuary: *Marine Biology*, v. 55, no. 1, p. 69-78.
- LeBlanc, R. J., and Hodgson, W. D., 1959, Origin and development of the Texas shoreline, in Russell, R. J., chm., *Coastal geography conference*, 2nd: Baton Rouge, Louisiana State University, Coastal Studies Institute, p. 57-101.
- Leopold, L. B., Wolman, M. G., and Miller, J. P., 1964, *Fluvial processes in geomorphology*: W. H. Freeman and Company, San Francisco, 522 p.
- Letzsch, W. S., and Frey, R. W., 1980, Deposition and erosion in a Holocene salt marsh, Sapelo Island, Georgia: *Journal of Sedimentary Petrology*, v. 50, no. 2, p. 529-542.
- Levinton, J. S., 1977, Ecology of shallow water deposit-feeding communities, Quisset Harbor, Massachusetts, in Coull, B. C., ed., *Ecology of Marine Benthos*: University of South Carolina Press, Columbia, South Carolina, p. 191-227.
- Lewis, F. G., and Stoner, A. W., 1983, Distribution of macrofauna within seagrass beds: an explanation for patterns of abundance: *Bulletin of Marine Science*, v. 33, n. 2, p. 296-304.
- Lie, U., and Evans, R., 1973, Long-term variability in the structure of subtidal benthic communities in Puget Sound, Washington, USA: *Marine Biology*, v. 21, p. 122-126.

- Liebbrand, N. F., 1987, Estimated sediment deposition in Lake Corpus Christi, Texas, 1972-85: Austin, Texas, U. S. Geological Survey open-file report 87-239, 26 p.
- Lohse, E. A., 1955, Dynamic geology of the modern coastal region, northwest Gulf of Mexico, in Hough, J. L. and Menard, H. W., Jr., eds., Finding ancient shorelines-a symposium: Society of Economic Paleontologists and Mineralogists Special Publication No. 3, p. 99-105.
- Lohse, E. A., 1952, Shallow-marine sediments of the Rio Grande Delta: The University of Texas at Austin, Ph.D. Dissertation, 113 p.
- Loi, T-n., and Wilson, B. J., 1979, Macroinfaunal structure and effects of thermal discharges in a mesohaline habitat of Chesapeake Bay, near a nuclear power plant: Marine Biology, v. 55, no. 1, p. 3-16.
- Longbottom, M. R., 1970, The distribution of *Arenicola marina* (L.) with particular reference to the effects of particle size and organic matter of the sediments: Journal of Experimental Marine Biology and Ecology, v. 3, p. 138-157.
- Looney, R. M., and Baker, V. R., 1977, Late Quaternary geomorphic evolution of the Colorado River on the inner Texas Coastal Plain: Gulf Coast Association of Geological Societies, Transactions, v. 27, p. 323-333.
- Loosanoff, V. L., 1962, Effects of turbidity on some larval and adult bivalves: Proceedings Gulf and Caribbean Fisheries Institute, 14th annual session, Miami Beach, Fla., p. 80-95.
- Love, K. M., Chafetz, H. S., Bruno, L., and Buczynski, C., 1985, Bay margin sand distribution, north Texas coast: A model for sediment distribution in microtidal bays and lagoons: Gulf Coast Association of Geological Societies Transactions, v. 35, p. 439-443.
- Lowery, G. H., 1974, The mammals of Louisiana and its adjacent waters: Louisiana State University Press, Baton Rouge, 565 p.
- Lund, E. J., 1957, Self-silting by the oyster and its significance for sedimentation geology: Publications of the Institute of Marine Science, v. 4, no. 2, p. 320-327.
- Lynts, G. W., 1966, Relationship of sediment-sized distribution to ecologic factors in Buttonwood Sound, Florida Bay: Journal of Sedimentary Petrology, v. 36, no. 1, p. 66-74.
- Mackin, J. G., 1961, Canal dredging and silting in Louisiana bays: The University of Texas, Institute of Marine Science Publications, v. 7, p. 262-314.
- Mackin, J. G., and Hopkins, S. H., 1961, Studies on oyster mortality in relation to natural environments and to oil fields in Louisiana: Publications of the Institute of Marine Science, v. 7, p. 1-131.
- Manka, L. L., 1970, A study of the sediments and depositional history of a portion of the delta of the Colorado River, Texas: Texas Christian University, Master's Thesis, 87 p.

- Manka, L. L., and Steinmetz, Richard, 1971, Sediments and depositional history of the southeast lobe of the Colorado River delta, Texas: Gulf Coast Association of Geological Societies, v. 21, p. 309-323.
- Marmer, H. A., 1954, Tides and sea level in the Gulf of Mexico, in Galtsoff, P. S., coordinator, Gulf of Mexico, Its origin, waters, and marine life: United States Department of the Interior, Fish and Wildlife Service, Fishery Bulletin 89, p. 101-118.
- Martinez, A. R., 1969-1970, Coastal hydrographic and meteorological study, coastal fisheries project report, Texas: Texas Parks and Wildlife Department, p. 227-344.
- Martinez, A. R., 1975, Coastal hydrographic and meteorological study, coastal fisheries project report, Texas: Texas Parks and Wildlife Department, p. 100-157.
- Martinez, A. R., 1973, Coastal hydrographic and meteorological study, coastal fisheries project report, Texas: Texas Parks and Wildlife Department, p. 111-172.
- Martinez, A. R., 1971, Coastal hydrographic and meteorological study, coastal fisheries project report, Texas: Texas Parks and Wildlife Department, p. 135-189.
- Martinez, A. R., 1972, Coastal hydrographic and meteorological study, coastal fisheries project report, Texas: Texas Parks and Wildlife Department, p. 146-197.
- Martinez, A. R., 1974, Coastal hydrographic and meteorological study, coastal fisheries project report, Texas: Texas Parks and Wildlife Department, p. 61-108.
- Mathewson, C. C., and Minter, L. L., 1976, Impact of water resource development on coastal erosion, Brazos River, Texas: Texas A&M University, Texas Water Resources Institute, Technical Report No. 77, 85 p.
- Matthews, G. A., Marcin, C. A., and Clements, George, 1975, A plankton and benthos survey of the San Antonio Bay system: Summary report for the period March 1972-July 1974: Texas Parks and Wildlife Department Coastal Fisheries Project Report to the Texas Water Development Board, 76 p.
- Matthews, G. A., Marcin, Carol, and Welch, Douglas, 1974, A plankton and benthos survey of the San Antonio Bay system: Second annual report, August 1972-July 1973: Texas Parks and Wildlife Department Coastal Fisheries Project Report to the Texas Water Development Board, 53 p.
- Maurer, Don, 1969, Pelecypod-sediment association in Tomales Bay, California: The Veliger, v. 11, no. 3, p.243-249.
- Maurer, D., Watling, L., Kinner, P., Leathem, W., and Wethe, C., 1978, Benthic invertebrate assemblages of Delaware Bay: Marine Biology, v. 45, p. 243-249.

- McBee, J. T., and Brehm, W. T., 1982, Spatial and temporal patterns in the macrobenthos of St. Louis Bay, Mississippi: Gulf Research Reports, v. 7, no. 2, p. 115-124.
- McCall, P. L., 1977, Community patterns and adaptive strategies of the infaunal benthos of Long Island Sound: Journal of Marine Research, v. 35, no. 2, p. 221-266.
- McEwen, M. C., 1969, Sedimentary facies of the modern Trinity delta, in Lankford, R. R., and Rogers, J. J. W., Holocene geology of the Galveston Bay area: Houston Geological Society, Chapter 3, p. 53-77.
- McGowen, J. H., 1971, Gum Hollow fan delta, Nueces Bay, Texas: The University of Texas at Austin, Bureau of Economic Geology Report of Investigations 69, 91 p.
- McGowen, J. H., Brown, L. F., Jr., Evans, T. J., Fisher, W. L., and Groat, C. G., 1976a, Environmental geologic atlas of the Texas Coastal Zone - Bay City-Freeport area: The University of Texas at Austin, Bureau of Economic Geology, 98 p., 9 maps.
- McGowen, J. H., Byrne, J. R., and Wilkinson, B. H., 1979, Geochemistry of bottom sediments, Matagorda Bay system, Texas: The University of Texas at Austin, Bureau of Economic Geology Geological Circular 79-2, 64 p.
- McGowen, J. H., Proctor, C. V., Jr., Brown, L. F., Jr., Evans, T. J., Fisher, W. L., and Groat, C. G., 1976b, Environmental geologic atlas of the Texas Coastal Zone--Port Lavaca area: The University of Texas at Austin, Bureau of Economic Geology, 107p.
- McGowen, J. H., and Brewton, J. L., 1975, Historical changes and related coastal processes, Gulf and mainland shorelines, Matagorda Bay area, Texas: The University of Texas at Austin, Bureau of Economic Geology Special Publication, 72 p., 16 maps.
- McGowen, J. H., and Morton, R. A., 1979, Sediment distribution, bathymetry, faults, and salt diapirs on the submerged lands of Texas: The University of Texas at Austin, Bureau of Economic Geology Special Publication, 31 p. 7 maps.
- McIntosh, R. P., 1967, An index of diversity and the relation of certain concepts to diversity: Ecology, v. 48, p. 392-404.
- McKinney, L. D., Bedinger, C. A., and Hopkins, S. H., 1976, The effects of shell dredging and siltation from dredging on organisms associated with oyster reefs, in Bouma, A. H., ed., Shell dredging and its influence on Gulf coast environments: Houston, Texas, Gulf Publishing Company, p. 281-303.
- McLusky, D. S., and Elliott, M., 1981, The feeding and survival strategies of estuarine molluscs, in Jones, N. V., and Wolff, W. J., eds., Feeding and survival strategies of estuarine organisms: New York, Plenum Press, p. 109-121.

- McMahan, C. A., 1968, Biomass and salinity tolerance of shoalgrass and manateegrass in Lower Laguna Madre, Texas: *Journal of Wildlife Management*, v. 32, no. 3, p. 501-506.
- McMillan, Calvin, 1984, The distribution of tropical seagrasses with relation to their tolerance of high temperatures: *Aquatic Botany*, v. 19, p. 369-379.
- McMillan, Calvin, and Moseley, F. N., 1967, Salinity tolerances of five marine spermatophytes of Redfish Bay, Texas: *Ecology*, v. 48, p. 503-506.
- Meade, R. H., 1969, Landward transport of bottom sediments in estuaries of the Atlantic coastal plain: *Journal of Sedimentary Petrology*, v. 39, no. 1, p. 222-234.
- Meade, R. H., and Emery, K. O., 1971, Sea level as affected by river runoff, eastern United States: *Science*, v. 173, p. 425-428.
- Meier, M. F., 1984, Contribution of small glaciers to global sea level: *Science*, v. 226, no. 4681, p. 1418-1421.
- Menge, B. A., and Sutherland, J. P., 1976, Species diversity gradients: synthesis of the roles of predation, competition, and temporal heterogeneity: *The American Naturalist*, vol. 110, no. 973, p. 351-369.
- Milliman, J. D., and Meade, R. H., 1983, World-wide delivery of river sediments to the oceans: *Journal of Geology*, v. 91, no. 1, p. 1-21.
- Minter, L. L., 1976, Impact of water resource development on the hydrology and sedimentology of the Brazos River system: Texas A&M University, College Station, Master's Thesis, 86 p.
- Mirabal, J., 1974, Suspended sediment load of Texas Streams: Compilation report, October 1965-September 1971: Texas Water Development Board Report 184, 119 p.
- Morgan, D. J., 1977, The Mississippi River Delta, legal-geomorphologic evaluation of historic shoreline changes: *Geoscience and Man*, v. 16, p. 196.
- Morgan, J. P., 1970a, Deltas--a resume: *Journal of Geological Education*, v. 18, no. 3, p. 107-117.
- Morgan, J. P., 1970b, Depositional processes and products in the deltaic environment, in Morgan, J. P., and Shover, R. H., eds., *Deltaic sedimentation, modern and ancient*: Society of Economic Paleontologists and Mineralogists, Special Publication 15, p. 31-47.
- Morgan, J. P., 1967, Ephemeral estuaries of the deltaic environment, in Lauff, G. H., ed., *Estuaries*: American Association for the Advancement of Science, p. 115-120.
- Morton, J. W., 1977, Ecological effects of dredging and dredge spoil disposal: a literature review: Washington, D.C., Department of the Interior, Fish and Wildlife Service, Technical paper, 33 p.

- Morton, R. A., 1972, Clay mineralogy of Holocene and Pleistocene sediments, Guadalupe delta of Texas: *Journal of Sedimentary Petrology*, v. 42, p. 85-88.
- Morton, R. A., 1977, Historical shoreline changes and their causes, Texas Gulf Coast: The University of Texas at Austin, Bureau of Economic Geology Geological Circular 77-6, p. 352-364.
- Morton, R. A., 1975, Shoreline changes between Sabine Pass and Bolivar Roads, an analysis of historical changes of the Texas Gulf shoreline: The University of Texas at Austin, Bureau of Economic Geology Geological Circular 75-6, 43 p.
- Morton, R. A., Pieper, M. J., and McGowen, J. H., 1976, Shoreline changes on Matagorda Peninsula (Brown Cedar Cut to Pass Cavallo), an analysis of historical changes of the Texas Gulf shoreline: The University of Texas at Austin, Bureau of Economic Geology Geological Circular 76-6, 37 p.
- Morton, R. A., and Donaldson, A. C., 1978, Hydrology, morphology, and sedimentology of the Guadalupe fluvial-deltaic system: *Geological Society of America Bulletin*, v. 89, p. 1030-1036.
- Morton, R. A., and McGowen, J. H., 1980, Modern Depositional Environments of the Texas Coast: The University of Texas at Austin, Bureau of Economic Geology, Guidebook 20, 167 p.
- Morton, R. A., and Paine, J. G., 1984, Historical shoreline changes in Corpus Christi, Oso, and Nueces Bays, Texas Gulf Coast: The University of Texas at Austin, Bureau of Economic Geology Geological Circular 84-6, 66 p.
- Morton, R. A., and Pieper, M. J., 1975, Shoreline changes in the vicinity of the Brazos River delta (San Luis Pass to Brown Cedar Cut), an analysis of historical changes of the Texas Gulf shoreline: The University of Texas at Austin, Bureau of Economic Geology, Geological Circular 75-4, 47 p.
- Morton, R. A., and Pieper, M. J., 1976, Shoreline changes on Matagorda Island and San Jose Island (Pass Cavallo to Aransas Pass), an analysis of historical changes of the Texas Gulf shoreline: The University of Texas at Austin, Bureau of Economic Geology Geological Circular 76-4, 42 p.
- Morton, R. A., and Pieper, M. J., 1977, Shoreline changes on Mustang Island and north Padre Island (Aransas Pass to Yarbrough Pass): The University of Texas at Austin, Bureau of Economic Geology Geological Circular 77-1, 45 p.
- Morton, R. A., and Price, W. A., 1987, Late Quaternary sea-level fluctuations and sedimentary phases of the Texas coastal plain and shelf: *Society of Economic Paleontologists and Mineralogists*, p. 181-198.
- Mullican, W. F., III, 1988, Subsidence and collapse at Texas salt domes: The University of Texas at Austin, Bureau of Economic Geology Geological Circular 88-2, 36 p.

- Nichols, F. H., 1977, Infaunal biomass and production on a mudflat, San Francisco Bay, California, in Coull, B. C., ed., Ecology of marine benthos: Columbia, South Carolina, University of South Carolina Press, p.339-357.
- Nichols, J., and Rowe, G., 1977, Infaunal macrobenthos off Cap Blanc, Spanish Sahara: Journal of Marine Research, v. 35, p. 525-536.
- Nichols, M. M., 1964, Characteristics of sedimentary environments in Moriches Bay: Papers in Marine Geology, Shepard commemorative volume, p. 363-383.
- Nichols, M. M., and Biggs, R. B., 1985, Estuaries, in Davis, R. A., Jr., ed., Coastal Sedimentary Environments: New York, Springer-Verlag, p. 77-186.
- Nichols, M. M., and Ellison, R. L., 1967, Sedimentary patterns of microfauna in a coastal plain estuary, in Lauff, G. H., ed., Estuaries: Washington, D.C., American Association for the Advancement of Science, Publication No. 83, p. 283-287.
- Nixon, S. W., 1980, Between coastal marshes and coastal waters -- a review of twenty years of speculation and research on the role of salt marshes in estuarine productivity and water chemistry, in Hamilton, Peter, and Macdonald, K. B., eds., Estuarine and wetland processes, p. 437-525.
- Nummedal, Dag, 1983, Barrier islands, in Komar, P. D., ed., CRC handbook of coastal processes and erosion: Boca Raton, Fla., CRC Press Inc., p. 77-121.
- O'Gower, A. K., and Wacasey, J. W., 1967, Animal communities associated with Thalassia, Diplanthera, and sand beds in Biscayne Bay I. analysis of communities in relation to water movements: Bulletin of Marine Science, v. 17, no. 1, p. 175-201.
- Odum, E. P., 1971, Fundamentals of ecology: Philadelphia, W. B. Saunders Company, 574 p.
- Odum, H. T., 1963, Productivity measurements in Texas turtle grass and the effects of dredging an intracoastal channel: Publications of the Institute of Marine Science, v. 9, p.48-58.
- Odum, H. T., and Hoskin, C. M., 1958, Comparative studies on the metabolism of marine waters: Publications of the Institute of Marine Science, v. 5, p. 16-46.
- Odum, H. T., and Wilson, R. F., 1962, Further studies on reaeration and metabolism of Texas Bays, 1958-1960: The University of Texas, Institute of Marine Science Publications, v. 8, p. 23-55.
- Oenema, Oene, and DeLaune, R. D., 1988, Accretion rates in salt marshes in the Eastern Scheldt, South-west Netherlands: Estuarine, Coastal and Shelf Science, v. 26, p. 379-394.

- Officer, C. B., Lynch, D. R., Setlock, G. H., and Helz, G. R., 1984, Recent sedimentation rates in Chesapeake Bay, in Kennedy, V. S., ed., *The estuary as a filter*: Orlando, Academic Press, p. 131-157.
- Orth, R. J., 1973, Benthic infauna of eelgrass, *Zostera marina*, beds: *Chesapeake Science*, v. 14, no. 4, p. 258-269.
- Orth, R. J., 1977, The importance of sediment stability in seagrass communities, in Coull, B. C., ed., *Ecology of marine benthos*: Columbia, South Carolina, University of South Carolina Press, p. 281-300.
- Orth, R. J., Heck, K. L., Jr., and van Montfrans, J., 1984, Faunal communities in seagrass beds: a review of the influence of plant structure and prey characteristics on predator-prey relationships: *Estuaries*, v. 7, n. 4A, p. 339-350.
- Paine, J. G., and Morton, R. A., 1986, Historical shoreline changes in Trinity, Galveston, West, and East Bays, Texas Gulf Coast: The University of Texas at Austin, Bureau of Economic Geology Geological Circular 86-3, 58 p.
- Parker, R. H., 1975, *The study of benthic communities, a model and a review*: Elsevier Scientific Publishing Company, New York, 279 p.
- Peet, R. K., 1974, The measurement of species diversity: *Annual Review of Ecology and Systematics*, v. 5, p. 285-307.
- Peres, J. M., and Picard, J., 1975, Causes of decrease and disappearance of the seagrass *Posidonia oceanica* on the French Mediterranean coast: *Aquatic Botany*, v. 1, p. 133-139.
- Persson, L., 1983, Temporal and spatial variation in coastal macrobenthic community structure, Hano Bay (southern Baltic): *Journal of Experimental Marine Biology and Ecology*, v. 68, p. 277-293.
- Pestrong, R., 1972, Tidal-flat sedimentation at Cooley Landing, southwest San Francisco Bay: *Sedimentary Geology*, v. 8, p. 251-288.
- Peterson, C. H., and Peterson, N. M., 1979, The ecology of intertidal flats of North Carolina: a community profile: Washington, D.C., U.S. Department of the Interior, Fish and Wildlife Service, Office of Biological Services, FWS/OBS-79/39, 73 p.
- Pethick, J. S., 1981, Long-term accretion rates on tidal marshes: *Journal of Sedimentary Petrology*, v. 51, no. 2, p. 571-577.
- Pickral, J. C., and Odum, W. E., 1976, Benthic detritus in salt marsh tidal creek, in Wiley, Martin, ed., *Estuarine Processes, Volume II*: Academic Press, New York, p. 280-292.
- Poff, M. J., 1973, Species composition, distribution and abundance of macrobenthic organisms in the intake and discharge areas after construction and operation of the Cedar Bayou Electric Power Station: Texas A&M University, Master's thesis, 300 p.

- Pratt, D. M., 1953, Abundance and growth of *Venus mercenaria* and *Callocardia morrhuana* in relation to the character of bottom sediments: *Journal of Marine Research*, v. 12, no. 1, p. 60-74.
- Pratt, W. E., and Johnson, D. W., 1926, Local subsidence of the Goose Creek oil field: *Journal of Geology*, v. 34, p. 577-590.
- Price, W. A., 1947, Equilibrium of form and forces in tidal basins of coast of Texas and Louisiana: *American Association of Petroleum Geologists*, v. 31, no. 3, p. 1616-1663.
- Price, W. A., 1956, Hurricanes affecting the coast of Texas from Galveston to the Rio Grande: U. S. Army Corps of Engineers, Beach Erosion Board Technical Memorandum 78, 17 p.
- Price, W. A., and Gunter, Gordon, 1943, Certain recent geological and biological changes in South Texas, with consideration of probable causes: *The Texas Academy of Science Proceedings and Transactions*, v. 26, p. 138-156.
- Pullen, E. J., and Trent, Lee, 1969, Hydrographic observations from the Galveston Bay system, Texas, 1958-1967: U. S. Fish and Wildlife Service data report 31, Washington, D.C., 151 p.
- Purdy, E. G., 1964, Sediments as substrates, in Imbrie, J., and Newell, N. D., eds., *Approaches to paleoecology*: New York, John Wiley, p.238-271.
- Ranwell, D. S., 1964, *Spartina* salt marshes in southern England. II. Rate and seasonal pattern in sediment accretion: *Journal of Ecology*, v. 52, p. 79-94.
- Ratzlaff, K. W., 1980, Land-surface subsidence in the Texas coastal region: U. S. Geological Survey Open-File Report 80-969, 19 p.
- Redfield, A. C., 1972, Development of a New England salt marsh: *Ecological Monographs*, v. 42, no. 2, p. 201-237.
- Rehkemper, L. J., 1969b, Sedimentology of Holocene estuarine deposits, Galveston Bay, Texas, in Lankford, R. R., and Rogers, J. J. W., compilers, *Holocene geology of the Galveston Bay area*: Houston Geological Society, p. 12-52.
- Rehkemper, L. J., 1969a, Sedimentology of Holocene estuarine deposits, Galveston Bay, Texas: Rice University, Houston, Texas, Ph.D. Dissertation, 68 p., 7 appendices.
- Reid, W. M., 1973, Active faults in Houston, Texas: The University of Texas at Austin, Ph. D. dissertation, 122 p.
- Reidenbaugh, T. G., Banta, W. C., Varricchio, M., Streiter, R. P., and Mendoza, S., 1983, Short-term accretional and erosional patterns in a Virginia salt marsh: *Gulf Research Reports*, vol. 7, no. 3, p. 211-215.

- Reinharz, Eli, and O'Connell, Arthur. 1983 Animal-sediment relationships of the upper and central Chesapeake Bay: Annapolis, Maryland, U.S. Environmental Protection Agency Final Report, EPA Grant No. R8059864, 49p.
- Renwick, W. H., and Ashley, G. M., 1984, Sources, storages, and sinks of fine-grained sediments in a fluvial-estuarine system: Geological Society of America Bulletin, v. 95, p. 1343-1348.
- Rhoads, D. C., 1974, Organism-sediment relations on the muddy sea floor: Oceanography and marine biology annual review, vol. 12, p. 263-300.
- Rhoads, D. C., McCall, P. L., and Yingst, J. Y., 1978, Disturbance and production on the estuarine seafloor: American Scientist, vol.66, p. 577-586.
- Rhoads, D. C., and Young, D. K., 1970, The influence of deposit-feeding organisms on sediment stability and community trophic structure: Journal of Marine Research, v.28, no.2, p.150-178.
- Rice Center for Community Design and Research, 1974, Environmental analysis for development planning, Chambers County, Texas, An approach to natural environmental analysis: Houston, Texas, Rice Center, Technical report, v. 1, 531 p.
- Rice, R. H., 1967, A study of the suspended load of the Trinity River, Texas: Rice University, Houston, Texas, Masters Thesis, 47 p.
- Rice, R. H., 1969, Suspended load of the Trinity River, chapter 4, in Lankford, R. R., and Rogers, John J. W., Holocene geology of the Galveston Bay area: Houston Geological Society, p. 78-84.
- Richard, G. A., 1978, Seasonal and environmental variation in sediment accretion in a Long Island salt marsh: Estuaries, v. 1, p. 29-35.
- Ritter, D. F., 1967, Rates of denudation: Journal of Geologic Education, v. 15, no. 4, p. 154-159.
- Rosen, P. S., 1977, Increasing shoreline erosion rates with decreasing tidal range in the Virginia Chesapeake Bay: Chesapeake Science, vol. 18, no. 4, p. 383-386.
- Rowe, G. T. and Smith, K. L., Jr., 1977, Benthic pelagic coupling in the mid-Atlantic bight, in Coull, B. C., ed., Ecology of marine benthos: Columbia, South Carolina, University of South Carolina Press, p.55-65.
- Rusnak, G. A., 1967, Rates of sediment accumulation in modern estuaries, in Lauff, G. H., ed., Estuaries: American Association for the Advancement of Science, Publication No. 83, p. 180-184.
- Rusnak, G. A., 1960, Sediments of Laguna Madre Texas, in Shepard, F. P., Phleger, F. B., and van Andel, T. H., eds., Recent sediments, northwest Gulf of Mexico: Tulsa, Oklahoma, American Association of Petroleum Geologists, p. 153-196.

- Russell, R. J., 1967, River and delta morphology: Louisiana State University, Coastal Studies Institute, Technical Report No. 52, 55p.
- Ryan, J. J., and Goodell, H. G., 1972, Marine geology and estuarine history of Mobile Bay, Alabama, Part 1. Contemporary sediments: Geological Society of America, Memoir 133, p. 517-554.
- Sanders, H. L., 1958, Benthic studies in Buzzards Bay. I. Animal - sediment relationships: *Limnology and Oceanography*, vol. 3, no. 3, p. 245-258.
- Sanders, H. L., 1968, Marine benthic diversity: a comparative study: *The American Naturalist*, vol. 102, no. 925, p. 1-40.
- Sanders, H. L., Goudsmit, E. M., Mills, E. L., and Hampson, G. E., 1962, A study of the intertidal fauna of Barnstable Harbor, Massachusetts: *Limnology and Oceanography*, vol. 7, no. 1, p.63-79.
- Santos, S. L., and Simon, J. L., 1974, Distribution and abundance of the polychaetous annelids in a south Florida estuary: *Bulletin of Marine Science*, vol. 24, no. 3, p. 669-689.
- Scaife, W. W., Turner, R. E., and Costanza, Robert, 1983, Coastal Louisiana recent land loss and canal impacts: *Environmental Management*, v. 7, no. 5, p. 433-442.
- Schaffner, L. C., Diaz, R. J., Olsen, C. R., and Larsen, I. L., 1987, Faunal characteristics and sediment accumulation processes in the James River estuary, Virginia: *Estuarine, Coastal and Shelf Science*, v. 25, p. 211-236.
- Schou, A., 1967, Estuarine research in the Danish moraine archipelago, in Lauff, G. H., ed., *Estuaries: American Association for the Advancement of Science Publication 83*, Washington, D.C., p. 129-145.
- Schubel, J. R., Carter, H. H., and Wise, W. M., 1979, Shrimping as a source of suspended sediment in Corpus Christi Bay (Texas): *Estuaries*, vol. 2, no. 3, p.201-203.
- Schubel, J. R., and Carter, H. H., 1984, The estuary as a filter for fine-grained suspended sediment, in Kennedy, V. S., ed., *The estuary as a filter: Academic Press*, New York, p. 81-105.
- Schubel, J. R., and Hirschberg, D. J., 1977, Pb210-determined sedimentation rate, and accumulation of metals in sediments at a station in Chesapeake Bay: *Chesapeake Science*, vol., 18, no. 4, p.379-382.
- Schultz, E. A., and Simmons, H. B., 1957, Fresh water-salt water density currents, a major cause of siltation in estuaries: U. S. Army Corps of Engineers, Technical bulletin no. 2, 28 p.
- Schumm, S. A., 1972, Fluvial paleochannels, in Rigby, J. K., and Hamblin, W. K., eds., *Recognition of ancient sedimentary environments: SEPM Special Publication No. 16*, p. 98-109.

- Schumm, S. A., 1968, Speculations concerning paleohydrologic controls of terrestrial sedimentation: Geological Society of America Bulletin, v. 79, p. 1573-1588.
- Scott, A. J., Hoover, R. A., and McGowen, J. H., 1969, Effects of Hurricane Beulah, 1967, on Texas coastal lagoons and barriers, in Castanares, A. A., and Phleger, F. B., eds., *Lagunas costeras, un simposio: Mexico, D. F., UNAM-UNESCO, Memoir Simposio International Lagunas Costeras*, Nov. 28-30, 1967, p. 221-236.
- Scrudato, R. J., Henningsen, B. F., and Estes, E. L., 1976, Trace metal concentrations of selected macrofauna from a southeast Texas estuary: The Texas Journal of Science, vol. 27, no. 4, p. 419-426.
- Scruton, P. C., 1960, Delta building and the deltaic sequence: in Shepard, F. P., Phleger, F. B., and van Andel, T. H., eds., *Recent sediments, northwest Gulf of Mexico: American Petroleum Institute and American Association of Petroleum Geologist*, Tulsa, p. 82-102.
- Seelig, W. N., and Sorensen, R. M., 1973, Historic shoreline changes in Texas: Texas A&M University Sea Grant, TAMU-SG-73-206, COE-165, 19p.
- Settlemyre, J. L., and Gardner, L. R., 1977, Suspended sediment flux through a salt marsh drainage basin: Estuarine and Coastal Marine Science, v. 5, p. 653-663.
- Shenton, E. H., 1957, A study of the foraminifera and sediments of Matagorda Bay, Texas: Gulf Coast Association of Geological Societies Transactions, v. 7, p. 135-147.
- Shepard, F. P., 1953, Sedimentation rates in Texas estuaries and lagoons: American Association of Petroleum Geology Bulletin, v. 37, no. 8, p. 1919-1934.
- Shepard, F. P., and Moore, D. G., 1960, Bays of central Texas coast, in Shepard, F. P., Phleger, F. B., and van Andel, T. H., eds., *Recent sediments, northwest Gulf of Mexico, Project 51, American Petroleum Institute: American Association of Petroleum Geologist*, Tulsa, Oklahoma, p. 117-152.
- Shepard, F. P., and Moore, D. G., 1955, Central Texas coast sedimentation: characteristics of environment, recent history, and diagenesis: American Association of Petroleum Geologists, Bulletin, v. 39, no. 8, p. 1463-1593.
- Sheridan, P. F., and Livingston, R. J., 1983, Abundance and seasonality of infauna and epifauna inhabiting a *Halodule wrightii* meadow in Apalachicola Bay, Florida: Estuaries, vol. 6, no. 4, p. 407-419.
- Sherk, J. A., Jr., 1971, The effects of suspended and deposited sediments on estuarine organisms: literature summary and research needs: Prepared for U.S. Army Corps of Engineers, Natural Resources Institute of the University of Maryland, Contribution no. 443, contract no. DACW73-70-C-0013, 73p.

- Shew, D. M., Baumann, R. H., Fritts, T. H., and Dunn, L. S., 1981, Texas barrier island region ecological characterization: environmental synthesis papers: Washington, D. C., U. S. Department of the Interior, Fish and Wildlife Service, Office of Biological Services, FWS/OBS-81/82, 413 p.
- Shideler, G. L., 1980, Reconnaissance observations of some factors influencing the turbidity structure of a restricted estuary: Corpus Christi Bay, Texas: Texas Journal of Science, vol. 32, no. 1, p. 59-71.
- Short, F. T., and Short, C. A., 1984, The seagrass filter: purification of estuarine and coastal waters, in Kennedy, V. S., ed., The estuary as a filter: New York, Academic Press, p. 395-413.
- Simons, D. B., and Simons, R. K., 1987, Differences between gravel- and sand-bed Rivers, in Thorne, C. R., Bathurst, J. C., and Hey, R. D., eds., Sediment transport in gravel-bed rivers: John Wiley Sons Ltd, New York, p. 3-11.
- Simpson, R. H., and Lawrence, M. B., 1971, Atlantic hurricane frequencies along the U. S. coastline: U. S. Department of Commerce, National Oceanic and Atmospheric Administration Technical Memorandum NWSSR-58, 14p.
- Smith, J. M., and Frey, R. W., 1985, Biodeposition by ribbed mussel, *Geukensia demissa*, in a salt marsh, Sapelo Island, Georgia: Journal of Sedimentary Petrology, vol. 55, no. 6, p. 817-828.
- Solomon, D. E., and Smith, G. D., 1973, Seasonal assessment of the relationship between the discharge of the Trinity River and the Trinity Bay ecosystem: Final report, U. S. Army Corps of Engineers, contract no. DACW 63-73-C-0059, prepared by Coastal Ecosystems Management, Inc., 147p.
- Sorenson, R. P., 1975, Clay mineralogy of suspended-load transport, Guadalupe River, Texas: The University of Texas at Austin, Masters Thesis, 66 p.
- Stevenson, J. C., Kearney, M. S., and Pendleton, E. C., 1985, Sedimentation and erosion in a Chesapeake Bay brackish marsh system: Marine Geology, v. 67, no. 3/4, p. 213-235.
- Stevenson, J. C., Ward, L. G., and Kearney, M. S., 1985, Vertical accretion in marshes with varying rates of sea level rise, in Wolfe, T. A., ed., Estuarine variability: Orlando, Academic Press, Inc., p. 241-259.
- Stevenson, J. C., Ward, L. G., and Kearney, M. S., 1988, Sediment transport and trapping in marsh systems: Implications of tidal flux studies: Marine Geology, v. 80, p. 37-59.
- Stevenson, J. C., Ward, L. G., and Kearney, M. S., 1986, Vertical accretion in marshes with varying rates of sea level rise, in Wolfe, D. A., ed., Estuarine variability: New York, Academic Press, p. 241-259.
- Stone, A. N., and Reish, D. J., 1965, The effect of fresh-water run-off on a population of extuarine polychaetous annelids: Bulletin of the Southern California Academy of Sciences, vol. 64, part 3, p. 111-119.

- Stoner, A. W., 1980, The role of seagrass biomass in the organization of benthic macrofaunal assemblages: *Bulletin of Marine Science*, v. 30, no.3, p. 537-551.
- Stoner, A. W., Greening, H. S., Ryan, J. D., and Livingston, R. J., 1983, Comparison of macrobenthos collected with cores and suction sampler in vegetated and unvegetated marine habitats: *Estuaries*, vol. 6, no. 1, p. 76-82.
- Stora, G., and Arnoux, A., 1983, Effects of large freshwater diversions on benthos of a Mediterranean lagoon: *Estuaries*, vol.6, no.2, p. 115-125.
- Stout, I. M., Bentz, L. C., and Ingram, H. W., 1961, Silt load of Texas streams: a compilation report, June 1889 --September 1959: Texas Board of Water Engineers, Bulletin 6108, 236 p.
- Strand, R. I., and Pemberton, E. L., 1982, Reservoir sedimentation: Technical guideline for Bureau of Reclamation: U. S. Department of the Interior, Bureau of Reclamation, Sedimentation and River Hydraulics Section, Engineering and Research Center, Denver, Colorado, 48 p.
- Stumpf, R. P., 1983, The process of sedimentation on the surface of a salt marsh: *Estuarine, Coastal and Shelf Science*, v. 17, no. 5, p. 495-508.
- Subrahmanyam, C. B., Kruczynski, W. L., and Drake S. H., 1976, Studies on the animal communities in two north Florida salt marshes: part II. macroinvertebrate communities: *Bulletin of Marine Science*, vol. 26, no. 2, p. 172-195.
- Suhayda, Joseph N., 1987, Sea level and subsidence: MMS Information Transfer Meeting, November, 1986, p. 48.
- Summerhayes, C. P., Ellis, J. P., and Stoffers, P., 1985, Estuaries as sinks for sediment and industrial waste--a case history from the Massachusetts coast: *Contributions to Sedimentology*, 14, Stuttgart, E. Schweizerbart'sche Berglugsbuchhandlung (Nagele u. Obermiller), 47 p.
- Swanson, R. L., and Thurlow, C. I., 1973, Recent subsidence rates along the Texas and Louisiana coasts as determined from tide measurements: *Journal of Geophysical Research*, v. 78, no. 5, p. 2665-2671.
- Tacconi, P., and Billi, P., 1987, Bed load transport measurements by the vortex-tube trap on Virginio Creek, Italy, in Thorne, C. R., Bathurst, J. C., and Hey, R. D., eds., *Sediment transport in gravel-bed rivers*: John Wiley & Sons Ltd, New York, p. 583-606.
- Tenore, K. R., 1977, Food chain pathways in detrital feeding benthic communities: a review, with new observations on sediment resuspension and detrital recycling, in Coull, B. C., ed., *Ecology of marine benthos*: Columbia, South Carolina, University of South Carolina Press, p. 37-53.
- Tenore, K. R., 1972, Macrobenthos of the Pamlico River estuary, North Carolina: *Ecological Monographs*, vol. 42, no. 1, p. 51-69.

- Texas Department of Water Resources, 1978b, Hydrological and biological studies of the Colorado River delta, Texas: Texas Department of Water Resources, LP-79, 73p.
- Texas Department of Water Resources, 1978a, Hydrological and biological studies of the Trinity River Delta, Texas: Texas Department of Water Resources, LP-70, 176p.
- Texas Department of Water Resources, 1980a, Lavaca-Tres Palacios estuary: a study of the influence of freshwater inflows: Texas Department of Water Resources, LP-106, 292p.
- Texas Department of Water Resources, 1981a, Nueces and Mission-Aransas estuaries: a study of the influence of freshwater inflows: Texas Department of Water Resources, LP-108, 308p.
- Texas Department of Water Resources, 1981c, Sabine-Neches estuary: a study of the influence of freshwater inflows: Texas Department of Water Resources: LP-116, 292p.
- Texas Department of Water Resources, 1980b, Guadalupe estuary: a study of the influence of freshwater inflows: Texas Department of Water Resources, LP-107, 277p.
- Texas Department of Water Resources, 1983, Laguna Madre estuary: a study of the influence of freshwater inflows: Texas Department of Water Resources, LP-182, 250p.
- Texas Department of Water Resources, 1982, The influence of freshwater inflows upon the major bays and estuaries of the Texas Gulf coast: Texas Department of Water Resources, Executive Summary, LP-115, 53 p.
- Texas Department of Water Resources, 1981b, Trinity-San Jacinto estuary: a study of the influence of freshwater inflows: Texas Department of Water Resources, LP-113, 310p.
- Texas Department of Water Resources, Division of Planning and Development, 1978, Present and future surface-water availability in the Colorado River basin, Texas: Texas Department of Water Resources, LP-60, 160p., 5 appendices.
- Texas Water Development Board 1973, Engineering data on dams and reservoirs in Texas, part II: Texas Water Development Board Report 126, unpaginated.
- Thorbjarnarson, K. W., Nittrouer, C. A., DeMaster, D. J., and McKinney, R. B., 1985, Sediment accumulation in a back-barrier lagoon, Great Sound, New Jersey: *Journal of Sedimentary Petrology*, v. 55, no. 6, p. 856-863.
- Thornthwaite, C. W., 1948, An approach toward a rational classification of climate: *Geographical Review*, v. 38, no. 1, p. 55-94.
- Thorson, G., 1957, Bottom communities (sublittoral or shallow shelf): *Geological Society of America, Memoir 67*, vol. 1, p. 461-534.

- Thurman, C. L., 1984, Ecological notes on fiddler crabs of South Texas, with special reference to *Uca subcylindrica*: *Journal of Crustacean Biology*, v. 4, no. 4, p. p. 665-681.
- Trent, L., Pullen, E., and Proctor, R., 1976, Abundance of macrocrustaceans in a natural marsh and a marsh altered by dredging, bulkheading, and filling: *Fishery Bulletin*, vol. 74, no. 1, p.195-200.
- Turner, R. E., 1977, Intertidal vegetation and commercial yields of penaeid shrimp: *Transactions of the American Fisheries Society*, vol. 106, no. 5, p. 411-416.
- U. S. Army Corps of Engineers, 1967, Interim report on Colorado River and tributaries, Texas, covering mouth of Colorado River: U. S. Army Engineer District, Galveston, Texas, 29 p., 6 plates, 4 appendices, attachments.
- U. S. Army Corps of Engineers, 1981, Mouth of Colorado River, Texas phase I: general design memorandum and environmental impact statement (diversion features): Galveston, Texas, U. S. Army Engineer district, 123 p.
- U. S. Army Corps of Engineers, 1977, Mouth of Colorado River, Texas, phase 1: general design memorandum (navigation features): Galveston, Texas, U. S. Army Engineer District, 292 p.
- U. S. Army Corps of Engineers, 1960, Report on second resurvey of sedimentation, Denison Dam and Reservoir (Lake Texoma), Red River, Texas and Oklahoma: Tulsa, Oklahoma, U. S. Army Engineer District, 16p.
- U. S. Army Corps of Engineers, 1956, Wave statistics for the Gulf of Mexico off Caplan, Texas: Beach Erosion Control Board, Technical Memorandum No. 86, 51 p.
- U. S. Army Corps of Engineers, undated Trinity River Project, Texas, general design memorandum, phase 1, Appendix 3, planning background, vol. III of VI.
- U. S. Department of Agriculture, 1959, Inventory and use of sedimentation data in Texas: Soil Conservation Service, U. S. Department of Agriculture, Bull. 5912, 85p.
- U. S. Department of Commerce, 1980, Climatological data, national summary, annual summary: National Oceanic and Atmospheric Administration, Environmental Data and Information Service, Asheville, N. C., v. 31, no. 13, 106 p.
- U. S. Department of Commerce, 1980, Tide current tables 1981, Atlantic coast of North America: National Oceanic and Atmospheric Administration, National Ocean Survey, 231 p.
- U. S. Department of Commerce, 1983, Tide current tables 1984, Atlantic coast of North America: National Oceanic and Atmospheric Administration, National Ocean Survey, 250 p.

- U.S. Department of Commerce, 1978, Tide current tables 1979, Atlantic coast of North America: National Oceanic and Atmospheric Administration, National Ocean Survey, 231 p.
- United States Department of Agriculture, 1972, Texas Coastal Basins: Interim report, Type IV Cooperative River Basin Survey, 262p.
- Van Beek, J. L., Gael, B. T., Harris, M. K., Irish, G. J., McLindon, C. D., Dugan, S. S., and Wicker, K. M., 1980, Delta building potential of the Colorado River Central Texas, in U. S. Army Corps of Engineers, 1981, Mouth of Colorado River, Texas phase I. General design memorandum and environmental impact statement (diversion features): Galveston, Texas, U. S. Army Corps of Engineers, 209 p.
- Van Beek, J. L., and Meyer-Arendt, K., 1981, Sediment--asset or liability, in Cross, R. D., and Williams, D. L., eds., Proceedings of the National Symposium on freshwater inflow to estuaries, volume II: U. S. Department of Interior, Fish and Wildlife Service, Slidell, Louisiana, FWS/OBS 81/04, p. 197-215.
- Van Heerden, I. Li., Wells, J. T., and Roberts, H. H., 1981, Evolution and morphology of sedimentary environments, Atchafalaya Delta, Louisiana: Gulf Coast Association of Geological Societies, Transactions, v. 31, p. 399-408.
- Van Heerden, I. Li., and Roberts, H. H., 1980, The Atchafalaya Delta--Louisiana's new prograding coast: Gulf Coast Association of Geological Societies, Transactions, v. 30, p. 497-506.
- Van Siclen, D., 1967, The Houston fault problem: Institute of Professional Geologists, Texas Section 3rd Annual Meeting, Proceedings, p. 9-31.
- Van Straatan, L. M. J. U., and Kuenen, P. H., 1958, Tidal action as a cause of clay accumulation: Journal of Sedimentary Petrology, v. 28, p. 406-413.
- Vanoni, V. A., 1975, Sedimentation engineering: American Society of Civil Engineers, New York, ASCE--Manuals and reports on engineering practice--No. 54, 745 p.
- Verbeek, E. R., and Clanton, U. S., 1981, Historically active faults in the Houston metropolitan area, Texas, in Etter, E. M. ed., Houston area environmental geology: surface faulting, ground subsidence, hazard liability: Houston Geological Society, p. 28-68.
- Virnstien, R. W., 1977, The importance of predation by crabs and fishes on benthic infauna in Chesapeake Bay: Ecology, v. 58, p. 1199-1217.
- Virnstien, R. W., Mikkelsen, P. S., Cairns, K. D., and Capone, M. A., 1983, Seagrass beds versus sand bottoms: the trophic importance of their associated benthic invertebrates: Florida Scientist, vol. 46, no. 3/4, p. 363-381.
- Wadsworth, A. H., Jr., 1966, Historical deltation of the Colorado River, Texas, in Shirley, M. L., ed., Deltas in their geologic framework: Houston Geological Society, p. 99-105.

- Wadsworth, A. H., Jr., 1941, The lower Colorado River, Texas: The University of Texas at Austin, Master's thesis, 61 p.
- Ward, G. H., Armstrong, N. E., and the Matagorda Bay Project Teams, 1980, Matagorda Bay Texas: its hydrography, ecology and fishery resources: U.S. Fish and Wildlife Service, Biological Services Program, Washington D.C., FWS/OBS-81/52, 230 p.
- Ward, L. G., Kemp, W. M., and Boynton, W. R., 1984, The influence of waves and seagrass communities on suspended particulates in an estuarine embayment: Marine Geology, vol. 59, p. 85-103.
- Warwick, R., and Price, R., 1975, Macrofauna production in an estuarine mud-flat: Journal Marine Biological Association of the United Kingdom, vol. 55, no. 1, p. 1-18.
- Watson, R. L., and Behrens, E. W., 1976, Hydraulics and dynamics of New Corpus Christi Pass, Texas: a case history, 1973-75: U. S. Army Coastal Engineering Research Center, GITI Report 9, 175 p.
- Wayne, C. J., 1974, Effect of artificial sea-grass on wave energy and near-shore sand transport: Gulf Coast Association of Geological Societies, Transactions, v. 24, p. 279-282.
- Weaver, P., and Sheets, M., 1962, Active faults, subsidence and foundation problems in the Houston, Texas, area, in Geology of the Gulf Coast and central Texas: Houston Geological Society Guidebook, p. 254-265.
- Welborn, C. T., 1967, Comparative results of sediment sampling with the Texas sampler and the depth-integrating samplers and specific weight of fluvial sediment deposits in Texas: Texas Water Development Board Report, 36, 109 p.
- Welborn, C. T., and Andrews, F. L., 1980, Sediment and water discharge of the Colorado River at Matagorda, Texas May 4-7, and June 7-11, 1979: U. S. Department of the Interior, Geological Survey, Open file report 80-1177, 21p.
- White, W. A., 1985, Textural and geochemical relationships in benthic sediments in three bay-estuary-lagoon systems in Texas: Offshore Technology Conference, Houston, May 6-9, 1985, OTC 4905, p. 67-74.
- White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., Littleton, T. G., McGowen, J. H., and Nance, H. S., 1987, Submerged lands of Texas, Beaumont-Port Arthur area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands: The University of Texas at Austin, Bureau of Economic Geology Special Publication, 110 p.
- White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., Littleton, T. G., McGowen, J. H., Nance, H. S., and Schmedes, K. E., 1986, Submerged lands of Texas, Brownsville-Harlingen area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands: The University of Texas at Austin, Bureau of Economic Geology Special Publication, 138 p.

- White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., Littleton, T. G., McGowen, J. H., Nance, H. S., and Schmedes, K. S., 1983, Submerged lands of Texas, Corpus Christi area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands: The University of Texas at Austin, Bureau of Economic Geology Special Publication, 154 p.
- White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., Littleton, T. G., McGowen, J. H., Nance, H. S., and Schmedes, K. E., 1985, Submerged lands of Texas, Galveston-Houston area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands: The University of Texas at Austin, Bureau of Economic Geology Special Publication, 145 p., 6 plates.
- White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., Littleton, T. G., McGowen, J. H., and Nance, H. S., in preparation, Submerged lands of Texas: Kingsville area: sediment, geochemistry, benthic macroinvertebrates, and associated wetlands: The University of Texas at Austin, Bureau of Economic Geology.
- White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., Littleton, T. G., McGowen, J. H., and Nance, H. S., in press a, Submerged lands of Texas, Bay City-Freeport area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands: The University of Texas at Austin, Bureau of Economic Geology.
- White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., Littleton, T. G., McGowen, J. H., and Nance, H. S., in press b, Submerged Lands of Texas, Port Lavaca area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands: The University of Texas at Austin, Bureau of Economic Geology.
- White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., and Littleton, T. G., 1984, State submerged lands of Texas--sediments, geochemistry, benthic macroinvertebrates, and associated wetlands, in Magoon, O. T., and others, Coastal zone '83: 3rd Symposium on Coastal and Ocean Management, post-conference volume, p. 1-17.
- White, W. A., and Calnan, T. R., in preparation, Marsh sedimentation Colorado and Trinity River deltas Texas Gulf Coast: field studies: The University of Texas at Austin, Bureau of Economic Geology, Draft report prepared for Texas Parks and Wildlife Department.
- White, W. A., and Galloway, W. E., 1977, Guide to modern barrier environments of Mustang and north Padre Islands and Jackson (Eocene) barrier/lagoon facies of the South Texas uranium district: The University of Texas at Austin, Bureau of Economic Geology Research Note 7, 51p.
- White, W. A., and Morton, R. A., 1987, Historical shoreline changes in San Antonio, Espiritu Santo, and Mesquite Bays, Texas Gulf Coast: The University of Texas at Austin, Bureau of Economic Geology Geological Circular 87-1, 41 p.
- Whiting, N. H., and Moshiri, G. A., 1974, Certain organism-substrate relationships affecting the distribution of *Uca minax* (Crustacea: Decapoda): *Hydrobiologia*, vol. 44, p. 481-493.

- Whitlatch, R. B., 1977, Seasonal changes in the community structure of the macrobenthos inhabiting the intertidal sand and mud flats of Barnstable Harbor, Massachusetts: *Biological Bulletin*, vol. 152, p. 275-294.
- Wiersema, J. M., Mitchell, R. P., and James, S. N., 1973, Sabine Power Station ecological program, v. 1, A. utilization of aquatic habitats in the GSU marsh, B. assessment of GSU thermal discharge on the biota of the Neches River: Austin, Texas, Tracor, Inc., document no. T73-AU-9507-U(R), submitted to Gulf States Utilities, 215 p.
- Wilkinson, B. H., and Basse, R. A., 1978, Late Holocene history of the central Texas coast from Galveston Island to Pass Cavallo: *Geological Society of America, Bulletin*, v. 89, no. 10, p. 1592-1600.
- Wilkinson, B. H., and Byrne, J. R., 1977, Lavaca Bay transgressive deltaic sedimentation in central Texas USA estuary, *American Association of Petroleum Geologists Bulletin*, v. 61, no. 4, p. 527-545.
- Williams, G. P., and Wolman, G. M., 1984, Downstream effects of dams on alluvial rivers: *U. S. Geological Survey Professional Paper* 1286, 64 p.
- Williams, R. B., and Thomas, L. K., 1967, The standing crop of benthic animals in a North Carolina estuarine area: *Journal of the Elisha Mitchell Scientific Society*, vol. 83, p. 135-139.
- Windom, H. L., Neal, W. J., and Beck, K. C., 1971, Mineralogy of sediments in three Georgia estuaries: *Journal of Sedimentary Petrology*, v. 41, p. 497-504.
- Winker, C. D., 1979, Late Pleistocene fluvial-deltaic deposition, Texas coastal plain and shelf: *The University of Texas at Austin, Master's thesis*, 187 p.
- Winslow, A. G., and Doyel, W. W., 1954, Land-surface subsidence and its relation to the withdrawal of ground water in the Houston-Galveston region, Texas: *Economic Geology*, v. 49, no. 4, P. 413-422.
- Winston, J. E., and Anderson, F. E., 1971, Bioturbation of sediments in a northern temperate estuary: *Marine Geology*, v.10, p. 39-49.
- Wohlschlag, D. E., and Copeland, B. J., 1970, Fragile estuarine systems--ecological considerations: *Water resources bulletin*, vol. 6, p. 94-105.
- Wolff, W. J., and de Wolf, L., 1977, Biomass and production of zoobenthos in the Grevelingen estuary, the Netherlands: *Estuarine and Coastal Marine Science*, vol. 5, p. 1-24.
- Woodward-Clyde Consultants, 1977, Sediment chemistry and benthic biological communities in Matagorda Bay, Texas: *San Fransico, California*, 60p.
- Woolley, B. W., 1985, Channel changes on the Brazos River below Whitney Dam, Bosque, Hill, and McLennan Counties, Texas: *Baylor University, Master's thesis*, 61 p.

Wright, L. D., 1985, River deltas, in Davis, R. A., Jr., Coastal Sedimentary Environments, 2nd revised, expanded edition:

Wright, L. D., 1977, Sediment transport and deposition at river mouths; a synthesis: Geological Society of America Bulletin, v. 88, no. 6, p. 857-868.

Wright, S. S., 1980, Seismic stratigraphy and depositional history of Holocene sediments on the central Texas Gulf coast: The University of Texas at Austin, Master's thesis, 123 p.

Young, D. K., and Rhoads, D. C., 1971, Animal-sediment relations in Cape Cod Bay, Massachusetts I. A transect study: Marine Biology, v. 11, p. 242-254.

Zetler, B. D., 1953, Some effects of the diversion of the Santee River on the waters of Charleston Harbor: Transactions, American Geophysical Union, vol.34, no.5, p. 729-732.

Zimmerman, R. J., Minello, T. J., and Zamora, G., Jr., 1984, Selection of vegetated habitat by brown shrimp, *Penaeus aztecus*, in a Galveston Bay salt marsh: Fishery Bulletin, vol. 82, no. 2, p.325-336.

Zimmerman, R. J., and Minello, T. J., 1984, Densities of *Penaeus aztecus*, *Penaeus setiferus*, and other natant macrofauna in a Texas salt marsh: Estuaries, vol. 7, no. 4A, p. 421-433.

APPENDIX

FLUVIAL-ESTUARINE SEDIMENTATION AND FLORAL/FAUNAL RELATIONSHIPS ANNOTATED BIBLIOGRAPHY OF SELECTED REFERENCES¹

Aksoy, S., 1967, Riverbed degradation downstream of large dams: Sedimentation, annotated bibliography of foreign literature, 1967-1968, Survey No: 5, U. S. Department of Agriculture and National Science Foundation, Washington D.C., # 3787, p. 5.

DAMS, RIVERS, EROSION, DEPOSITION

This original article is in Turkish with an English abstract. Construction of dams effects existing regime -- all transported solids will be trapped; released clear water downstream is more uniform in distribution. The riverbed will erode until new equilibrium is reached. Presented in the paper are results of a theoretical study of the rate of channel deepening, and an experimental investigation in a laboratory flume. Also, an equation that can be used for calculating the degradation. (This information is not presented in the annotated bibliography).

Anderson, R. Y., 1977, Short term sedimentation response in lakes in western United States as measured by automated sampling: Limnology and Oceanography, v. 22, p. 423-433.

SEDIMENTATION, SEDIMENT TRAP, LAKES

Describes sediment traps that amplify rate of sedimentation and automatically time depositional process. The trap uses a steep-sided cone (funnel) to amplify the rate of sedimentation and an automatic device that releases a marking layer into a collecting tube to set off, or sample, equal increments of time in an accumulating column of collected material.

Armstrong, N. E., Harris, S. E., and Gordon, V. N., 1977, Exchange rates for carbon, nitrogen, and phosphorus in the Trinity River delta marshes: The University of Texas at Austin, Center for Research in Water Resources, CRWR-154, EHE 77-07, 63 p.

TRINITY RIVER, DELTA, MARSHES, CARBON, NUTRIENTS

Report has some data on total suspended solids (table IV-9) and organic vs. inorganic sediments at some sites in marshes.

Armstrong, N. E., and Ward, G. H., 1981, Effects of alterations of freshwater inflows into Matagorda Bay, Texas, in Cross, R. D., and Williams, D. L., Proceedings of the National Symposium on Freshwater Inflow to Estuaries, vol. 2, p. 179-196.

¹ An annotated bibliography was not required under the IAC but a partial one was completed early in the project to assist in report preparation. Many more references were used in the report than are cited here.

DIVERSION, OYSTERS, SEDIMENTATION, SILTATION, GILLS, MATAGORDA BAY, PREDATOR, PARASITE, SALINITY

This is an excerpt from only a portion of the biological impact from the diversion--"The freshwater diversion will have mixed effects on oysters, in that while it will increase the flow of fresh water and may increase nutrient influx into Matagorda Bay, it will also increase sedimentation. The results will be, on one hand, lowered salinity and reduced predator and parasite impacts but, on the other hand, increased siltation and turbidity, hence clogging of gill structures of the oysters. The most serious impacts will be due to sedimentation."

Bader, R. G., 1954, The role of organic matter in determining the distribution of pelecypods in marine sediments: *Journal of Marine Research*, v.13, p. 32-47.

PELECYPODS, ORGANIC CONTENT, SEDIMENT, DECOMPOSITION, REFRACTORY, NONREFRACTORY, OXYGEN, LIGNIN, DENSITY, DEPTH, SALINITY, TEMPERATURE, SILT, CLAY, CURVILINEAR, RESINS, CELLULOSE, MT. DESERT ISLAND, MAINE

Twenty-seven samples were taken in the bays and sounds around Mt. Desert Island, Maine. The most important variables controlling pelecypod density are organic content and its state of decomposition. Density decreases as decomposition increases. Beyond 3% organic content, products of decomposition and decline in available oxygen become limiting variables. Maximum populations are present in sediments containing 1.18 to 3.00% organics. "If the type of organics are known, then it is possible to determine the state of decomposition." Lipids, resins, celluloses, etc. are nonrefractory and decompose rapidly, while lignin is refractory and decomposes slowly. The ratio of the refractory to nonrefractory times the total amount of organic matter that is available equals the coefficient of decomposition. From Bader's samples, those with decomposition coefficient of less than one had more than 48 pelecypod individuals. Thus there is a curvilinear relationship between amount of organics and density--as organics increase so do individuals until coefficient of decomposition is high and population decreases.

Bartberger, C. E., 1976, Sediment sources and sedimentation rates, Chincoteague Bay, Maryland and Virginia: *Journal of Sedimentary Petrology*, v. 46, no. 2, p. 326-336.

SEDIMENTATION RATES, BAY, MARYLAND, VIRGINIA

Sediment load from stream discharging into the bay is predominantly mud. Assuming a bulk density of 2.0 g/cu cm for sediment accumulation in the bay, annual sediment yield from the 270 sq km watershed is about 5,000 cu m. Sediment contributed from shoreline erosion is estimated at 40,000 cu m, which is also principally mud. Total mud/yr = 45,000 cu m. Based on sand-to-mud ratio which is 1:1 as determined by grain size analyses, an equal volume of sand must also have been contributed to the system (45,000 cu m). The principal immediate source of the sand appears to be the barrier, Assateague Island, with storm washover being twice as effective as eolian transport.

However, this source today appears much too low. Sand transported through two active inlets is important only in the immediate vicinity of the inlets. Bay sedimentation rate based on volume of 90,000 cu m/yr is only 0.3 mm/yr over the 310 sq km area of the bay. Over past 5000 yrs an estimate of the actual rate based on borehole data is 1.5 mm/yr. Difference maybe due to past active tidal inlets which are now closed, resulting in present low sed rate and marsh decline

Bascescu, M. C., 1972, Substratum - Animals, in Kinne, O., ed., Marine ecology, v. 1, p. 1291-1322.

SUBSTRATE, MUD, SAND, BENTHOS, STABILITY, DAMS, COMMUNITIES

Textbook discussion of influence of substrate on animals. Discussions on structural responses to substrate, distribution and substrate, larval and reproductive response to various substrates, and functional responses to substrates, among others. Mentions example of a dam on Rance River in France and its influence on benthos in estuary of river. Influence included water stagnation, a slight reduction in salinity and silting of sandy areas causing death of several populations and their replacement by more calcareous forms.

Baumann, R. H., Day, J. W., Jr., and Miller, C. A., 1984, Mississippi deltaic wetland survival: sedimentation versus coastal submergence: Science, v. 224, no. 4653, p. 1093-1095.

WETLANDS, DELTA, MARSH, SEDIMENTATION RATES

Report documents the differences in the mechanisms of sedimentation at two wetland locations in the Mississippi deltaic plain. Success of marshes to combat the effects of apparent sea-level rise and erosion is dependent on riverine input because basinal sedimentary processes are insufficient to combat the effects. Deteriorating marshes are accreting at a faster rate (15 mm/yr at streamside, 9 mm/yr at inland areas) than stable marshes (13 mm/yr at streamside, 6 mm/yr at inland areas), but the deteriorating marshes are not maintaining their intertidal elevation as well as the stable marshes relative to local apparent sea-level rise as measured by local tide gauges. The mean annual sedimentation rates are in accord with rates of 137 Cs for saline marshes in the area (Delaune, Raddy, Patrick, 1978). Hurricanes and winter storms are major depositional and erosional agents.

Bell, S. B., 1980, Meiofauna-macrofauna interactions in a high salt marsh habitat: Ecological Monographs, v. 50, no. 4 p. 487-505.

MEIOFAUNA, MACROFAUNA, SALT MARSH, PREDATION

From abstract-"Biological interactions between meiofauna and macroepifauna were studied in a high salt marsh habitat using manipulative techniques. When macroepifauna were excluded from experimental cage sites, meiofauna densities increased significantly compared to open, noncaged areas...My experimental results support the hypothesis that macroepifauna predation/disturbance has an important effect on meiofaunal assemblages, thus providing insight into possible competitive associations between meiofauna taxa."

Beschta, R. L., 1987, Conceptual models of sediment transport in streams, in Thorne, C. R., Bathurst, J. C., and Hey, R. D., Sediment transport in gravel-bed rivers: John Wiley & Sons Ltd, New York, p. 387-408.

SEDIMENT TRANSPORT, STREAMS, BEDLOAD, SUSPENDED LOAD,

Factors influencing the movement of sediments in streams include variable source areas of sediments, transient flows, variable particle sizes, non-uniformity of channel geometry and flow, and dynamic/adjusting channels. Suspended load and bed load transport becomes indistinct for particles ranging in diameter from about 0.1 to 1 mm. Bed load transport has high temporal and spatial variability making accurate prediction of transport difficult using hydraulically-based formulae.

Biggs, R. B., and Howell, B. A., 1984, The estuary as a sediment trap: alternate approaches to estimating its filtering efficiency, in Kennedy, V. S., ed., The estuary as a filter: New York, Academic Press, p. 107-129.

ESTUARY, SEDIMENT, SEDIMENTATION, BENTHOS, DEPOSITION

"The purpose of this section is to evaluate the hypothesis that biologically mediated sedimentation processes are as important, if not more so, than the mechanical or physical processes which lead to the deposition of fine sediments."

Biggs, Robert, B., 1967, The sediments of Chesapeake Bay, in Lauff, G. H., ed., Estuaries, Washington, D.C., American Association for the Advancement of Science, Publication No. 83, p. 239-260.

SEDIMENTS, CHESAPEAKE BAY, ESTUARIES, ORGANIC CARBON,

One of the findings was an increase of organic carbon in surface sediments with increasing water depth which may be attributed to several factors including higher rate of sedimentation in shallow water diluting organic carbon, higher rate of inorganic oxidation in shallow waters due to higher oxygen in water and circulation in coarser sediment, scavenging by organisms in shallow water, and higher physical energy.

Bloom, S. A., Simon, J. L., and Hunter, V. D., 1972, Animal-sediment relations and community analysis of a Florida estuary: Marine Biology, v. 13, p. 43-56.

TAMPA BAY, FLORIDA, SEDIMENT, COMMUNITY, DEPOSIT FEEDER, FILTER FEEDER, GRAIN-SIZE, TROPHIC-GROUP AMENSALISM

This study supports Sanders--optimal median grain-size for filter feeders was 0.174mm (Sanders-0.18mm) and that the assemblage most dominated by deposit feeders has a high variance in the silt-clay fraction (1.3-22.09%). This study also supports Rhoads and Young (1970) in that suspension feeders are largely confined to sandy or firm mud-bottoms, while deposit feeders attain

high densities on soft, muddy substrates. In this study, filter feeders were positively correlated to sorting and positively correlated to skewness. "Based on the demonstration of an ecotone community and the lack of a consistent multi-species dominance pattern, a view of communities as abstractions from continua of distributions was supported over a Peterson-type community concept."

Boesch, D. F., 1973, Classification and community structure of macrobenthos in the Hampton Roads area, Virginia: Marine Biology, v. 21, p. 226-244.

MACROBENTHOS, COMMUNITY, CLASSIFICATION, HAMPTON ROADS, VIRGINIA, BENTHOS, SEDIMENT, SPECIES DIVERSITY, SAND, MUD, SPECIES RICHNESS, TEMPORAL, ELIZABETH RIVER, SUBSTRATE, SUSPENSION-FEEDER, EPIFAUNA, INFAUNA

"Within Hampton Roads proper, the overwhelming factors responsible for the observed spatial patterns of species distribution and diversity were those pertaining to the substrate." Most species in Hampton Roads were either not bottom-type specific or restricted in varying degrees to sand bottoms. Species that were specific to particular sediment were distributed according to silt and clay content in sediment or the availability of hard substrates in the form of shell or gravel. Suspension feeders were excluded from mud bottoms. Species richness was highest in sands due to the large number of epifauna and differences in deposit-feeding infaunal species richness. Epifauna of sands more diverse because more suitable hard substrates occur in sands and substrates will not be silted over. "Greater spatial heterogeneity allowed greater species diversity on stable sand bottoms than on mud." Most of the numerically dominant species were ubiquitous or markedly seasonal.

Boesch, D. F., 1972, Species diversity of marine macrobenthos in the Virginia area: Chesapeake Science, v. 13, no.3, p. 206-211.

DIVERSITY, MACROBENTHOS, SHANNON'S FORMULA, SEDIMENT, STABILITY, POLYHALINE, MESOHALINE, OLIGOHALINE, SPECIES RICHNESS, EQUITABILITY, SALINITY, MUD, SAND, VIRGINIA, HAMPTON ROADS, JAMES RIVER, CHESAPEAKE BAY

Five areas with different salinity regimes were sampled:(1)OCS off Virginia and North Carolina (2) inner shelf off Virginia (3)Hampton Roads port area(4)Chesapeake Bay-York River-Pamunkey River estuary(5)oligohaline part of James River. Shannon's diversity values were highest on outer shelf. Values for inner shelf were considerably lower than those for the outer shelf or polyhaline areas. Low diversity on shallow shelf may be due to physically controlled environment. In Hampton Roads area, diversity highest in sands than that of muds, presumably because of greater variety of microhabitats.

Boesch, D. F., Day, J. W., Jr., and Turner, R. E., 1984, Deterioration of coastal environments in the Mississippi Deltaic Plain: options for management, in Kennedy, V. S., ed., The estuary as a filter: Orlando, Academic Press, p. 447-466.

MARSHES, DELTA, ACCRETION RATES, MARSH LOSS, MISSISSIPPI DELTA, CHANNELIZATION, SPOIL, WETLANDS

Table 1 is a summary of marsh accretion rates for Louisiana and east coast. Rapid changes in coastal environments in the Mississippi deltaic plain in southeastern Louisiana are attributed to (1) natural decay of delta lobes, (2) channelization of river flow, and (3) dredging activities. The result is the conversion of coastal wetlands to open water and uplands at a rate of 100 sq km/yr, which is 85% of the U. S. loss rate. Among options to reduce wetland loss: "management of new delta accretion and regulatory control and rehabilitation of channelized wetlands are likely the most effective."

Boesch, D. F., and Turner, R. E., 1984, Dependence of fishery species on salt marshes: the role of food and refuge: *Estuaries*, v. 7, no. 4a, p. 460-468.

SALT MARSHES, REFUGE, NURSERIES, FISH, CRUSTACEANS, PRODUCTIVITY, HABITATS, INVERTEBRATES, VASCULAR PLANTS, TROPHIC

Evidence supports the concept that juveniles of many economically important fishes and invertebrates heavily utilize the shallow habitats associated with the edges of salt marshes and that this is coincident with regional trends of increased yield of those species in areas with extensive salt marshes. Many small crustaceans, annelids, and meiofauna, which are potential prey of fishery species, are known to consume organic detritus. More than 98% of the detrital material exported from a Gulf Coast salt marsh consisted of amorphous aggregates which were produced primarily by benthic macroflora on the marsh surface rather than by microbial decomposition of the vascular plants. "Furthermore, marsh grass production is essential to the perpetuation of this structurally complex habitat, because marshes must continue to accrete sediments and vascular plant detritus (peat) and to grow upward to survive. (Hatton et al. 1983)."

Boucot, A. J., 1981, Principles of benthic marine paleoecology: Academic Press, New York, 463 p.

SUBSTRATE, SEDIMENT, PALEOECOLOGY, BENTHOS, TURBIDITY, COMMUNITIES, PREDATION, SAND, MUD, COARSE GRAIN, INFAUNA, EPIFAUNA

Textbook containing good discussion of substrate-faunal relationships, including substrate selection, antibiosis, changes in substrate during ontogeny, level-bottom environments and general considerations of animal-substrate relationships, turbidity, sediment stability, and many others. This text applies to paleo and recent benthic communities.

Brehmer, M. L., 1965, Turbidity and siltation as forms of pollution: *Journal of Soil and Water Conservation*, v. 20, no. 4, p.132-133.

TURBIDITY, SILTATION, POLLUTION, ESTUARY, SUSPENDED SEDIMENT

General article on turbidity and effects on biota. "The productivity of the estuarine environment is basically dependent upon the primary producers within the system. These planktonic plants are

dependent upon sunlight as the energy source for photosynthesis: as the suspended solids content of the water increases, the depth of light penetration decreases. Therefore, as the depth of light penetration decreases the compensation depth-the level at which the rate of photosynthesis equals the rate of respiration-decreases, or approaches the surface. Theoretically, if the respiration exceeds the productivity, the system is characterized by a negative energy flow and the biological community undergoes degradation. The temporary suspensoids may undergo peptization and dispersion during the period of suspension. Prior to settling, flocculation and aggregation occur. In the process, a large proportion of the microscopic plants are mechanically entrapped and are carried to the bottom with the particles. The effects of inorganic suspensoids on zooplankton and higher aquatic life are even....."

Byrne, J. R., 1975, Holocene depositional history of Lavaca Bay, central Texas Gulf Coast: The University of Texas at Austin, Ph.D. dissertation, 149 p.

LAVACA BAY, TEXAS, HOLOCENE, DEPOSITIONAL HISTORY, SEDIMENTS, TEXTURES, CLAYS, SUSPENDED LOAD, COLORADO RIVER, BRAZOS RIVER, LAVACA RIVER, NAVIDAD RIVER, SEDIMENT BUDGET

Bay-estuarine sediments of Lavaca Bay have three major sources of mud: (1) suspended load transported by the Colorado-Brazos fluvial complex, which was discharged into marine and estuarine bodies about 83 km (50 mi) to the northeast and transported southwest by longshore currents, (2) erosion of Pleistocene sediments along the bay margins, and (3) suspended load from the Lavaca-Navidad fluvial complex. It is estimated that there has been sufficient sediment from the Lavaca-Navidad fluvial complex and bluffs along the bay margin to have filled the bay since stillstand, however the sediment accumulation rate in the bay is slow because significant quantities of sediment are transported out of the bay by waves and surface currents. The Lavaca-Navidad fluvial-deltaic complex has prograded over transgressive sediments since stillstand. A rough argillaceous sediment budget is presented in the dissertation.

Cammen, L. M., 1976, Macroinvertebrate colonization of *Spartina* marshes artificially established on dredge spoil: Estuarine and Coastal Marine Science, v. 4, p. 357-372.

MACROINVERTEBRATES, SPARTINA, DREDGE SPOIL, MARSHES, BIOMASS, DIVERSITY, SEDIMENT, PARTICLE SIZE, ELEVATION, SEDIMENTATION RATE, FAUNAL AFFINITY, SPECIES RICHNESS

From abstract-"Core samples were taken from dredge spoil planted with *Spartina alterniflora*, spoil left bare and nearby natural marsh in two locations in order to determine the factors influencing the development of the spoil fauna and to investigate the relationship of the spoil fauna with the natural marsh fauna. Based on calculation of total numbers, total biomass, sample diversity and species richness and faunal affinity between plots, the difference in elevation between the bare and planted spoil plots appeared to be the major factor determining the degree of similarity in their fauna. Five

factors are suggested to control the development of the planted spoil fauna: the similarity of the spoil to natural marsh in elevation and sediment particle size, the natural sedimentation rate in the area, the proximity of the spoil to natural marsh and the relative maturity of the natural marsh faunal community."

Cammen, L. M., 1979, The macro-infauna of a North Carolina salt marsh: The American Midland Naturalist, v. 102, no. 2, p. 244-253.

NORTH CAROLINA, SALT MARSH, SPARTINA ALTERNIFLORA, INFAUNA, BIOMASS, EPIFAUNA, DETRITUS

From abstract-"Annual production of the macro-infauna was estimated to be 5.9g ash-free dry weight m², similar to that found in other North Carolina salt marshes. Production values for salt-marsh infauna are comparable to those found for salt-marsh epifauna." Infauna was generally most abundant in late winter and early spring and least in summer and early autumn. The salt marshes in North Carolina were dominated by only a few infaunal taxa. All the most important species in the marsh infauna are selective deposit or detritus feeders or omnivores with the possible exception of Eteone heteropoda and Glycera americana. Much of the sediment and detritus in the marsh is reworked each year by the infauna.

Carriker, M. R., 1967, Ecology of estuarine benthic invertebrates: a perspective, in Lauff, G. H., ed., Estuaries, Publication 83, American Association for the Advancement of Science, p. 442-487.

ESTUARINE, BENTHOS, ECOLOGY

This chapter is a thorough literature review on the ecology of the estuarine benthos, including suspended sediments and sedimentation and estuarine bottom sediments. Statement from pg. 466-"The observations of Brett, McNulty, and Sanders explain in general terms the nature of the faunally rich "muddy sand" frequently mentioned in the literature. In estuaries, these appear to be more or less optimum mixtures of clays, silts, fine sands, and organic matter which support greater densities of megabenthic populations than either clean coarse unstable sands and gravels at the mouth or soft slurry muds in the quiet sheltered reaches of estuaries."

Chapman, V. J., 1960, Salt marshes and salt deserts of the world: Leonard Hill, New York, 392 p.

SALT MARSHES

Sedimentation rates reported: 13 mm/yr *Spartina patens*; 6 mm/yr *Spartina alterniflora*

Coastal Ecosystems Management Inc., 1972, Environmental assessment of the Trinity River discharge on productivity in Trinity Bay: Final report prepared for Fort Worth District, Army Corps of Engineers, contract no. DACW63-72-C-0142, 73 p.

TRINITY BAY, PRODUCTIVITY, SEDIMENT, TRINITY RIVER, BENTHIC INVERTEBRATES, SAND, DIVERSITY

"The patterns of invertebrate populations are similar to those found in other Texas bays, with the highest populations being found in poorly sorted, sandy to silty sand sediments. The more loosely compacted the sediments, the more interstitial spaces are available for animal life....Total populations of invertebrates are greater in Trinity Bay than in other Texas bays."

Copeland, B. J., 1970, Estuarine classification and responses to disturbances: Transactions, American Fisheries Society, v. 4, p. 826-835.

MARINE GRASS, MARSH, PLANKTON-BASED SYSTEM, RIVER MOUTH, TURBIDITY, IMPOUNDMENT, CLAM FLAT

Response of six coastal systems are related to impoundment and turbidity. The 6 systems are grass bottom, plankton-based, river-mouth, coral reef, marsh, and clam flat.

Crisp, D. J., 1984, Energy flow measurements, in Holme, N. A., and McIntyre, A. D., eds., Methods for the study of marine benthos: London, Blackwell Scientific Publications, p. 284-372.

BIOMASS, PRODUCTIVITY, BENTHOS, ENERGY FLOW

Lengthy discussion on methods and uses of biomass and energy flow measurements.

Driscoll, E. G., 1967, Attached epifauna-substrate relations: Limnology and Oceanography, v.12, p. 633-641.

SUBSTRATE, SEDIMENT, EPIFAUNA, SUSPENSION FEEDER, BYSSUS, SHELL, SILT, CLAY, SAND, BUZZARDS BAY, MASSACHUSETTS, CALCIUM CARBONATE, BORING SPONGE, DEPTH, BIVALVE, CHITON, GASTROPOD, INFAUNAL

Samples were taken at 51 stations in Buzzards, Mass. from June to September, 1965. Nine of the most abundant epifaunal suspension feeders, including *Anomia simplex*, *Crepidula convexa*, *C. fornicata*, *C. plana*, *Anadara transversa*, *Cliona celata*, and others, were examined to determine distribution in relation to temperature, depth, mean grain diameter, silt-clay content, dead shell abundance, and calcium carbonate content. All species except *Crepidula* are more abundant on bottoms with low silt-clay content and a mean grain diameter in the medium sand range. "They are more plentiful in sediments coarser than those in which infaunal suspension feeders reach their maximum abundance. It is suggested that the stronger currents associated with coarser sediments provide an increased food supply that results in their more plentiful occurrence."

Driscoll, E. G., 1975, Sediment-animal-water interaction, Buzzards Bay, Massachusetts: Journal of Marine Research, v. 33, no. 3, p. 275-302.

BUZZARDS BAY, MASSACHUSETTS, SEDIMENT, BENTHOS, MICROORGANISMS, FEEDBACK, ORGANICS, STANDING CROP, FECAL FORMATION, DEPOSIT FEEDERS, BIOTURBATION, BIODEPOSITION, TEMPERATURE

"Characteristics of sediments, benthic animals and bottom water are all interrelated in Buzzards Bay. Seasonal variations in the organic content of the sediment are due largely to variations in the standing crop of benthic microorganisms. Bioturbation and fecal formation by deposit feeders result in increased surface area for colonization by microorganisms. Biodeposits further represent a readily available nutrient source for such organisms. It is suggested that a feedback relationship exists, increasing microorganisms resulting in increasing deposit feeder abundance. The latter, through bioturbation, and biodeposition, produce an environment suitable for increasingly abundant microorganisms. The rate of this feedback is temperature dependent."

Espey, Huston and Associates, Inc., 1978, Hydrological and biological studies of the Colorado River delta, Texas: Texas Department of Water Resources, LP-79, 165 p.

COLORADO RIVER, DELTA, STREAMFLOW, MARSHES

Among the points of interest made in this report are: Model studies show that marsh zones on the delta are not inundated by riverine flows (p. 1-3); "Model computations and field observations confirm that, with few exceptions, no part of the delta west of the river is inundated by freshwater overbanking from the Colorado River during flood events. All nutrients transported by the river are contributed directly to either the Gulf of Mexico or Matagorda Bay." (p. 1-5) Simulated inflow studies of the Colorado River under typical tidal conditions showed that approximately 80% of the river flow above Tiger Island Cut was diverted through the cut at river flows of 2,000 cu-ft/sec. This percentage decreased as river flow increased to a rate of approximately 8,000 cu-ft/sec; at flow rates greater than this the diversion through Tiger Island Cut remained at a constant 62% (p. 6).

Faris, O. A., 1933, The silt load of Texas streams: U.S. Department of Agriculture Technical Bulletin No. 382, Washington, D.C., 71 p.

STREAMS, SEDIMENT LOAD, TEXAS, COLORADO RIVER, RESERVOIR SEDIMENTATION

Estimated that the mean annual volume of silt entering the Austin Reservoir from the Colorado river for a three year period before 1933 was 11,500 acre-ft/yr (p.45). This is the figure used by van Beek and others (1981) to determine suspended sediment load carried by the Colorado River before 1933.

Flessa K. F., Constantine, K. J., and Cushman, M. K., 1977, Sedimentation rates in a coastal marsh determined from historical records: Chesapeake Science, v. 18, no. 2, p. 172-176.

SEDIMENTATION RATES, MARSH, SPARTINA ALTERNIFLORA, NEW YORK

Historical records indicate that Flax Pond, a small *Spartina alterniflora* marsh located on the north shore of Long Island, New York, was opened to marine waters in 1803. This opening is recorded in the sediments by a sharp transition from sedge (brackish or fresh water) peat to *Spartina* (salt marsh) peat. This dated horizon was used to calculate an average net rate of vertical accretion of 2.5 mm/yr and a maximum net rate of 4.7 mm/yr. These estimates may serve to predict the rate at which *Spartina* marshes are able to recover from physical disturbance."

Flint, R. W., 1985, Long-term estuarine variability and associated biological response: *Estuaries*, v. 8, no. 2A, p. 158-169.

ESTUARIES, BIOMASS, SHRIMP, CORPUS CHRISTI BAY, BENTHOS, NUTRIENTS, PRODUCTIVITY

Article concerning role of benthos in Corpus Christi estuary. From the section on "biological response"-"After the episodic rainfall of September 1979, benthic macroinfaunal communities exhibited large increases in abundance and biomass of dominant populations. Paralleling the 1979 increases in benthos production were significantly larger catches of shrimp. On a monthly basis, increased shrimp yields, were observed at the end of 1979 and early part of 1980, periods which immediately followed the September rainfall and correlated with increased production of the benthic community."

Flint, R. W., 1984, Phytoplankton production in Corpus Christi Bay: *Contributions in Marine Science*, v. 27, p. 65-83.

CORPUS CHRISTI BAY, PHYTOPLANKTON, TURBIDITY, ESTUARY, PRODUCTIVITY

From p. 77-"The indication that water column turbidity limits the amount of primary production realized by the Corpus Christi Bay estuary, both from the regression analyses and the comparison of 0.5 m with 1.0 m production rates, best illustrated why this estuarine ecosystem is not as productive as others around the world. Based upon data from a number of estuaries around North America, the Corpus Christi Bay estuary was relatively oligotrophic in comparison. The greatest annual primary productivity for an estuary was estimated at 532g C/m²/yr. The Corpus Christi Bay estuary, which over a three-year study period exhibited a mean phytoplankton productivity rate of 174.1 g C/m²/yr, was at the lower end of the scale."

Flint, R. W., Rabalais, S. C., and Kalke, R. D., 1981, Quantification of extensive freshwater input to estuarine benthos: The University of Texas Marine Science Institute, Port Aransas, report to Texas Department of Water Resources, contract no. IAC(80-81), 55 p.

CORPUS CHRISTI BAY, CHANNEL, SHOAL, BENTHOS, SEDIMENT, SAND, MUD

Long-term sampling was conducted at 3 channel and 3 shoal stations in Corpus Christi Bay. Muddy and shelly sands were present at shoal stations and clays, fine sands, and silts dominated the channel

stations. In the channel a 1-2cm layer of oxidized brown silt was present at the very surface of these sediments. Little biogenic activity observed at channel stations, but there were numerous tube structures in the shoal sands. The channel stations exhibited much lower diversities and densities than the shoal stations. Differences in the influences of dredging, tanker traffic, and shrimp trawling also affected the benthos of the two areas.

Flint, R. W., Rabalais, S., and Kalke, R., 1982, Estuarine benthos and ecosystem functioning, in Davis, J. R., ed., Proceedings of the symposium on recent benthological investigations in Texas and adjacent states: Texas Academy of Science 85th annual meeting, San Angelo, Texas, p. 185-201.

ESTUARY, BENTHOS, ECOSYSTEM, CORPUS CHRISTI BAY, BIOMASS, TURBIDITY, SECONDARY PRODUCTION, SEDIMENT, SEDIMENT METABOLISM, PRIMARY PRODUCTIVITY

"Trubidity of the water column appeared to play a principal role in regulating primary productivity rates. On numerous occasions, especially at Station 2, photosynthesis was decreased by as much as 30% at 1 m depth compared to rates measured at 0.5 m depth. In contrast, there appeared to be relatively little relationship between primary productivity rates and corresponding water column ammonia-nitrogen concentrations. The correlations between ammonia concentration and photosynthetic rate was $r=0.23$, which was not significant at p less than 0.10. Sediment metabolism, indicative of benthic community activity and production, ranged from a low at station 2 in January to a high at station SR4" Benthic communities at station SR4 had the highest carbon production rates. Percent sand at station SR4 was greater than 70%, with 30% at station 2 in Nueces Bay, and 5% at station 7 in mid-Corpus Christi Bay.

Flint, R. W., and Kalke, R. D., 1985, Benthos structure and function in a South Texas estuary: Contributions in Marine Science, v. 28, p. 185-201.

BENTHOS, BIOMASS, CORPUS CHRISTI BAY, NUECES BAY, ECOSYSTEM, SEDIMENT, MACROINFAUNA, FLUVIAL

From abstract-"The Corpus Christi Bay estuary in the northwestern Gulf of Mexico was investigated for spatial and temporal variation of benthos community structure and function. Four stations were sampled quarterly for 2.5 years to investigate for macroinfaunal species assemblage changes as well as changes in benthic metabolism and nutrient regeneration. Cluster analysis of species assemblages illustrated that community structure changed from the riverine-influenced end of the estuary to the oceanic-influenced end. Taxa number increased away from the fluvial source while total abundance decreased. Maximum biomass was observed in the middle estuary region. Although spatial patterns were evident, no consistent seasonal patterns were observed for community structure characteristics from one site to the next. Community functional processes were not significantly different between sites. Metabolism did not show consistent temporal patterns but sediment nutrient flux always exhibited peak rates during the summer at all sites. Sediment texture differences as well as variability in salinity between sites were thought....."

Frey, R. W., and Basan, P. B., 1985, Coastal salt marshes, in Davis, R. A., Jr., ed., Coastal sedimentary environments, second revised, expanded edition: New York, Springer-Verlag, p. 225-301.

SALT MARSHES, SEDIMENTATION, VEGETATION, GULF COAST, ATLANTIC, PACIFIC, SOURCES OF SEDIMENT, PROCESSES

Good discussion on processes affecting sedimentation in salt marshes, including sources of sediments, water movement on marshes, hydrographic setting, tidal hydraulics and sedimentation, sediment trapping, flocculation, seasonality of sediment supply, physiochemical conditions, biological conditions, deposition and erosion along channels, dynamics of the marsh surface, and sedimentary structures. "In terms of hydraulics, the main difference between river flood and tidal flood is the duration; river slack allows suspended sediments much more time to settle out than does tidal slack." The observation that where water is impounded, artificially or naturally, the deposition rates are significantly higher suggests that suspended sediments ordinarily do not have sufficient time to settle before tide-turn but would settle if the duration of slack or sluggish water were somehow increased. Even in densely vegetated Danish marshes, no more than half the suspended matter is deposited (Schau, 1967).

Funicelli, N. A., 1984, Assessing and managing effects of reduced freshwater inflow to two Texas estuaries, in Kenedy, V. S., ed., The estuary as a filter: Orlando, Academic Press, p. 435-446.

FRESHWATER INFLOW, TEXAS, ESTUARIES, NUECES BAY, CORPUS CHRISTI BAY, MATAGORDA BAY, DELTA INUNDATION

Paper includes table showing historic vs. projected marsh inundation frequency for the Nueces-Corpus Christi Bay system marshes. Table 2. Comparison of historical and projected freshwater inundations. Historical inundation (1958-1979) Projected Average flow duration (days) 9.8 13 Average one-day flood volume 17.0 19.4 (thousand acre-ft) Number of occurrences 50 26 Occurrences/yr 2.3 1.2

Gagliano, S. M., Meyer-Arendt, K. J., and Wicker, K. M., 1981, Land loss in the Mississippi River deltaic plain: Gulf Coast Association of Geological Societies, Transactions, v. 31, p. 271-273.

WETLANDS, LAND LOSS, DELTAS, MISSISSIPPI RIVER, LOUISIANA, MARSHES

Determination of land loss was based on systematic measurements and comparisons of historic maps and photographs. The comparisons show that the long-term trend of net progradation persisting through most of the past 5000 yrs reversed during the late 19th century, and land-loss rates have accelerated geometrically during the 20th century. In the study area land-loss rates have increased from approximately 6.7 sq mi/yr in 1913 to 39.4 sq mi/yr in 1980. Greatest loss has been in wetlands. Apparent causes include harnessing Mississippi River by levees and control structures reducing the natural processes such as upstream diversion, overbank flooding, and crevassing

(processes responsible for extensive sedimentation and deltaic plain progradation) and funneling valuable sediments to deep offshore water. The elimination of the natural processes coupled with extensive canalization and accelerated subsidence related to hydrocarbon extraction has led to the serious land loss problems.

Gillard, R. M., 1974, Distribution, abundance, and species diversity of macrobenthic and meiobenthic invertebrates in relation to Houston Ship Channel pollution in upper Galveston Bay and Tabbs Bay, Texas: Texas A&M University, Ph.D. Dissertation, 186 p.

GALVESTON BAY, TABBS BAY, MACROBENTHOS, MEIOBENTHOS, SEDIMENT, PERCENT SAND, DIVERSITY, INFAUNA, EPIFAUNA

Benthic samples were collected at 16 stations in upper Galveston and Tabbs Bays over a 18 month period. In the sediment section, a statement from page 65- "the total number of species reached a peak when the percent sand was approximately 44 to 54% and decreased when the percent sand was higher or lower. Although there were more fluctuations, a similar relationship occurred with the infaunal and primarily epifaunal species. It is interesting that an increase in epifaunal species was generally accompanied by an increase in infaunal species. There was essentially the same relationship between the relative amount of sand and the mean number of species as there was with the total number of species." Diversity indices were highest at stations having between 44 to 51% sand.

Gilmore, G., Dailey, J., Garcia, M., Hannebaum, N., and Means, J., 1976, A study of the effects of fresh water on the plankton, benthos, and nekton assemblages of the Lavaca Bay System, Texas: Texas Parks and Wildlife Department, Technical report to the Texas Water Development Board, January, 1973-June, 1975, 113 p.

LAVACA BAY, BENTHOS, SEDIMENT, STANDING CROP, DIVERSITY,

"Benthic standing crop was significantly related to salinity, turbidity, total carbon, organic nitrogen, and nitrate. Thus inflow (freshwater) may have been indirectly related to standing crop." Diversity declined from the high salinity lower bay to the low salinity upper bay and river area. Diversity was highest during late winter and early spring when sustained freshwater inflow was generally low. Low inflow was associated with high salinity, low turbidity, and low nutrient concentrations. Species diversity was positively related to bottom salinity and negatively related to bottom turbidity and nutrients. Diversity at each site did not appear to be related to sediment particle size distribution at that site.

Gilmore, G., and Trent, L., 1974, Abundance of benthic macroinvertebrates in natural and altered estuarine areas: Seattle, Washington, U.S. Department of Commerce, National Oceanic and Atmospheric Administration Technical Report NMFS SSRF-677, 13 p.

WEST GALVESTON BAY, TEXAS, MACROINVERTEBRATES, DREDGING, BULKHEADING, FILLING, MARINE GRASSES, CANALS, MUD, ALGAE, MARSH, OPEN BAY

Numeric and volumetric abundance was examined for a marsh, canal and open bay area in West Bay. Benthic invertebrates were slightly more abundant and over twice as abundant volumetrically in the marsh than in the canals; they were least abundant in the bay. The marine grass taken in the samples was *Halodule wrightii*.

Ginsburg, R. N., and Lowenstam, H. A., 1958, The influence of marine bottom communities on the depositional environment of sediments: *Journal of Geology*, vol. 62, p. 249-265.

SEAGRASSES, COMMUNITIES, SEDIMENT, HABITAT, STABILIZING

Marine grasses help trap and stabilize sediment. "In addition to the trapping and fixing of sediment, marine grass carpets also create a habitat for other organisms. In this way the presence or absence of the grasses can produce modifications in the distribution of organisms."

Goldman, C. R., 1971, Biological implications of reduced freshwater flows on the San Francisco Bay-Delta system, in Seckler, D., ed., *California water, a study in resource management*: University of California Press, Berkeley, p. 109-124.

SAN FRANCISCO BAY, DELTA, FRESHWATER INFLOW, SEDIMENT, EUTROPHICATION, NUTRIENTS, CALIFORNIA, SUSPENDED SEDIMENT, TURBIDITY, ESTUARY, TRANSPARENCY

From p. 116-"a diversion of fresh water from the estuary will greatly reduce the sediment inflow. Although a lesser amount of sediment means increased transparency of the water, inorganic turbidity in the Bay waters provides certain distinct ecological benefits. Decreased turbidity would result naturally from decreased inflow of sediment-bearing water, a longer time for sedimentation to occur, and decreased water turbulence which aids in maintaining sediment in suspension. From p. 120-"Even if the turbidity per unit volume of water were the same, a reduced rate of flow would bring less sediment into the Bay and Delta during the same period of time. This alone would increase the transparency of the water. In the unlikely event that nutrient conditions could remain unchanged, we would then have a large increase in the growth of higher aquatic plants such as pond weeds and bulrushes, which are now shaded out in deeper water by the sediment and algae in suspension. But in all probability, the decrease in inorganic turbidity would be accompanied by an in planktonic algae in response..

Gray, J. S., 1974, Animal-sediment relationships, in Barnes, H., ed., *Oceanography and Marine Biology-an Annual Review*, v. 12, p. 335-344.

DIVERSITY, STRUCTURAL COMPLEXITY, SEDIMENT, MEAN PARTICLE SIZE, STABILITY-TIME HYPOTHESIS

Structurally complex sediments are the most diverse. "Mean particle size clearly does not reflect structural complexity. Characters such as sorting coefficients give a better indication of the grain size complexity." Data from the sediment infauna of the European boreal region show that the coarse deposits have low diversity, the medium size deposits the highest diversity, while in muds diversity falls. Mixed sediments have the highest diversity. Coarse and heterogeneous sediments seem to have higher diversities than homogeneous sediments. Another factor relating to diversity is the stability-time hypothesis. This hypothesis states that physical instability in an environment or short time-periods prevent the establishment of diverse communities. In a stable environment with time, speciation and immigration will lead to high diversity. This hypothesis would be applicable in a bay environment.

Harper, D. H., and Hopkins, S. H., 1976, The effects of oyster shell dredging on macrobenthic and nektonic organisms in San Antonio Bay, in Bouma, A. H., ed., Shell dredging and its influence on Gulf Coast environments: Houston, Texas, Gulf Publishing Company, p. 232-279.

SHELL DREDGING, OYSTER, CRASSOSTREA VIRGINICA, SAN ANTONIO BAY, BENTHOS, SALINITY

Benthic organisms were divided into two groups, those in the low-salinity upper bay and those in the high-salinity, lower bay. Benthic populations increased dramatically following a spring flood, and they concluded that the bloom was due to large increases in nutrients from increased river flow and only indirectly to decreased salinities.

Harrison, E. Z., and Bloom, A. L., 1977, Sedimentation rates on tidal salt marshes in Connecticut: Journal of Sedimentary Petrology, v. 47, no. 4, p. 1484-1490.

SALT MARSHES, SEDIMENTATION RATES, ARTIFICIAL MARKERS, LONG ISLAND SOUND, CONNECTICUT

Rates of sediment accumulation from 1963-1973 on five high marsh sites on the Connecticut Coast of Long Island Sound ranged from 2.0 mm/yr to 6.6 mm/yr. The rates are correlated with tidal range (which varied from 0.75m to 2.2m)--the highest sedimentation rates are on marshes with the greatest tidal range. Several ounces of glitter were teased over 1 m² plots with a new layer of different colored glitter added each year from 1962 to 1966. Much of the inorganic marsh is derived from a marine source as indicated by decreasing grain size of sediment upstream away from the Sound and illite:vermiculite ratio of the sediment. A sedimentation rate of 7.1 mm/yr from 1963-1973 was measured where *Spartina patens* salt marsh is replaced by *Phragmites communis*. Years with fewer average storms show less sediment accretion suggesting that a few storms each year may account for most of the annual sedimentation. Month to month observation would be required to test this hypothesis.

Hatton, R. S., DeLaune, R. D., and Patrick, W. H., Jr., 1983, Sedimentation, accretion, and subsidence in marshes of the Barataria Basin, Louisiana: *Limnology and Oceanography*, v. 28, no. 3, p 494-502.

MARSHES, SEDIMENTS, SEDIMENTATION RATES, 137 CS, LOUISIANA.

Vertical accretion rates in marshes measured vary from about 13 mm/yr in levee areas to 7 mm/yr in back marshes. Mineral sediment content of the marsh soil profile decreased with distance from the coast. Autochthonous organic matter appears to be an important factor defining the process and rate of vertical accretion, especially in freshwater marshes. Except in natural levee areas, marsh accretion rates are less than subsidence measured by water level data, however, this alone cannot account for observed land-loss patterns in the basin area.

Hatton, R. S., Patrick, W. H., Jr., and DeLaune, R. D., 1982, Sedimentation, nutrient accumulation, and early diagenesis in Louisiana Barataria basin coastal marshes, in Kennedy, V. S., ed., *Estuarine comparisons*: Academic Press, New York, p. 255-267.

SEDIMENTATION, RATES, NUTRIENT ACCUMULATION, DIAGENESIS, MARSHES, LOUISIANA, BARATARIA BASIN, CS 137, CHEMICAL ANALYSIS

Abstract: "Vertical marsh accretion, along with sediment and nutrient accumulation, were examined using 137 Cs dating and chemical analysis of soil cores taken from various marshes in Louisiana's Barataria Basin. Vertical accretion averaged 1.3 cm/yr and 0.7 cm/yr in levee and backmarsh areas respectively. Inorganic sedimentation increased fourfold from freshwater through salt marshes along a transect perpendicular to the Gulf Coast. The accretionary role of inorganic sediments is variable according to marsh type. Mineral sediment being deposited in the marsh is the principle source of iron, manganese, and phosphorus, but not of organic carbon or nitrogen. Manganese and phosphorus appear to be subject to substantial post-depositional remobilization."

Haven, D., and Morales-Alamo, R., 1972, Biodeposition as a factor in sedimentation of fine suspended solids in estuaries, in Nelson, B., ed., *Environmental framework of coastal plain estuaries*: Geological Society of America Memoir No.133, p. 121-130.

BIODEPOSITION, SUSPENDED SOLIDS, ESTUARIES, SEDIMENTATION, BIODEPOSITS, SUSPENSION-FEEDERS, DETRITUS, JAMES RIVER, VIRGINIA, OYSTERS, BARNACLES, FECAL PELLETS, PSEUDOFECES,

Biodeposits from suspension-feeders form a recognizable and measurable component of the total suspended load in the James River, Virginia. "Quantity of total solids and pellets in suspension varied with time, with maximum quantity occurring at peak current velocities and minimum values occurring at slack water. This fluctuation implied alternate suspension and deposition of biodeposits. The process of biodeposition may be at least as important as the abiotic process in initiating deposition of fine suspended solids." Lund (1957) suggested that one acre of oysters might deposit 7.6 metric tons of detritus annually.

Hines, A. H., and Comtois, K. L., 1985, Vertical distribution of infauna in sediments of a subestuary of central Chesapeake Bay: *Estuaries*, v. 8, no. 3, p. 296-304.

INFAUNA, CHESAPEAKE BAY, SEDIMENT, VERTICAL DISTRIBUTION, SAND, MUD, RHODE RIVER, DEPOSIT FEEDERS, INFAUNA, BIOTURBATION

The vertical distribution of macroinfaunal species in sand and mud sediments of the Rhode River, a subestuary of central Chesapeake Bay, is described. Infaunal communities of the two sediment types were composed of 15 common species, as well as several relatively uncommon species. The total density and standing crop of sediment type were similar. There was extensive overlap in the species inhabiting both sediment types, with only four species in mud and five species in sand not occurring in the other sediment type. For both sediment types, most organisms occurred in the upper 5 cm of sediment, with few occurring deeper than 20 cm. However, the greatest fraction of biomass of infauna was composed of a few relatively large organisms occurring from 10 to 30 cm in mud and from 10 to 25 cm in sand. Almost no organisms or biomass were found deeper than 30 cm in either sediment. Some species exhibited significant temporal changes in density, but not biomass and some had significant temporal changes in vertical distribution of both density and biomass.

Holland, A. F., Mountford, N. K., and Mihursky, J. A., 1977, Temporal variation in upper bay mesohaline benthic communities: mesohaline benthic communities: 1. the 9-m mud habitat: *Chesapeake Science*, v. 18, no. 4, p. 370-378.

CHESAPEAKE BAY, BENTHIC, MESOHALINE, MUD, COMMUNITIES, HABITAT, SEASONAL, RECOLONIZATION, SUBSTRATE

"This study demonstrated that spatial distribution was primarily controlled by substrate characteristics which change with depth. Along the substrate depth-gradient several communities were identified. These were: a 3-m sand community; a 6-m shell community; a 9-m mud community; and a 6-m muddy-sand transitional region."

Holland, J. S., Maciolek, N. J., Kalke, R. D., and Oppenheimer, C. H., 1975, A benthos and plankton study of the Corpus Christi, Copano and Aransas Bay systems: report on data collected during the period July 1974-May 1975 and summary of the three-year project: The University of Texas at Port Aransas Marine Science Institute, Final report to the Texas Water Development Board, 171 p.

BENTHOS, SEDIMENT, SHELL, CORPUS CHRISTI BAY, NUECES BAY, COPANO BAY, ARANSAS BAY, STANDING CROP, OYSTER REEF, CLUSTER

"In our study area the monthly benthos standing crop values are generally strongly correlated with the amounts of shell found in the sediment." Depauperate sites in bays had several factors in common; little or no shell in the sediment and periodically lowered salinities. "Benthic populations seem to be

directly correlated to salinity, high salinities yielded larger standing crops and greater diversities at all sites. Lowered salinity regimes after extreme floods always showed lowered standing crops and diversities at most sites so affected." There was also a cluster of organisms consistently associated with live oyster clumps.

Isphording, W. C., 1986, Apalachicola Bay: dynamic sedimentation in a Gulf Coast estuary: Transactions of the Gulf Coast Association of Geological Societies, v. 36, p. 471-488.

SEDIMENTS, ESTUARY, BAY, SEDIMENTATION RATES, RIVERS, DAMS, SAND, SILT, CLAY

Changes in sediment regimen in Apalachicola Bay over past 25 yrs traced to dams on rivers. Changes are toward sediments rich in sand and devoid of silt. Concludes that removal of sand and silt by reservoirs increases erosive power of river downstream from dams (closest dam about 100 mi away from bay). Clay passes over spillway and sand is eroded as channel is deepened and widened. More sand now being supplied to delta where accumulation rates exceed 10 mm/yr; overall rate for entire bay is 6.7 mm/yr. This is the highest rate for any bay in the entire northern Gulf of Mexico. As noted by Bedosky (1986), this rate combined with lack of appreciable submergence will infill the bay many years before it would have been infilled by natural geologic processes. Rates of deposition for other bays presented in table 4: Narragansett--0.65mm (Morton, 1967); Delaware--1.8mm (Ootsdam, 1971); Chesapeake Bay--3.7mm (Biggs, 1970); Mobile--0.52mm (Ryan, 1969). With regard to river sediment load: Davis (1978) concluded that probable average distribution for all rivers in eastern U.S.-- 10% sand, 45% silt, 45% clay for a ratio of 2:9:9.

Johnson, R. G., 1971, Animal-sediment relations in shallow water benthic communities: Marine Geology, v.11, p.93-104.

BENTHOS, SEDIMENT, POLYCHAETES, MOLLUSKS, TOMALES BAY, CALIFORNIA, DIVERSITY, BRILLOUIN'S EQUATION, ENVIRONMENTAL STABILITY, COMMUNITY, SANDS, SILT, CLAY, SUBSTRATE, MUD, CORRELATION COEFFICIENT, SUSPENSION, DEPOSIT

Most benthic species are found in a particular substrate type. Individuals occasionally found in other substrates. Mud species are better able to invade sands than sand species invade mud. This is probably related to feeding and respiration as most clean sand species probably cannot tolerate large amounts of silt and clay while mud species have special adaptations to do so. Many of the sand species are suspension feeders and mud species are deposit feeders. "Mud species are found to occur primarily in association with the most diverse assemblages in muddy sands and clean sands. Samples collected in Tomales Bay, California. Substrate was treated as an index of environment. Substrate reflects water movement, distance from shore, food supply, and other features of environment. In a local situation, however it does not reflect salinity or temperature except as these variables might be geographically related to sediment distribution.

Kalke, R. D., 1986, Benthos, in Jones, R. S., and others, Studies of freshwater inflow effects on the Lavaca River Delta and Lavaca Bay, Texas: Final report to Texas Water Development Board, contract no. IAC(86-87)0757, p. 5.1-5.41.

LAVACA BAY, LAVACA RIVER, FRESH WATER, BIOMASS, BENTHOS, CHIRONOMID LARVAE, MIDGES, INFLOW, MOLLUSCS

From conclusions p.5.6-"Benthic communities in upper Lavaca Bay varied between the two years but it is difficult to relate these variables to freshwater inflow. The only species which were affected by inflow on a short term was the aquatic chironomid larva and *Hobsonia florida* which had a lag response to inflow. The molluscs, *Mulinia lateralis* and *Macoma mitchelli* both increased in the low flow period which resulted in higher benthic-biomass. Once these effects were separated from the rest of the benthos no apparent differences between the two years were discernable. Higher biomass in year 2 was a result of increased mollusc populations while the remaining biomass for the rest of the benthos was similar in Year 1 and 2. Lower density and higher biomass in the deeper sediment indicates that most of the deep burrowing infauna are large individuals. Although some distributional patterns were apparently related to fluctuations in freshwater inflow, longer term monitoring is necessary to distinguish between seasonal cycles and freshwater inflow events."

Kanes, W. H., 1970, Facies and development of the Colorado River delta in Texas, in Morgan, J. P., and Shaver R. H., eds., Deltaic sedimentation, modern and ancient: Society of Economic Paleontologists and Mineralogists Special Publication No. 15, p. 78-106.

COLORADO RIVER, DELTA, SEDIMENTATION, FACIES, TEXAS, BENTHOS, PROGRADATION

Documents the progradational development of the modern Colorado River delta, and maps and describes the lithologic and biologic facies of the delta and bay depositional complex. A delta front-sheet sand was deposited during the early stages and bar-finger sands were formed when the digitate lobe was actively prograding. Distribution of sand (barfinger versus sheet) is probably related to variations in discharge and load resulting from dredging of a channel when the latter delta lobe was forming. Estimated subsidence from compaction since initial delta formation is 1 to 2 ft, but locally as much as 4 ft is documented. The sequence of facies from the surface downward are: (1) delta-plain clay and silt, root mottled, (1 to 5 ft thick) locally massive and burrowed clays of the channel-fill environment; (2) delta-front sand (2 to 8 ft thick), grades laterally into finer grained sediment; (3) prodelta reddish-brown laminated silty clays (1 to 5 ft thick); (4) bay laminated and burrow-mottled silty clays and clayey silts (10 to 14 ft thick), which lie unconformably on Pleistocene sediments.

Lane, E. W., and Borland, W. M., 1951, Estimating bed load: Transactions American Geophysical Union, 32 (1), p. 121-123.

STREAMS, BED LOAD

Ratio of bed load to suspended load: In sand-bed channels, the bed load may form 5-20% of the total load (e.g. Lane and Borland, 1951).

Larsen, P. F., 1979, The shallow-water macrobenthos of a northern New England estuary: Marine Biology, v. 55, no. 1, p. 69-78.

MACROBENTHOS, ESTUARY, NEW ENGLAND, BIOMASS, DENSITY, SILT, CLAY, SAND, SEDIMENT

"In November 1973 the benthic macrofauna of two shallow-water areas in the lower Sheepscot River estuary (New England, USA) was quantitatively sampled....Only two of the stations exhibited less than 72% silt and all but 3 of the stations could be classed as silt or clayey-silt using Shepard's nomenclature." Mean density (m²) of site-groups ranged from 289 to 2,129 and mean biomass ranged from 1.6 to 8.3 g/m². Mean diversity of site groups ranged from 3.12 to 3.82. Diversity measured by Margalef and Pielou method.

Letzsch, W. S., and Frey, R. W., 1980, Deposition and erosion in a Holocene salt marsh, Sapelo Island, Georgia: Journal of Sedimentary Petrology, v. 50, no. 2, p. 529-542.

SALT MARSH, DEPOSITION, EROSION, GEORGIA, ARTIFICIAL MARKERS

Two year study using polyester plastic markers in salt marsh indicated marsh, although quite variable, as a whole gained an estimated 6.5 mm of sediment the 1st yr and 2 mm the next. Maximum deposition was in low marsh. Seasonal deposition was minimum in autumn and maximum in summer. Seasonal erosion was affected primarily by storm frequency; rainfall intensity had little apparent influence. Paper has a good summary table of various studies and types of markers used to study sedimentation rates.

Lie, U., and Evans, R., 1973, Long-term variability in the structure of subtidal benthic communities in Puget Sound, Washington, USA: Marine Biology, v. 21, p. 122-126.

BENTHIC, PUGET SOUND, BIOMASS, DIVERSITY, ASSEMBLAGES, STABILITY

From abstract--"Data on benthic infauna from 4 permanent stations in Puget Sound off Seattle, USA, collected during 1963-1964, 1967, and 1969, revealed considerable stability in numbers of species and specimens and in diversity within stations among sampling dates. The species composition of the faunal assemblages also remained rather constant during the period of investigation, but the relative dominance among the numerically important species varied somewhat. Biomass data did not differ significantly in 1964 and 1969, but the 1967 data were considerably lower at all stations."

Loi, T.-n., and Wilson, B. J., 1979, Macroinfaunal structure and effects of thermal discharges in a mesohaline habitat of Chesapeake Bay, near a nuclear power plant: Marine Biology, v. 55, no. 1, p. 3-16.

SEDIMENT, ORGANISMS, NUMBER OF ORGANISMS, CALVERT CLIFFS, CHESAPEAKE BAY, NUCLEAR POWER PLANT, SAND, SILT, CLAY, ORGANIC CARBON

Three stations on three transects were sampled monthly in Chesapeake Bay near a nuclear power plant. "There were clear indications of differences associated with sediment types in total number of species, total number of individuals, and abundances of the dominant species. The shallower and intermediate habitats had higher sand-clay ratios, lower organic carbon contents, larger numbers of organisms than the deeper clay habitat. Within each depth, apparent effects caused by the thermal discharges on total number of species, total number of individuals, abundances of the dominant species, species diversity or evenness were found."

Loosanoff, V. L., 1962, Effects of turbidity on some larval and adult bivalves: Proceedings Gulf and Caribbean Fisheries Institute, 14th annual session, Miami Beach, Fla., p. 80-95.

TURBIDITY, LARVAL, BIVALVE, AMERICAN OYSTER, CRASSOSTREA VIRGINICA, SILT, GILLS, MOLLUSKS, ADULTS, KAOLIN

Adults and larvae of *Crassostrea virginica* and *Venus mercenaria* were used to study the effects of turbidity on these species. Turbidity was caused by natural silt and kaolin. Exposures lasted from several hours to weeks. "Very small quantities of silt sometimes stimulated normal activities of adult and larval mollusks. This may be due to absorption by normal activities of particles of suspended materials of toxic substances present in the water and to slight mechanical stimulation of the gills of the adults. However, concentrations as small as 0.1 gram/liter significantly reduced the rate of water pumping and strongly affected the character of shell movements of the adults. Different species of mollusks, their eggs and larvae were affected to different degrees by the same concentrations of the same turbidity-creating substances. Data indicated that bivalves feed most effectively in relatively clear water."

Lowery, G. H., 1974, The mammals of Louisiana and its adjacent waters: Louisiana State University Press, Baton Rouge, 565 p.

LOUISIANA, MAMMALS, NUTRIA

Includes discussion of the habits of nutria.

Mackin, J. G., 1961, Canal dredging and silting in Louisiana bays: The University of Texas, Institute of Marine Science Publications, v. 7, p. 262-314.

LOUISIANA BAYS, CANALS, DREDGING, SILTING, OYSTERS, TURBIDITY, FORAMINIFERA, WINNOWING, EROSION, SUBSIDENCE, CORE, SPOIL

Mackin studied forams from cores and related them to siltation rates and other processes. "Studies of foraminifera in the bottom muds often indicate concentrations far beyond what would be possible if siltation is taking place at a rate of 0.1 foot per year." Mackin also states that winnowing of the fine sediments concentrates shell materials and forams in the surface layers. "Foraminiferan shells were found in all cores taken in Louisiana which were composed of marine muds deposited at a normal

rate of speed. Because most bottoms are in equilibrium, or depositing rates are slow, foraminiferan shells accumulate in the surface layers." Layers that are deposited rapidly do not contain forams, nor do they contain shells of molluscs, woody debris or any other particles too large and heavy to be moved by an ordinary current. "The data indicate that forams may not live to reach mature size if the mean rate of deposit is about four inches per month."

Maurer, Don, 1969, Pelecypod-sediment association in Tomales Bay, California: *The Veliger*, v. 11, no. 3, p.243-249.

PELECYPOD, SEDIMENT, TOMALES BAY, CALIFORNIA, CORRELATION, SIZE, DISTRIBUTION, ABUNDANCE, MUD, SAND, SILT, CLAY, COARSE, FINE, TELLINIDS, BIVALVE

During June, July, and August of 1961 through 1963, 142 samples were taken in Tomales Bay, California. Sediment samples were taken with a van Veen grab and washed with a 1.0mm screen. Five pelecypod species, *Tellina buttoni*, *T. salmonea*, *Mysella tumida*, *Transennella tantilla*, and *Lyonsia californica*, were found, and their distribution and size were correlated with sediment. Results indicate that the distribution and abundance of all species except *Transennella tantilla* appear to be strongly influenced by sediment type. Also, as sediment particle increases, size of the tellinids living in sediment increases. For *Mysella tumida*, size decreases as particle size increases. No explanation was given for the influence of sediment on pelecypod size.

Maurer, D., Watling, L., Kinner, P., Leathem, W., and Wethe, C., 1978, Benthic invertebrate assemblages of Delaware Bay: *Marine Biology*, v. 45, p. 243-249.

BENTHIC, DELAWARE BAY, DENSITY, BIOMASS, ASSEMBLAGES, SEDIMENT

"The benthic invertebrates of Delaware Bay were related to the cosmopolitan mode of estuarine faunas. Faunal assemblages were identified by cluster analysis. The assemblages were associated with sediment type and salinity. It was concluded that Delaware Bay comprises a mosaic of animal assemblages, some of which have relatively sharp boundaries similar to classical level bottom type communities, whereas the boundaries of others are almost impossible to detect, and these represent species distributed along an environmental continuum."

McBee, J. T., and Brehm, W. T., 1982, Spatial and temporal patterns in the macrobenthos of St. Louis Bay, Mississippi: *Gulf Research Reports*, v. 7, no. 2, p. 115-124.

MISSISSIPPI, MACROBENTHOS, TEMPORAL, SPATIAL, COMMUNITY, SEDIMENT, BAY, ESTUARINE, ENDEMIC, OPPORTUNISTS, PREDATION, CLAY, EURYHALINE

"The station-date dendrogram and the Duncan's multiple range test for sand show no consistent pattern between the groupings of stations based on percent sand and the distribution of benthic infauna in

St. Louis Bay. Station groupings in the station-dendrogram, determined mainly by the distribution of deposit feeders, show no relationship to those groups based on percentage of clay in the sediment. Most estuarine organisms have a wide range of habitat selection. Sedimentary effects in estuaries are confounded with other environmental parameters such as depth, salinity, and seasonal changes...."

McCall, P. L., 1977, Community patterns and adaptive strategies of the infaunal benthos of Long Island Sound: *Journal of Marine Research*, v. 35, no. 2, p. 221-266.

LONG ISLAND SOUND, INFAUNAL, BENTHOS, COMMUNITY, OPPORTUNISTS, ECOLOGIC SUCCESSION, DEPOSIT-FEEDERS, SEDENTARY, DEFAUNATED, COLONIZATION, DEPOSIT-FEEDERS, SUSPENSION-FEEDERS, PREDATOR EXCLUSION, HIGH WATER CONTENT

"Local disturbances of the bottom which result in catastrophic mortality are proposed to be the cause of many spatial and temporal faunal changes encountered in Long Island Sound. This phenomenon was studied by placing samples of defaunated mud on the bottom of two sites in central Long Island Sound to simulate a local disaster." Opportunistic species are the first to colonize the defaunated areas. Opportunists survive in unpredictable environments possibly by their ability to discover new habitats quickly, to develop rapidly and to reproduce many times per year. Opportunists were *Streblospio*, *Capitella*, *Owenia* and the amphipod *Ampelisca*. These species were in all sediment types but were restricted to water less than 20m. Five of seven suspension feeders found on natural bottom were absent after 57 days of sediment deposition and 10 of the 13 deposit feeders remained. "Secondary successional changes are probably common in nearshore environments such as intertidal, shallow subtidal, and polluted habitats-all environments subject to frequent physical and biological disturbances."

McEwen, M. C., 1969, Sedimentary facies of the modern Trinity delta, in Lankford, R. R., and Rogers, J. J. W., *Holocene geology of the Galveston Bay area*: Houston Geological Society, Chapter 3, p. 53-77.

TRINITY RIVER, DELTA, FACIES, CORES, GALVESTON BAY, SEDIMENTS, GEOLOGY

Following facies identified from cores taken in the Trinity River delta and Trinity Bay area: Channel and point bar, natural levee, channel mouth and submarine extension of natural levee, blocked channel, marsh, lake, interdistributary bay, delta front, bay or prodelta, and bay margin. Delta-front churned sands are volumetrically most important unit. These churned sands are thickest at oldest part of delta (NW part)--14-15 ft thick and base is 15-16ft below sea level. In central part of delta this unit is about 8 ft thick and base is 8-10 ft below sea level. On southeast part of delta, coalescing laminated deposits of numerous distributaries are more common. At eastern end the transition of delta-front to prodelta not apparent because of material from erosion of Pleistocene scarp. No evidence that thicker part of delta due to compaction or displacement of underlying sediment. No measurable subsidence has occurred during growth of delta as indicated by thickness and depth of marsh and levee deposit --no core encountered marsh or levee deposit at depths greater than two ft below present sea level (suggests stable sea level)

McKinney, L. D., Bedinger, C. A., and Hopkins, S. H., 1976, The effects of shell dredging and siltation from dredging on organisms associated with oyster reefs, in Bouma, A. H., ed., Shell dredging and its influence on Gulf coast environments: Houston, Texas, Gulf Publishing Company, p. 281-303.

SHELL DREDGING, OYSTER REEF, SAN ANTONIO BAY, MUD FLOW, DISCHARGE, TURBIDITY, SPAT, OYSTER-REEF ASSOCIATES

" In conclusion, our observations and experimental results from the dredging study agreed with all other studies of the discharge from operating dredges and of the effects of such discharges on oysters and their associated fauna. All studies concurred on two major points:(1)high-density mud flow affected only those organisms on the bottom, level with the discharge;(2)at levels more than two feet above the bottom at point of discharge, turbidities resulting from dredging were not greater than those which might occur under natural conditions for comparable length of time and these turbidities did not kill oysters or the common oyster reef animals, which nature has enabled to survive such adversities. In addition, we found that:(3) baskets of oysters and oysters on reef flank areas low enough and close enough to the dredge to be killed by burial in mud were soon repopulated by reef and mud bottom animals and that (4) the mud did not remain on the reef indefinitely but was gradually washed away by waves and currents so that shells were again available for attachment of oyster spat and other sessile animals..."

Meade, R. H., 1969, Landward transport of bottom sediments in estuaries of the Atlantic coastal plain: Journal of Sedimentary Petrology, v. 39, no. 1, p. 222-234.

ESTUARY, SEDIMENT, ATLANTIC COASTAL PLAIN, LANDWARD TRANSPORT, MINERALS, SAND, WATER CIRCULATION, RIVERS, FRESH-WATER INFLOW, SUSPENDED MATTER

Estuaries of the Atlantic Coastal Plain are being filled with sediment. River-borne sediment is partially trapped in the estuaries by the predominantly landward flow of estuarine bottom waters. Measurements during tidal cycles indicates sediment is being moved landward from shelf. However, field data is based on average day-to-day conditions and not extreme conditions when river flooding carries largest load; during these times sediment may be transported through estuary and out to sea. Influence of man has increased river-borne sediment; conversions of forest to farms has caused at least a tenfold increase in sediment yield of the land (Wark and Keller, 1963). Urbanization has added enormous loads of sediment to rivers (Wolman and Schick, 1967). Even with reservoirs intercepting part of the sediment load from major rivers, present loads that enter the Atlantic Coast estuaries are greater than they would be if the drainage areas were still in their natural state.

Meade, R. H., and Emery, K. O., 1971, Sea level as affected by river runoff, eastern United States: Science, v. 173, p. 425-428.

SEA LEVEL, RIVER FLOW

From Coastal-estuarine and nearshore processes, an annotated bibliography: variations in annual river inflow account for 7 to 21 percent of the total variation in average annual sea level along the Atlantic and Gulf of Mexico coasts. This compares with 29 to 68% of the total variation that can be attributed to the secular rise of sea level and with 10 to 50% of the variation that cannot be attributed either to river inflow or the secular rise.

Morgan, J. P., 1970b, Depositional processes and products in the deltaic environment, in Morgan, J. P., and Shover, R. H., eds., Deltaic sedimentation, modern and ancient: Society of Economic Paleontologists and Mineralogists, Special Publication 15, p. 31-47.

DEPOSITION, PROCESSES, DELTAS, MISSISSIPPI RIVER, MEKONG RIVER

Factors controlling and influencing delta formation: (1) river regime -- principally particle size and amount of material transported by the river to the delta as well as seasonal variations in these properties due to fluctuations in flow, (2) coastal processes -- the influence of waves, tides, and currents on the margins of the deltaic environment, (3) structural behavior and the relation of sea level to the depositional site -- is the delta prograding across a stable platform, or is subsidence occurring contemporaneously with the deposition, and is sea level rising, falling, or stationary, (4) climatic factors -- especially those that affect delta vegetation.

Morton, J. W., 1977, Ecological effects of dredging and dredge spoil disposal: a literature review: Washington, D.C., Department of the Interior, Fish and Wildlife Service, Technical paper, 33 p.

DREDGING, SPOIL, LITERATURE REVIEW, ANIMAL-SEDIMENT RELATIONSHIPS, SEDIMENT, FILTER FEEDERS, COMMUNITY, INDICATOR SPECIES, DISPOSAL

This report includes a good discussion on biological effects of dredging and spoil disposal on estuarine environments, including a section on benthos and gross effects, effects of suspended sediment on filter feeders, community response, animal-sediment relationships, and indicator species.

Nichols, F. H., 1977, Infaunal biomass and production on a mudflat, San Francisco Bay, California, in Coull, B. C., ed., Ecology of marine benthos: Columbia, South Carolina, University of South Carolina Press, p.339-357.

INFAUNA, BIOMASS, MUDFLAT, SAN FRANCISCO BAY, CALIFORNIA, ORGANICS, MARSH, STANDING CROP, BIRDS

From page 354- "Our observations of a large standing crop of infaunal invertebrates throughout the year indicate that the mudflat acts as a sink for much of the organic matter produced in the marsh, in the water column, and on the mudflat itself; it also provides abundant food for resident and migratory

birds. As such, the mudflat invertebrate community is an important link in the cycling of organic matter of San Francisco Bay estuary." In the mudflats of San Francisco Bay, molluscs were the chief primary consumers. Bird predation is big factor in the early mortality of many of the invertebrates.

Nichols, J., and Rowe, G., 1977, Infaunal macrobenthos off Cap Blanc, Spanish Sahara: *Journal of Marine Research*, v. 35, p. 525-536.

INFAUNA, MACROBENTHOS, CAP BLANC, SPANISH SAHARA, SEDIMENTATION, BIOMASS, BENTHIC, UPWELLING, ORGANIC MATTER, PERCENT SIMILARITY, ORGANIC CARBON, BENTHIC PRODUCTION, PRIMARY PRODUCTIVITY

"Cap Blanc, Spanish Sahara is characterized by a nearshore benthic faunal province physically controlled by rates of sedimentation, conservatively estimated at 4cm/yr." Sedimentation rates estimated with sediment traps suspended at various heights off bottom. For comparison, figures for sedimentation rates in Buzzard Bay were given as an average rate of 4.55mm/yr. "Sedimentation rates at Cap Blanc are less than those shown experimentally in Buzzards Bay to prohibit escape from burial (Nichols et al., in prep.). However, the constant expenditure of energy to escape burial could explain the observed small size of individuals nearshore. It has been suggested elsewhere that small size is an advantage when organisms are buried in fine sediments (Thayer, 1975). *Spio* sp. A dominant offshore was not eliminated from the nearshore community by burial, but the competitive advantages were shifted to species better able to cope with burial, sediment instability and high turbidity."

Nichols, M. M., 1964, Characteristics of sedimentary environments in Moriches Bay: *Papers in Marine Geology*, Shepard commemorative volume, p. 363-383.

TEXAS, BAYS, SEDIMENTS, SEDIMENT SOURCES

Compares Moriches Bay with Texas bays

Nichols, M. M., and Ellison, R. L., 1967, Sedimentary patterns of microfauna in a coastal plain estuary, in Lauff, G. H., ed., *Estuaries*: Washington, D.C., American Association for the Advancement of Science, Publication No. 83, p. 283-287.

VIRGINIA, RAPPAHANNOCK ESTUARY, FORAMS, SEDIMENTATION RATES, MARSHES

Discussion of forams; brief discussion of sedimentation rates based on ratio of live/dead specimens. A method used by Phleger (1960). Author found little relationship between sedimentation rates estimated from successive changes of depth on hydrographic charts and foram ratios. Has information of marsh forams in addition to estuarine forams.

Odum, E. P., 1971, Fundamentals of ecology: Philadelphia, W. B. Saunders Company, 574 p.

ECOLOGY, BIOMASS, STANDING CROP, BENTHOS

This is a textbook on ecology.

Odum, H. T., 1963, Productivity measurements in Texas turtle grass and the effects of dredging an intracoastal channel: Publications of the Institute of Marine Science, v. 9, p.48-58.

THALASSIA TESTUDINUM, TURTLEGRASS, DREDGING, TURBIDITY, PRODUCTIVITY, PHOTOSYNTHESIS, REDFISH BAY, SILT

"Data reported here indicate a marked seasonal cycle in productivity with considerable year to year variation possible. The values for the Redfish Bay Turtle Grass were not as high as some grass flat data in the lower Laguna Madre of Texas where higher values were recorded associated with clearer waters. The decreased productivity and imbalance of respiration over photosynthesis of the summer of 1959 may have been associated with the silts turned loose over the flats by the start of dredging that spring. Certainly, during the weeks of dredging nearby, light penetration was much reduced and the plant beds may have received a set-back at this time."

Odum, H. T., and Wilson, R. F., 1962, Further studies on reaeration and metabolism of Texas Bays, 1958-1960: The University of Texas, Institute of Marine Science Publications, v. 8, p. 23-55.

PHOTOSYNTHESIS, TURBIDITY, MARINE GRASS, TEXAS BAYS, METABOLISM

From summary section--"Maximum efficiencies, productivities, and sustained respiration were found in natural, high salinity, grass flat systems with values equal to those in artificial sewage ponds. Values were as great as in tropical reefs with photosynthetic efficiencies as high as other aquatic communities of maximum fertility on earth. To produce maximum total photosynthesis in all of the waters of Texas, measures for management should include: reducing turbidity, eliminating irregular flushing of flood waters, developing grass bottoms, retaining wind driven circulation, and adjusting water depths of shallow and deep areas toward an average depth of 0.5m."

Officer, C. B., Lynch, D. R., Setlock, G. H., and Helz, G. R., 1984, Recent sedimentation rates in Chesapeake Bay, in Kennedy, V. S., ed., The estuary as a filter: Orlando, Academic Press, p. 131-157.

ESTUARY, SEDIMENTATION RATES, BIOTURBATION, Pb, Cs, CHESAPEAKE BAY, SEDIMENT SOURCES, SUBSIDENCE, SEA LEVEL, BATHYMETRIC CHANGES, RIVER

Good discussion of method used to determine sedimentation rates which takes into account bioturbation and compaction among other variables. Bay sedimentation rates were determined

using nonlinear parameter estimation techniques from Pb-210, Cs-137, and Pu-239,240, geochemical tracer profiles. The upper bay, with the Susquehanna River as the principal sediment source, had the highest rates at 0.3-1.2 gm/sq-cm/yr; the middle bay had modest rates at 0.1-0.3 gm/sq-cm/yr; and the lower bay, with principal source the ocean and lower bay environments, had modest to high rates of 0.1-0.8 gm/sq-cm/yr. With regard to total sediment budget, results agree with published evidence that episodic sedimentation events are important. Chesapeake Bay traps essentially all the river-derived sediments and is also a sink for ocean-derived sediments.

Orth, R. J., 1977, The importance of sediment stability in seagrass communities, in Coull, B. C., ed., Ecology of marine benthos: Columbia, South Carolina, University of South Carolina Press, p. 281-300.

SEDIMENT STABILITY, DIVERSITY, SEAGRASS, INFAUNA, SAND, LARVAE, LEAF BLADES, RHIZOMES, BAFFLING EFFECT

Sandy environments are generally unstable, and this accounts for the lack of species and individuals in this habitat. Marine grasses increase stability and increase species number and density of individuals. Once larvae have settled in the grass bed, neither physical or biological processes can remove juveniles and adults from the habitat. "Leaf blades can increase the complexity of the substrate, and may therefore be important to sessile species requiring an attachment site for initial settlement." Other quotes--"In many exposed intertidal and shallow subtidal environments, sediments resuspended and transported by wave energy and tidal currents are the dominant forces preventing successful recruitment of many benthic forms. Associated with these habitats is a fauna characterized by low species diversity and density. Survival in this environment requires a species to be either a rapid burrower or a tube builder....Seagrasses are well known for their ability to stabilize sediments by baffling currents and damping wave action."

Parker, R. H., 1975, The study of benthic communities, a model and a review: Elsevier Scientific Publishing Company, New York, 279 p.

BENTHOS, INFAUNAL, SEDIMENT, HADLEY HARBOR, MASSACHUSETTS, TEXAS, FAUNAL COUNTS, CLAY, SILT, SAND, GRASS

From p. 241-"Faunal counts were made from a series of samples taken in Texas coastal habitats and compared with counts taken from Hadley Harbor samples. Highest counts for all stations in Texas and Massachusetts habitats were found in the grass flats of Cape Cod, averaging 110,000 animals taken from grass flats within south Texas bays. Lowest counts characterized samples from the highly unstable river and bayou estuaries and the low-salinity secondary and tertiary bays of the upper T and tertiary bayey sediments produced the lowest number of organisms in both the Cape Cod and Texas studies, while mixed sediments characterized areas of high benthic standing crops in all areas samples." Fig. 92 shows counts for sand clay and sandy clay for bays on Texas coast and for Massachusetts.

Persson, L., 1983, Temporal and spatial variation in coastal macrobenthic community structure, Hano Bay (southern Baltic): *Journal of Experimental Marine Biology and Ecology*, v. 68, p. 277-293.

BALTIC, TEMPORAL, SPATIAL, MACROBENTHOS, BIOMASS, SEDIMENT, DIVERSITY, SIMILARITY, INSTABILITY

Lowest biomass at 5m depth station. All three stations in fine sand. Highest biomass at 15m station. Highest density at shallow station. "The low biomass found at the 5-m depth indicates more unstable conditions which do not allow the long-lived molluscs to develop a normal population structure with a high number of old individuals. This contrasts with the theory presented by Jackson (1972), who postulated large infaunal bivalves to be common in intertidal and shallow subtidal sediments due to their ability to burrow deeper and escape the effects of sediment instability. The reason might be that the Baltic species of bivalves are too slow-growing to be able to reach a large size between the recurring storms causing catastrophies at the 5-m depth."

Peterson, C. H., and Peterson, N. M., 1979, The ecology of intertidal flats of North Carolina: a community profile: Washington, D.C., U.S. Department of the Interior, Fish and Wildlife Service, Office of Biological Services, FWS/OBS-79/39, 73 p.

NORTH CAROLINA, INTERTIDAL FLAT, SEDIMENT, INFAUNA, VEGETATED, UNVEGETATED, PREDATION

This book gives a good general discussion on the ecology of the intertidal flat environment, including faunal-sediment relationships. "Infauna of intertidal flats in North Carolina follow a very general pattern found broadly among the benthic infauna of shallow bottoms. The infauna on unvegetated flats is far less abundant than the infauna of marine grass beds at the same tidal height." The reasons for this may be that the infauna of grass beds get more food because as the grass blades baffle and slow the passing currents, the rate of organic deposition increases. Also, densities may be greater because the grass provides a qualitatively different and new habitat which permits the addition of new species. Finally, the last explanation for increased infauna abundance in grasses may be that predator density is less than in unvegetated environments. Other sections of book provide useful look at sediment-faunal relationships and the infauna of the intertidal environment.

Poff, M. J., 1973, Species composition, distribution and abundance of macrobenthic organisms in the intake and discharge areas after construction and operation of the Cedar Bayou Electric Power Station: Texas A&M University, Master's thesis, 300 p.

RUPPIA MARITIMA, WIDGEON GRASS, TRINITY BAY, CEDAR BAYOU

Poff made observations on the distribution of *Ruppia* in Trinity Bay around the discharge area of the Cedar Bayou electrical generating plant. Observations of grass beds were made from February 1970 through May 1972. *Ruppia* was limited to the shoreline discharge areas and extended to

approximately 300m offshore. *Ruppia* sprouted in February 1970 and grew to height of 53cm by August but vanished by November 1970. In 1971 *Ruppia* was neither as long or as thick as in 1970. By early October 1971, the beds were very sparse and vanished completely by early November. No growth of *Ruppia* was observed during the remainder of the study (through May 1972).

Pratt, D. M., 1953, Abundance and growth of *Venus mercenaria* and *Callocardia morrhuana* in relation to the character of bottom sediments: *Journal of Marine Research*, v. 12, no. 1, p. 60-74.

LARVAE, BIVALVE, SETTING, CURRENT, SPAT

Larvae of *Venus mercenaria* survive best when current velocities are low--"presumably a low current velocity favors setting, but it probably also results in three conditions adverse to survival:(1)rapid deposition of fine silt which may bury or smother the young set, (2)low rate of supply of food and oxygen, and (3)slow removal of metabolic wastes. On the other hand, a high current velocity probably allows only limited setting of the larvae and may cause a grinding action of coarse particles severe enough to injure the spat, but undoubtedly it favors survival by high rates of food and oxygen supply and waste removal."

Purdy, E. G., 1964, Sediments as substrates, in Imbrie, J., and Newell, N. D., eds., *Approaches to paleoecology*: New York, John Wiley, p.238-271.

SEDIMENT, SUBSTRATE, LITHOLOGY, DEPOSIT FEEDER, SUSPENSION FEEDER, VAGILE, ORGANIC CONTENT, SILT, CLAY, INFAUNA, SUBSTRATE MOBILITY, TOXIC DECOMPOSITION PRODUCTS, SESSILE, INTERSTITIAL

This is fairly early and basic discussion of relationship between sediment, feeding type, and organic content of substrate. In shifting sands the fauna is characterized by low population density and few micro-and macrofauna. The species present consist largely of suspension feeders, the lack of deposit feeders because of the small amount of organics in the substrate. Reduced current velocities favor the accumulation of mud and allow the deposition of organic detritus, and the proportion of deposit feeders increases. Suspension feeders will remain the dominant feeding type in sandy sediment and "suspension feeders present are sessile as well as vagile." With increase in finer sediments the proportion of suspension feeders decreases and the abundance of deposit feeders increases until finally the deposit feeders are more numerous than the suspension feeders. "In extremely fine grained sediments the fauna is generally characterized by low population density and low taxonomic diversity." This is because of poor interstitial circulation which results in accumulation of toxic waste products or loss of oxygen.

Redfield, A. C., 1972, Development of a New England salt marsh: *Ecological Monographs*, v. 42, no. 2, p. 201-237.

SALT MARSH, SEDIMENTATION, SPARTINA ALTERNIFLORA, SPARTINA PATENS, AGGRADATION RATES, SEA-LEVEL RISE

Redfield studied the development of a salt marsh at Barnstable, Mass., from the seeding of bare sand flats through an intertidal marsh to a mature high marsh. Tidal range in this area is from 9 to 10 ft. The source of sediments, primarily sand, was from the erosion of extensive sea cliffs. The few streams entering the marsh do not carry an appreciable sediment load. The large tidal range produces a broad tidal zone suitable for marsh development. Average rates of marsh accretion listed in a table are 52 mm/yr (initial development), 18 mm/yr, 2.7 mm/yr, 2.1 mm/yr, 1.5 mm/yr (translated from his units given in ft/yr). Differences in rate due to differences in elevation, availability of sediment, and distance from open water. Presumably vertical accretion is rapid on margins where levees form, but as margins get higher rate of accretion decreases and interior of marsh catches up so the ultimate effect is that the entire surface becomes level. Development of a salt marsh depends on: (1) physiology of the local halophytic vegetation, (2) behavior of the tide (3) processes of sedimentation, and (4) changes of sea level relative to land.

Reidenbaugh, T. G., Banta, W. C., Varricchio, M., Streiter, R. P., and Mendoza, S., 1983, Short-term accretional and erosional patterns in a Virginia salt marsh: Gulf Research Reports, vol. 7, no. 3, p. 211-215.

SALT MARSH, VIRGINIA, SPARTINA ALTERNIFLORA, ACCRETION, EROSION, SEDIMENT, TIDAL WRACK, MARSH, TIDAL FLATS

From abstract-"We estimated 3-year rates of accretion and erosion in different vegetation zones of a juvenile *Spartina alterniflora* salt marsh a Wallops Island, Virginia, by precise releveled of a fixed grid. Seaward of the marsh there was extremely variable accretion and erosion in tidal flat, as a result of winter ice scouring and transport. At the lower limit of the marsh, tall *Spartina* edge marsh accreted at about 6.2 mm/yr, well in excess of relative sea level rise, supplied by mineral sediments. At the upper limit, levee *Spartina* and high marsh accreted at about 1.6mm/yr, in equilibrium with sea level rise. Accretion there was supplemented by organic sediments from tidal wrack. At mid elevations, medium *Spartina* middle marsh eroded slightly at about 0.6 mm yr, and low density *Spartina* and bare soil eroded rapidly at about 5.3 mm/yr. These zones may be relatively sediment-starved. The most severe erosion resulted from vegetation diebacks beneath tidal wrack. Patterns of accretion and erosion show that this site is maturing topographically from a juvenile foreshore marsh to a creek-drained marsh."

Reinharz, Eli, and O'Connell, Arthur, 1983 Animal-sediment relationships of the upper and central Chesapeake Bay: Annapolis, Maryland, U.S. Environmental Protection Agency Final Report, EPA Grant No. R8059864, 49p.

SEDIMENT, CHESAPEAKE BAY, BENTHOS, BIOMASS, DENSITY, MUD, SAND, MIXED SEDIMENT, BIOTURBATION, ANOXIA, VERTICAL DISTRIBUTION, OLIGOHALINE, MESOHALINE, BOX CORER, X-RADIOGRAPHIC, PB 210

Fifty-two box cores were taken in upper and central Chesapeake Bay in Fall 1978 and summer 1979. Greatest species populations were in mixed-sediment type. Benthos were concentrated in top 10cm of sediment with penetration to 30cm or more. Bioturbation was substantially reduced in upper

region of bay because of high sedimentation rates. Prevalence of physical structures over biological features in the deep-water channel regime is due to fluid muds and periodic anoxia. Density of individuals and biomass greatest in mixed-sediment habitat. All habitats with exception of oligohaline sands, contain deep-dwelling fauna. Deep penetrating species are most abundant in mixed sediments. Biogenic diversity greatest in stable mud environment. The stable mud environment is primarily inhabited by a sedentary tube and relatively permanent burrow-dwelling community.

Renwick, W. H., and Ashley, G. M., 1984, Sources, storages, and sinks of fine-grained sediments in a fluvial-estuarine system: Geological Society of America Bulletin, v. 95, p. 1343-1348.

FLUVIAL, ESTUARINE, SEDIMENTS, SEDIMENT BUDGET, RARITAN RIVER, NEW JERSEY, SEDIMENTATION RATES.

Among conclusions are that the estuary is a sediment and pollution sink where the rate of accumulation is 1.5 to 3.4 mm/yr. "Sediments collected in the estuary averaged approximately 15% sand, and this figure may be used to adjust the total for bed-load contribution". The sediment budget was calculated. For main stem Raritan River, a discharged weighted estimate of mean annual suspended sediment load is about 140,000 tons/yr. Added loads of tributaries entering below a certain location were determined to be 46% of this tonnage by assuming uniform sediment yield per unit drainage area. Using the bed load figure noted above adjustments were made to determine total bed-load contributions. Altogether, these values suggest a mean annual sediment input from upland sources of approximately 235,000 tons/yr (approx. 196,000 cu m/yr, assuming a bulk density of 1.2 tons/cu m (Marshall and Holmes, 1979). The authors determined the annual total sediments transported landward into the estuary to be 88,000 tons, and removed by maintenance dredging, 240,000 tons. Marsh area probably receiving sediments = 10 km, at 3 mm/yr yields 36,000ton/yr.

Rhoads, D. C., 1974, Organism-sediment relations on the muddy sea floor: Oceanography and marine biology annual review, vol. 12, p. 263-300.

SEDIMENT, SUBSTRATE, SUSPENSION FEEDING, DEPOSIT FEEDING, FAECAL PELLETS, BIOTURBATED, FEEDING TYPE, BOTTOM STABILITY, BIODEPOSITION, SILT, CLAY, DIVERSITY

Contains good discussion of biologic modifications of the bottom. "It is not surprising to find that the surface of the muddy sea floor is passed through the benthos at least once, and in some cases, several times each year...At higher sedimentation rates, the influence of benthos on the sedimentary particles decreases. Moore and Scruton (1957) found that the effect of burrowing organisms on sedimentary properties decreased substantially when sedimentation rates exceeded about 12 cm/yr on the Mississippi pro-delta."

Rhoads, D. C., McCall, P. L., and Yingst, J. Y., 1978, Disturbance and production on the estuarine seafloor: American Scientist, vol.66, p. 577-586.

DREDGE, LONG ISLAND SOUND, SEDIMENT, ESTUARY, COLONIZERS, OPPORTUNISTS, PREDATION, BENTHOS, DISTURBANCE, DREDGE-SPOIL

Although article is about dredging and benthos, could be applied to any benthic community where either a disturbed or an equilibrium community is present and how disturbance affects benthic productivity. Excerpts from article follow: "Three patterns of colonization of shallow-water marine sediments by invertebrate species can be identified. Group 1 colonizers which arrive within a few days, reach peak abundances within a relatively short period of time. In the absence of further disturbance, Group 1 species experience high mortality and may disappear locally as a result of competition and/or predation. Group 3 colonizers may appear early in succession, but they maintain more or less constant, relatively low population densities. Individuals of these species persist over long periods of time. Group 2 species represent a mode of colonization intermediate to groups 1 and 3. Benthic assemblage on disturbed bottoms are dominated by species with short life spans, frequent reproduction and high dispersability." Other important sections in article highlighted in photocopy.

Rhoads, D. C., and Young, D. K., 1970, The influence of deposit-feeding organisms on sediment stability and community trophic structure: *Journal of Marine Research*, v.28, no.2, p.150-178.

SEDIMENT, STABILITY, COMMUNITY, TROPHIC, DEPOSIT-FEEDING, SUSPENSION-FEEDING, INFAUNA, BENTHIC, MUDS, SANDS, FECAL MATTER, TROPHIC-GROUP AMENSALISM, BUZZARDS BAY, MASSACHUSETTS

Deposit feeders modify fine-grained sediments in the following ways: (1) high turnover rate of bottom muds by a process of resuspension that is driven by weak tidal currents (2) high turbidity at the sediment-water interface (3) "production of textural and compositional grading in reworked sediments corresponding to the maximal depth of biologic reworking". Sediment instability is stressful to suspension feeding benthos by clogging filtering mechanisms, resuspending and burying newly settled larvae, and discouraging the settlement of suspension-feeding larvae. This process (the exclusion of one trophic group in an area occupied by another) is called trophic-group amensalism. In Buzzards Bay, three trophic groupings occur (1) homogeneous suspension feeding--inadequate food for deposit feeders (2) homogeneous deposit feeders--trophic-group amensalism (3) Mixed trophic group--physical stability of bottom allows diverse suspension-feeder population on mud bottom. A reworked mud surface limits suspension feeders only when surface becomes mobile. Sediment instability and turbidity limiting to suspension feeders.

Rice, R. H., 1967, A study of the suspended load of the Trinity River, Texas: Rice University, Houston, Texas, Masters Thesis, 47 p.

TRINITY RIVER, SUSPENDED LOAD, TEXAS

Abstract-"The Trinity River drains approximately 17,500 square miles of north-central and eastern Texas. Its average annual discharge is 5,166,000 acre-feet and in water year 1965 it carried

5,308,000 acre-feet of water. The river's average annual suspended load contribution to its delta is roughly 3000 acre-feet. During 1965 over 50% of the Trinity River's discharge and over 80% of its suspended load were transported by two major floods. The suspended material delivered to the delta contains about 25% sand, 30% silt, and 45% clay." Author notes that because results are based on such limited data they should be viewed as qualitative rather than quantitative.

Rice, R. H., 1969, Suspended load of the Trinity River, chapter 4, in Lankford, R. R., and Rogers, John J. W., Holocene geology of the Galveston Bay area: Houston Geological Society, p. 78-84.

TRINITY RIVER, SUSPENDED LOAD, DELTA, MASS BALANCE

Distance of Trinity River from headwaters to mouth is roughly 400 miles. It drains an area of approximately 17,500 sq. miles -- underlain by sedimentary rocks. Ave. annual rainfall is about 27 in. at western edge of headwaters and 50 in. near delta. Maximum flow recorded during 41 yr period from 1924-1965 was 12.3 mill. acre ft (1940-41); minimum flow 0.661 mill acre ft (1924-25); yearly ave. 5.166 mill acre ft. Reasonable estimate of grain-size distribution of sediments delivered to the delta each yr. is 25% sand, 30% silt, and 45% clay. Calculations indicate that the river delivers about 3000 acre ft. of suspended sediment to the delta each year; yearly average estimated in "The Silt Load of Texas Streams" (1955) was 3662 acre ft. Based on investigation of floods during 1965, author estimated that about 82% of the annual suspended load was carried by floods. Size of delta-6 sq. mi., thickness 15 ft. which equals 57,600 acre-ft of sediment. Assuming that 75 % of delta is sand author found imbalance. Sand delivered by river over 500 yrs = 375,000 ac-ft but contains max of 43,200. Concludes aggradation of lower valley.

Rosen, P. S., 1977, Increasing shoreline erosion rates with decreasing tidal range in the Virginia Chesapeake Bay: Chesapeake Science, vol, 18, no. 4, p. 383-386.

SHORELINE EROSION, TIDES, CHESAPEAKE BAY, TIDAL RANGE

Abstract-"The Virginia Chesapeake Bay has a large variation (200%) in tidal range (.36 m to 1 m) over a relatively short (120 km) distance. Analysis of a regional sampling of shoreline morphology and processes demonstrates that decreasing tidal ranges result in increasing long-term erosion rates. This results because larger tidal range areas form a higher elevation beach. Thus, in larger tidal range areas, water levels increased by storm surges have a lower probability of reaching the elevation of the fastland (bluff or dune) material to augment erosion. Secondly, the wave energy is distributed over a greater distance in the course of a tidal cycle."

Rowe, G. T. and Smith, K. L., Jr., 1977, Benthic pelagic coupling in the mid-Atlantic bight, in Coull, B. C., ed., Ecology of marine benthos: Columbia, South Carolina, University of South Carolina Press, p.55-65.

NUTRIENTS, BENTHOS, SEDIMENT, AMMONIA, REGENERATION, WATER COLUMN, BACTERIA, PELAGIC,

"The purpose of this paper is to explore the hypothesis that benthic nutrient regeneration couples the bottom with processes in the water column."

Rusnak, G. A., 1967, Rates of sediment accumulation in modern estuaries, in Lauff, G. H., ed., Estuaries: American Association for the Advancement of Science, Publication No. 83, p. 180-184.

ESTUARIES, SEDIMENTATION RATES, SEA LEVEL RISE

Positive-filled estuaries--those filled entirely by river-transported sediment inflow. Inverse-filled estuaries--those filled solely or to the largest extent by sediments entering from the sea. Neutral-filled (non-filling) basins--those in which sediments in estuary are being redistributed, say from margins to basin. Rate of sediment accumulation in a basin dependent on source of supply, geographic position, and climate. Rate of accumulation related to sea level rise, which has declined over last 15,000 yrs. Considering the past 6,000 yrs, for humid regions, estimated rates of accumulation vary from high of 300m/1000 yrs for prograding Mississippi River (Shepard, 1953, Scruton, 1960) to low of 2 to 4m/1000 yrs for bays and estuaries (Shepard 1953, Shepard and Moore, 1960). For semiarid lagoons, rate is only 1m/1000 yrs (Rusnak, 1960). These are rates averaged over entire basin; inlets which may cut and fill at extremely rapid rates are excluded.

Sanders, H. L., 1958, Benthic studies in Buzzards Bay. I. Animal - sediment relationships: Limnology and Oceanography, vol. 3, no. 3, p. 245-258.

BUZZARDS BAY, MASSACHUSETTS, SEDIMENT, DEPOSIT FEEDERS, SUSPENSION FEEDERS, SILT, CLAY, SAND, INFAUNAL, BENTHOS

Abstract-"The distribution of certain dominant deposit-feeders in Long Island Sound and Buzzards Bay was poorly correlated with the silt clay fraction of the sediment. However, when clay alone was used, a much better agreement was obtained. It was suggested that clay is probably the most valid criterion for the distribution of deposit feeders. The distribution of infaunal filter feeders seemed related to the degree of sorting and the median grain size of the sediment, with largest populations present in well-sorted fine sand. The hydrodynamic implications of this distribution are discussed." Sediments with large amount of silt-clay reflect feeble currents that pass over them and organic matter can thus settle out. There is therefore a smaller amount of organic matter in suspension to supply food for the filter-feeders. Stations with well sorted fine sands indicate intensity of current over bottom must be fairly constant, probably deviating only slightly from a velocity of two centimeters per second. Currents of this intensity can support large populations of filter feeders.

Sanders, H. L., Goudsmit, E. M., Mills, E. L., and Hampson, G. E., 1962, A study of the intertidal fauna of Barnstable Harbor, Massachusetts: Limnology and Oceanography, vol. 7, no. 1, p.63-79.

BARNSTABLE HARBOR, MASSACHUSETTS, INTERTIDAL, DEPOSIT FEEDERS, SUSPENSION FEEDERS, STABILITY, OMNIVORES, CURRENTS, BIOMASS, CHLOROPHYLL, DIATOMS, DINOFLAGELLATES

The deposit feeder is as common in the sand as in the softer sediments. The distribution of the feeding type is better correlated with chlorophyll concentration than with total organic content. The deposit are using as food the abundant supply of benthic diatoms and dinoflagellates on the sediment. "The sediment must be considered as an indicator of the availability of food, and not as a first order factor directly determining the distribution of feeding types."

Schubel, J. R., Carter, H. H., and Wise, W. M., 1979, Shrimping as a source of suspended sediment in Corpus Christi Bay (Texas): Estuaries, vol. 2, no. 3, p.201-203.

SUSPENDED SEDIMENT, CORPUS CHRISTI BAY, SHRIMPING, TEXAS, DREDGING

"The total amount of sediment disturbed each year by shrimping activity in Corpus Christi Bay is 10-100 times greater than that dredged in an average year for channel maintenance. Shrimping is a more consistent activity than dredging and also affects a greater area of the bay than dredging and dredged material disposal. This is not to suggest that the increased levels of suspended sediment resulting from either of these important activities has a deleterious effect on the biota of Corpus Christi Bay."

Schubel, J. R., and Hirschberg, D. J., 1977, Pb210-determined sedimentation rate, and accumulation of metals in sediments at a station in Chesapeake Bay: Chesapeake Science, vol., 18, no. 4, p.379-382.

SEDIMENTATION RATE, CHESAPEAKE BAY, PB210

Abstract-"This paper contains the first published radiometrically determined contemporary sedimentation rate, a rate characteristic of the past few decades, for any part of the Chesapeake Bay estuarine system. Cores collected from a station in the main body of the Chesapeake Bay off Tilghman Island were analyzed for Pb210, organic carbon, nitrogen, and selected metals. A mean sedimentation rate of between 0.09-0.12 cm/yr was inferred from the Pb210 data. The distribution of C and N are typical of many areas of the Bay. The distributions of extractable Ni, Cr, Cu, and Pb increased in the upper 10-15 cm of the core. The increases in metal concentrations over the past century or so may be a result of man's activities."

Schultz, E. A., and Simmons, H. B., 1957, Fresh water-salt water density currents, a major cause of siltation in estuaries: U. S. Army Corps of Engineers, Technical bulletin no. 2, 28 p.

SILTATION, DENSITY CURRENTS, FRESH WATER, SALT WATER, ESTUARIES, SEDIMENT, HYDRAULICS, UPLAND DISCHARGE

From general conclusions, p. 24--"The degree of mixing of salt and fresh water in estuaries plays an important role in establishing their hydraulic regimens. Because of incomplete mixing of salt and fresh water in estuaries, the predominance of flow in the bottom strata is almost always upstream while that in the surface strata is downstream; the degree of such predominance is dependent on the degree of mixing, being most prominent in the highly stratified estuary and least prominent in the well-mixed type. Changes in the upland discharge, tidal prism, and physical configurations of estuaries will frequently change the degree of mixing of salt and fresh water therein and thus affect such important features as the vertical distribution of current velocities, the direction and degree of flow predominance, the amount of shoaling, and the location of major shoal areas. Available data indicate that lightweight sediments are supplied to estuaries principally through the medium of upland discharge."

Scudato, R. J., Henningsen, B. F., and Estes, E. L., 1976, Trace metal concentrations of selected macrofauna from a southeast Texas estuary: The Texas Journal of Science, vol. 27, no. 4, p. 419-426.

TRACE METALS, OYSTERS, MACROFAUNA, SABINE LAKE, SOUTHEAST TEXAS, CRAB, SHRIMP, CADMIUM, CHROMIUM, COPPER, LEAD, ZINC

From abstract--"Cadmium, chromium, copper, lead and zinc concentrations of *Crassostrea virginica*, *Callinectes sapidus*, and *Peneus setiferus* from a shallow water, southeast Texas estuary (Sabine Lake) indicates levels significantly above associated estuarine waters...."

Shepard, F. P., 1953, Sedimentation rates in Texas estuaries and lagoons: American Association of Petroleum Geology Bulletin, v. 37, no. 8, p. 1919-1934.

ESTUARIES, SEDIMENTATION RATES, TEXAS

The shoaling rates of Texas bays and lagoons were estimated by comparing 20,518 points on bathymetric surveys conducted in the last century with recent surveys. The estimated rate of bay shoaling derived from these comparisons is 1.26 ft/century (3.8 mm/yr). Only the Trinity and Colorado Rivers of Texas have appreciably extended their deltas since the time of the old surveys. The silt load of Texas streams, which is probably deposited in the bays, is too high for the estimated rate of shoaling. This indicates that submergence has likely occurred. Using a submergence rate of 1.8 ft/century (5.4 mm/yr) based on average relative of sea-level rise per century for four Gulf coast tide stations) the total estimated rate of sedimentation in the bays is 3 ft/100yrs (9 mm/yr). Because the deposited material has such a high water content, it yields only about half as much sedimentary rock depending on compaction.

Shepard, F. P., and Moore, D. G., 1960, Bays of central Texas coast, in Shepard, F. P., Phleger, F. B., and van Andel, T. H., eds., Recent sediments, northwest Gulf of Mexico, Project 51, American Petroleum Institute: American Association of Petroleum Geologist, Tulsa, Oklahoma, p. 117-152.

BAYS, SEDIMENTATION RATES, TEXAS, TRINITY DELTA

Figure 3 on page 120 of this article shows depth changes in Texas bays, estimates of stream silt transport per 100 yrs, and solid fill in bays. Averages for each bay indicates that most parts of the bays have shoaled at rates of 1 ft/100yrs (3 mm/yr). Near large rivers the bays have shoaled at a faster rate, especially where deltas have grown forward and filled the bays. The lower bays and bays not connected to rivers have had quite variable changes. The Trinity delta has built forward about 0.3 mi since 1855 (p. 122).

Sherk, J. A., Jr., 1971, The effects of suspended and deposited sediments on estuarine organisms: literature summary and research needs: Prepared for U.S. Army Corps of Engineers, Natural Resources Institute of the University of Maryland, Contribution no. 443, contract no. DACW73-70-C-0013, 73p.

SEDIMENT, SUSPENDED SEDIMENT, BENTHOS, EUPHOTIC ZONE, HABITAT LOSS, PRIMARY PRODUCTION, COMMUNITY, OYSTERS, GRASSFLATS, NUTRIENTS

Good review of literature on effects of increased amount of sediment on estuarine organisms. Includes discussions on loss of habitat, decrease in euphotic zone, oxygen demand, nutrient sorption and release, primary production, community disruption, mortality, and effects of sediment on filter-feeding organisms.

Shideler, G. L., 1980, Reconnaissance observations of some factors influencing the turbidity structure of a restricted estuary: Corpus Christi Bay, Texas: Texas Journal of Science, vol. 32, no. 1, p. 59-71.

CORPUS CHRISTI BAY, TEXAS, NUECES BAY, NUECES RIVER, SUSPENDED SEDIMENT, TURBIDITY, TIDES, FLUVIAL, SEDIMENT, WIND

"Turbidity characteristics along the longitudinal trend of Corpus Christi Bay were highly variable in time and space. Turbidity toward the bayhead sector appeared to be largely influenced by wind, whereas the rate of fluvial-sediment influx from the Nueces River had no discernible influence on bay turbidity. A working hypothesis is suggested whereby sediment influx from the Nueces River enters adjacent Nueces Bay, which appears to function as a shallow storage basin that entraps sediment at times when winds have southerly or southeasterly onshore components. In essence, the influx of fluvial sediments into the estuarine system could be regulated mainly by wind direction. Winds further influence turbidity structures by generating waves that resuspend bottom sediments, especially within the bayhead sector."

Simons, D. B., and Simons, R. K., 1987, Differences between gravel- and sand-bed Rivers, in Thorne, C. R., Bathurst, J. C., and Hey, R. D., eds., Sediment transport in gravel-bed rivers: John Wiley Sons Ltd, New York, p. 3-11.

SEDIMENT, TRANSPORT, BED LOAD, SAND, GRAVEL, RIVERS

For sand-bed rivers, a very large number of variables affect sediment transport, however, most equations utilize a limited number of variables. A further constraint is that most of these theories have been developed using straight, small streams under steady flow. Hence, equations for sediment transport suffer severe limitations considering the complexities of the actual sediment transport process. For sand-bed channels, common equations ignore effects of non-uniform flow, unsteady flow, non-uniform velocities, wash load, river alignment and river dynamics, seepage forces, vegetation, etc. However, in the simplest form, it can be concluded that sediment in sand-bed channels varies as a function of the velocity to approx. the fifth power. Gravel-bed less responsive than sand-bed river because sand-bed material can be transported when flow velocities are of the order of 0.3 m/s or more. Therefore, there is virtually always transport with changes in position of bars and bed forms, etc.

Solomon, D. E., and Smith, G. D., 1973, Seasonal assessment of the relationship between the discharge of the Trinity River and the Trinity Bay ecosystem: Final report, U. S. Army Corps of Engineers, contract no. DACW 63-73-C-0059, prepared by Coastal Ecosystems Management, Inc., 147p.

TRINITY BAY, TRINITY RIVER, FRESH WATER, TIDAL INUNDATION, MARSH, SEDIMENT, BENTHOS, SAND, MUD, STANDING CROP, COARSE SEDIMENTS, FINE SEDIMENTS, BACTERIA, SILT, CLAY, INFAUNA

From the conclusions on p.130-134-"Bacterial counts in the sediments were three orders of magnitude higher than those from the water column. Populations this large possess a biomass large enough to affect the entire ecosystem. High populations and biomass of bacteria, correlated with high total organic carbon plus large amounts of organic matter flushed into the bay, support the thesis that bacteria may form the base of primary production in Trinity Bay. The standing crop of marsh vegetation was not related to river flow except when flooding prevented initial growth. Benthic invertebrates populations did not vary with the river flow. Large numbers thic invertebrates always occurred in samples with large amounts of detritus present. The diversity index of benthic communities was very stable throughout the entire study, in spite of fluctuations of all environmental parameters. Sediments showed a decrease in percentage of clay and an increase in percentage of sand, while the silt fraction remained stable. These sedimentary composition changes were observed to be directly proportional to river flow."

Sorenson, R. P., 1975, Clay mineralogy of suspended-load transport, Guadalupe River, Texas: The University of Texas at Austin, Masters Thesis, 66 p.

**GUADALUPE RIVER, SUSPENDED LOAD, CLAY MINERALOGY, SAN ANTONIO RIVER
GUADALUPE RIVER, ORGANICS**

The San Antonio River, compared to the Guadalupe River, carries a higher suspended load; in addition, there is a higher proportion of inorganics relative to organics in the suspended load. More organics are carried by the Guadalupe. Organic particles make up a large proportion of the suspended load

during normal flow. Amount of inorganic material in transport increases rapidly after storms but the organics in suspension increase only a small amount or even decrease. The high organic content of suspended material during normal flow suggests that measurement of total suspended solids may drastically overestimate the rate of sediment accumulation in a delta because the organic debris is unlikely to be preserved in substantial quantities. Changes in contributions of the two rivers could effect the organic content in the delta. Author suggests that focculation of organics and clay may occur by means of large organic molecules. Significance of this deposition process uncertain but it could be important because of the much greater settling velocity imparted to the clays as a result of attachment to the organic particles.

Stevenson, J. C., Kearney, M. S., and Pendleton, E. C., 1985 , Sedimentation and erosion in a Chesapeake Bay brackish marsh system: *Marine Geology*, v. 67, no. 3/4, p. 213-235.

BRACKISH MARSH, EROSION, SEDIMENTATION, CHESAPEAKE BAY, MARYLAND

Characteristics of sediment transport and deposition were studied in a marsh that has experienced extensive erosion (2600 ha) over last 4 decades. The marsh system is dominated by ebb tides which contribute to a net output of 720,000 MT of sediment/ yr. (13.8 kg/msq/yr). Little inorganic sediment reaches interior marsh because of export. Vertical accretion is primarily by peat deposition. Accretion rates range between 1.7 and 3.6 mm/yr (based on 210 Pb) in active area of marsh loss. The marsh has not kept pace with sea-level rise since 1940 (3.9 mm/yr). Heavy grazing by muskrats may have contributed to the deterioration of the 0.1 to 0.5 m thick organic mat overlying 1 to 4 m thick unconsolidated organic ooze that composes the marsh. Once ponds form they are enlarged in a NW-SE direction by NW tracking storms. The amplitude of astronomical tides are approximately 0.5 m, and meteorological are up to 2 m.

Stora, G., and Arnoux, A., 1983, Effects of large freshwater diversions on benthos of a Mediterranean lagoon: *Estuaries*, vol.6, no.2, p. 115-125.

ETANG DE BERRE, FRANCE, DIVERSION,BENTHOS,HYDROELECTRIC PLANT, DEGRADATION, EURYHALINE, MACROFAUNA, DURANCE RIVER, MOLLUSCS, MUD, TURBIDITY, SUSPENDED SEDIMENT

The Etang de Berre is the largest French Mediterranean lagoon. Lagoon sampled before and after the commencement of operation of a hydroelectric plant and the resulting diversion of fresh water from the Durance River into the lagoon. After several years the marine benthic communities were replaced by a euryhaline and eurythermal assemblage and large areas of the lagoon were devoid of macrobenthos due to oxygen depletion. Sedimentation periodically caused mass mortalities of the molluscs over the entire lagoon. "When the ratio of freshwater to sediment volume discharged (the inverse of suspended sediment concentration) was lower than 1,500 mortalities of molluscs, at least at the northern stations, were observed. In addition to the direct smothering of molluscs due to rapid covering of fine sediments, high suspended matter concentrations, such as exist when the freshwater to sediment volume is less than 10,000, may prove stressful to filter feeding molluscs and lower their resistance to other environmental factors."

Strand, R. I., and Pemberton, E. L., 1982, Reservoir sedimentation: Technical guideline for Bureau of Reclamation: U. S. Department of the Interior, Bureau of Reclamation, Sedimentation and River Hydraulics Section, Engineering and Research Center, Denver, Colorado, 48 p.

SEDIMENTATION, RESERVOIRS, RIVER CHANNEL DEGRADATION, SEDIMENT TRANSPORT,

Discusses methods of determining sediment inflow, reservoir sediment deposition, and downstream channel effects. Sediment yield is the portion of eroded material that makes it through the drainage network to a downstream measuring or control point. Sediment yield rate factors: (of a watershed) include (1) rainfall amount and intensity, (2) soil type and geologic formation, (3) ground cover, (4) land use, (5) topography, (6) upland erosion (nature of drainage network--density, slope, shape, size, and alignment of channels), (7) runoff, (8) sediment characteristics--grain size, mineralogy, etc., and (9) channel hydraulic characteristics. Sediment transport characteristics of a river are influenced by the first 6 factors but reflect a more direct combination of factors 7, 8, and 9. Systems of weighting these individual factors have been devised (Pacific Southwest Interagency Committee, 1968) to arrive at a sediment yield rate for an individual drainage basin. The report gives a rating chart of factors affecting sediment yield in table 1.

Subrahmanyam, C. B., Kruczynski, W. L., and Drake S. H., 1976, Studies on the animal communities in two north Florida salt marshes: part II. macroinvertebrate communities: Bulletin of Marine Science, vol. 26, no. 2, p. 172-195.

SALT MARSH, MACROINVERTEBRATES, FLORIDA, COMMUNITIES, DENSITY, JUNCUS, BIOMASS, DETRITUS, LOW MARSH, UPPER MARSH, HIGH MARSH

Abstract--"The mean density of marsh invertebrates in trap samples was 475/m² and average densities were significantly higher in low marsh (540/m²) than in the upper marsh (381/m²). Peaks of abundance were observed in winter and fall. Mean biomass for the year was 123g/m². Species diversity was significantly different among seasons with peaks in spring and fall. Species richness and equitability closely followed the seasonal trends of diversity. Mean homogeneity of species occurrence was 33% between the low and upper marsh zones. A total of 51 species were encountered and marsh zones exhibited distinct species assemblages. Based on biological index values, the lower and upper marsh were Littorina-Cyathura-tanaidacean communities with an abundance of three polychaetes. High marsh zones were characterized by an abundance of Uca spp., Melampus bidentatus, and Cerithidea scalariformis. Data from 1-m² areas along the two transects revealed a distinct horizontal zonation of species distribution."

Summerhayes, C. P., Ellis, J. P., and Stoffers, P., 1985, Estuaries as sinks for sediment and industrial waste--a case history from the Massachusetts coast: Contributions to Sedimentology, 14, Stuttgart, E. Schweizerbart'sche Verlagsbuchhandlung (Nagel u. Obermiller), 47 p.

ESTUARY, SEDIMENTS, SEDIMENT SINKS, MASSACHUSETTS, TRACE METALS

Like many other estuaries with moderate to low river discharge the Achushnet estuary is gradually being filled with silt transported landward from the continental shelf. Flood tidal currents are stronger than ebb tidal currents which tend to fractionate the silt and clay fraction with silt remaining in the harbor and clay being exported into the adjacent bay. The clay is accompanied by trace metals which form a thin blanket (20 Cm) over the bay floor (Buzzards Bay). Indicates that the harbor is a leaky sink because about 25% of the industrial waste has escaped down the estuary.

Swanson, R. L., and Thurlow, C. I., 1973, Recent subsidence rates along the Texas and Louisiana coasts as determined from tide measurements: *Journal of Geophysical Research*, v. 78, no. 5, p. 2665-2671.

SEA LEVEL, SUBSIDENCE, TIDES, TEXAS, LOUISIANA

Rates of sea level rise were determined by comparing tide records with long-term tide records at Pensacola, Florida, which was assumed to be stable. Sea level and tide level rose during the years 1959-1970 from as little as 0.5 cm/yr at Port Isabel to a maximum of 4.3 cm/yr at South Pass, Louisiana. Where records were available before 1959, sea level rise was greater between 1959 and 1970 than before 1959. The rise was attributed to subsidence. ESTIMATED RISE IN SUBSIDENCE

LOCATION	TIME PERIOD	RATE CM/YR	LOCATION	TIME PERIOD	RATE CM/YR		
Sabine Pass	1960-1969	1.25	Gal. Pier 21	1948-1959	0.24		
1959-1971	1.12	Port Aransas	1959-1969	1.28	Padre Island	1959-1971	-0.08
1948-1959	-0.02	1959-1971	0.49				

Tacconi, P., and Billi, P., 1987, Bed load transport measurements by the vortex-tube trap on Virginio Creek, Italy, in Thorne, C. R., Bathurst, J. C., and Hey, R. D., eds., *Sediment transport in gravel-bed rivers*: John Wiley & Sons Ltd, New York, p. 583-606.

BED LOAD, RIVERS, SEDIMENT TRAPS, SEDIMENT TRANSPORT

Among results: study demonstrates the pulsating nature of bed load transport and the very low correlations between flow, bed load and grain size characteristics of the sediment.

Tenore, K. R., 1977, Food chain pathways in detrital feeding benthic communities: a review, with new observations on sediment resuspension and detrital recycling, in Coull, B. C., ed., *Ecology of marine benthos*: Columbia, South Carolina, University of South Carolina Press, p. 37-53.

DETRITUS, BENTHIC, COMMUNITIES, SEDIMENT RESUSPENSION, ECOSYSTEMS, SUBSTRATE, FECAL PELLETS, BACTERIA, PSEUDOFECES, DEPOSIT FEEDERS, MACROPHYTES, ORGANICS

A considerable amount of organic matter becomes detritus derived from fecal pellets and pseudofeces. Up to 30% of the organic carbon ingested by the zooplankton ends up in fecal pellets. However,

due to their high rate of primary productivity, macrophytes are perhaps the major primary source of detrital material in most coastal environments. It has been demonstrated that the observed correlation between particle size of the sediment and the density of deposit feeders is due to associated levels of organic matter. Sediment resuspension, consisting mostly of zooplankton fecal pellets, accounted for a significant portion of high values of sedimentation (Steele and Baird, 1972). There is good section on sediment resuspension and detrital recycling (p.44).

Tenore, K. R., 1972, Macrobenthos of the Pamlico River estuary, North Carolina: Ecological Monographs, vol. 42, no. 1, p. 51-69.

PAMLICO RIVER, NORTH CAROLINA, MACROBENTHOS, DIVERSITY, SALINITY, ANOXIC, DENSITY, BIOMASS, STABILITY

"A preliminary sampling program in July 1968 showed that the macrobenthos was characterized by both low diversity and low density because of unfavorable environmental conditions of salinity and sediment."

Texas Department of Water Resources, 1978b, Hydrological and biological studies of the Colorado River delta, Texas: Texas Department of Water Resources, LP-79, 73p.

COLORADO RIVER, MARSH VEGETATION, NUTRIENT EXCHANGE, TIDES, PRIMARY PRODUCTION, FLOW RECORDS, BAY CITY, SPARTINA ALTERNIFLORA

Contains information on plant species composition of marsh vegetation zones. Also, contains flow records for parts of April, May, and July, 1977. Most of the information is on nutrients. Some information on tidal elevations at Culver Cut, Tiger Island Cut, mouth of Colorado, and ICWW east of Matagorda.

Texas Department of Water Resources, Division of Planning and Development, 1978, Present and future surface-water availability in the Colorado River basin, Texas: Texas Department of Water Resources, LP-60, 160p., 5 appendices.

COLORADO RIVER, RESERVOIR, SEDIMENT, MATAGORDA, MONTHLY FLOWS

Contains some sediment data relating to reservoir capacity (see p. V-24 and V-25). Also contains flow data for Colorado at Matagorda from 1941 to 1965 (p. IX-5 and in appendix).

Thorson, G., 1957, Bottom communities (sublittoral or shallow shelf): Geological Society of America, Memoir 67, vol. 1, p. 461-534.

SHELF, SUBLITTORAL, COMMUNITIES, SUBSTRATE, EPIFAUNAL, INFAUNAL, BIOMASS, LARVAE, PRODUCTIVITY, MEIOBENTHOS

Pioneering study of benthic communities and their relationship with substrate.

Trent, L., Pullen, E., and Proctor, R., 1976, Abundance of macrocrustaceans in a natural marsh and a marsh altered by dredging, bulkheading, and filling: Fishery Bulletin, vol. 74, no. 1, p.195-200.

WEST BAY, GALVESTON, DREDGING, BULKHEADING, FILLING, MACROCRUSTACEANS, MARSH, DETRITUS, MACROINVERTEBRATES, EUTROPHIC, SPARTINA ALTERNIFLORA

From abstract-"Indices of abundance of macrocrustaceans during March-October 1969 in West Bay, Texas were determined for day and night and statistically compared between 1)a natural marsh area, 2)upland and bayward canal areas of a housing development, and 3)an open bay area....Catches of brown shrimp, *Penaeus aztecus*, white shrimp, *P. setiferus*, blue crab, *Callinectes sapidus*, and pink shrimp, *P. duorarum*, were significantly greater at night than during the day at one or more stations in the marsh....The generally lower catches of each species in the open bay and upland canal areas when compared with the marsh and bayward canal areas were attributed to: 1) permanent loss of intertidal vegetation in the housing development; 2) low abundance of detrital material and benthic macroinvertebrates in the open bay and upland canal areas; and 3) eutrophic conditions in the upland canal area."

Turner, R. E., 1977, Intertidal vegetation and commercial yields of penaeid shrimp: Transactions of the American Fisheries Society, vol. 106, no. 5, p. 411-416.

INTERTIDAL, VEGETATION, SHRIMP, MARSH, ESTUARINE, INTERTIDAL

"The analysis of these data supports the hypothesis that the abundance and type of commercially valuable quantities of penaeid shrimp are directly related to the absolute area and the type of estuarine-intertidal vegetation. The percent of brown shrimp are directly correlated with the percent saline vegetation."

U. S. Army Corps of Engineers, 1967, Interim report on Colorado River and tributaries, Texas, covering mouth of Colorado River: U. S. Army Engineer District, Galveston, Texas, 29 p., 6 plates, 4 appendices, attachments.

COLORADO RIVER, FLOW, SEDIMENTS

Studies indicate that practically all of the sediment presently flowing into the Gulf originates from the watershed area below a series of dams constructed at and above Austin, which control 89 % of the 42,344 sq. mi. Colorado River basin. The present sediment load at the mouth of the river is estimated to be about 1,665 acre-ft/yr, based on unit sediment weight of 70 lbs/cu-ft. This sediment load would be greatly reduced by recommended Columbus Bend Reservoir and Matagorda Reservoir. Colorado River discharge normally varies annually between 1,000 cfs and 40,000 cfs. Discharges above 50,000 second-ft overflow the river banks above Matagorda and flow across the flood plain to the bay. Watershed sediment production rates published by the TWDB indicate under present conditions of watershed development, the sediment load at the mouth of the river is 2,700,000 cu-yds/yr.

U. S. Army Corps of Engineers, undated Trinity River Project, Texas, general design memorandum, phase 1, Appendix 3, planning background, vol. III of VI.

TRINITY RIVER, TEXAS, SEDIMENT

"At present only about 2 percent of the sediment entering Lake Livingston passed through the spillway, and most sediment contributions from tributaries below Livingston dam will be inhibited by Wallisville dam. It seems likely that, in the future, aggradation and progradation of the delta will occur only during high flows, and that the primary agents in the shaping of the delta will be wind-generated wave erosion and deposition". p. A-10

United States Department of Agriculture, 1972, Texas Coastal Basins: Interim report, Type IV Cooperative River Basin Survey, 262p.

SEDIMENTATION, SOILS, RIVER BASIN, EROSION, DEPOSITION, TEXAS, SEDIMENT LOAD, SHEET EROSION, SHORELINE EROSION

Book from reading room--information on soils in coastal basins. Also, contains sections on sedimentation and graphs on suspended sediment loads for Brazos, Trinity, and Colorado Rivers from 1930 to 1965. There is table on estimated annual sediment yield for bays and estuaries for 1972.

Van Beek, J. L., and Meyer-Arendt, K., 1981, Sediment--asset or liability, in Cross, R. D., and Williams, D. L., eds., Proceedings of the National Symposium on freshwater inflow to estuaries, volume II: U. S. Department of Interior, Fish and Wildlife Service, Slidell, Louisiana, FWS/OBS 81/04, p. 197-215.

COLORADO RIVER, DELTA, SEDIMENTATION, MISSISSIPPI DELTA, NAVIGATION CHANNELS, PROGRADATION RATE

Discusses deltaic processes and the effects of dredging and spoil disposal on sediment distribution and retention. Notes that delta growth at the mouth of the Colorado River practically terminated as a result of the extension of the channel across the peninsula in 1936, maintenance dredging and spoil disposal, and construction of reservoirs along the river upstream. Exception is limited growth of the delta at Tiger Island Cut since 1950. With diversion, estimate delta growth of 40 acres/yr (16 ha), contingent upon one or more years of rapid subaerial growth during major floods alternating with longer periods of subaqueous growth. In Atchafalaya Bay, comparison of sediment input and total deposition for the period of 1967-1977 indicates less than 30% retention which is well below the 70% average calculated for Mississippi River subdeltas (Gagliano and van Beek, 1976). Modification of major deltaic processes by navigation channels and associated spoil deposits reduces delta development potential. Delta progradation occurs progressively seaward along single channel where greater water depth reduces subaerial growth.

Van Heerden, I. Li., Wells, J. T., and Roberts, H. H., 1981, Evolution and morphology of sedimentary environments, Atchafalaya Delta, Louisiana: Gulf Coast Association of Geological Societies, Transactions, v. 31, p. 399-408.

DELTA, SEDIMENTS, DEPOSITIONAL ENVIRONMENT, ATCHAFALAYA DELTA, LOUISIANA

Used cores and bathymetric charts to investigate the morphology and sedimentary environments of the delta. The ratio of the fluvial to basinal energy regime determines the morphology and sedimentary parameters of each depositional environment. Levee environments: 1. High sedimentation rate deposits ($> 0.3\text{m/annual flood}$); channel flank core of delta showed 1.5 m of deposition in 1979 flood (van Heerden, 1980); 2. Medium sedimentation rate (0.15 to $0.30\text{ m/annual flood}$); 3. Low sedimentation ($< 0.15\text{ m/annual flood}$). Because floods of 1976, 1977, and 1978 were below normal, sedimentation rate was low and basinal processes dominated. Winter and flood deposits are discernible for each year since the 1976/1977 winter. Winter material contains less clay and organic material--erosion and redistribution of levee material related to water level setup/setdown and wind and wave action with frontal passage. Sedimentation during rest of year occurs by levee overtopping; leaf litter and roots yield high organics.

Van Heerden, I. Li., and Roberts, H. H., 1980, The Atchafalaya Delta--Louisiana's new prograding coast: Gulf Coast Association of Geological Societies, Transactions, v. 30, p. 497-506.

DELTA, SEDIMENTATION, GROWTH RATE, FLOODS, EROSION, ATCHAFALAYA, LOUISIANA

Presented evidence that the rate of subaerial growth in the Atchafalaya Delta is related to the annual flood discharge. Large increases in area occurred following major floods in 1973, 1974, 1975, and 1979. From 1976 to 1978, lobes were eroded, while subaqueous levees and abandoned channels became subaerial.

Virnstein, R. W., Mikkelsen, P. S., Cairns, K. D., and Capone, M. A., 1983, Seagrass beds versus sand bottoms: the trophic importance of their associated benthic invertebrates: Florida Scientist, vol. 46, no. 3/4, p. 363-381.

SEAGRASS, BENTHIC INVERTEBRATES, TROPHIC, SAND, EPIFAUNAL, INFAUNAL, PREDATORS, MACROBENTHOS, INVERTEBRATES, VEGETATED, UNVEGETATED, SEDIMENT, HABITAT, FOOD, PROTECTION, MACROFAUNA, FLORIDA, INDIAN RIVER

From abstract--"Seagrass beds in the Indian River lagoon, Florida, had 3 times the density of macrobenthic invertebrates found in unvegetated sediments a few meters away. Epifaunal abundance was 13 times greater in seagrass than sand. Most epifaunal species were uncommon in the unvegetated sediments, presumably due to their dependence on seagrass for providing habitat, food, nursery area, and/or protection from predators. Not only were macrofaunal densities much higher in natural seagrass than in sand, those animals which were more abundant (primarily the epifauna) were also more heavily preyed upon and thus are trophically more important than

infauna...The grazing epifauna of seagrass meadows forms a major trophic pathway to higher predators via the decapods; such a pathway is lacking in sand bottom communities."

Ward, G. H., Armstrong, N. E., and the Matagorda Bay Project Teams, 1980, Matagorda Bay Texas: its hydrography, ecology and fishery resources: U.S. Fish and Wildlife Service, Biological Services Program, Washington D.C., FWS/OBS-81/52, 230 p.

MATAGORDA BAY, TEXAS, HYDROGRAPHY, SEDIMENTS, MARSHES, STREAMS, SEDIMENT LOAD, COLORADO RIVER

Ward and Armstrong discuss briefly the sedimentary processes of the bay system and illustrate the annual silt loads of major rivers discharging into Matagorda Bay. The history of delta development of the Colorado River is also presented. Authors state that after 1940, no further substantial development of the Colorado River delta occurred, which they attributed to completion of the Highland Lakes chain and the fact that only 8% of the Colorado River watershed occurs downstream from Austin. The sediment rate of 550 acres per yr which occurred between 1930 to 1941 could not be sustained with a sediment load that is an order of magnitude lower than prior to 1940. With regard to the Lavaca delta organic carbon import/export, the authors note that it appears that large amounts of detrital material decompose in the vegetative zone and perhaps go into the production of peat (p. 77). (No data on peat presented).

Warwick, R., and Price, R., 1975, Macrofauna production in an estuarine mud-flat: Journal Marine Biological Association of the United Kingdom, vol. 55, no. 1, p. 1-18.

MUDFLAT, ESTUARY, MACROFAUNA, PRODUCTION, BENTHIC COMMUNITY

From abstract--"The total production was 13.311g/m²/year, which coincidentally almost exactly equalled the total mean biomass of 13.240g/m². Comparison with other areas shows that this production figure is about average for benthic communities. However, it is thought that the meiofauna production must be high in this situation because the top macrofauna producer, *Nephtys hombergi*, is a carnivore."

Welborn, C. T., and Andrews, F. L., 1980, Sediment and water discharge of the Colorado River at Matagorda, Texas May 4-7, and June 7-11, 1979: U. S. Department of the Interior, Geological Survey, Open file report 80-1177, 21p.

SEDIMENT, COLORADO RIVER, MATAGORDA, TEXAS, FLOOD, SUSPENDED LOAD, BED LOAD

Abstract--"The U.S. Army Corps of Engineers requested the U.S.G.S. to collect data on the Colorado River at Matagorda, Texas, to determine the total sediment discharge into Matagorda Bay if all the water from the Colorado River were diverted into the bay. Data were collected from two floods (May 4-7, and June 7-11, 1979) with peak discharges of 24,900 cubic feet per second and 40,900

cubic feet per second. The total suspended-sediment discharge from the first flood was approximately 201,000 tons, and the total suspended-sediment discharge from the second flood was 630,000 tons. The volume of sediment from these two floods was approximately 970 acre-feet, with approximately 20 acre-feet of this total being the bed load or 2.1% of total sediment load." "If all the water from these two floods had been channeled into Matagorda Bay, approximately 831,000 tons of suspended sediment would have entered the bay."

Whitlatch, R. B., 1977, Seasonal changes in the community structure of the macrobenthos inhabiting the intertidal sand and mud flats of Barnstable Harbor, Massachusetts: Biological Bulletin, vol. 152, p. 275-294.

BARNSTABLE HARBOR, MASSACHUSETTS, COMMUNITY, MACROBENTHOS, SANDFLAT, MUDFLAT, SEASONAL, SUSPENSION FEEDER, DEPOSIT FEEDER, INSTABILITY, GRAIN SIZE

Data pretty well supports Sander's work--suspension feeders in fine sands and deposit feeders in muds. Sanders said optimal grain size for suspension feeders was 0.18mm but in this study, optimal size was 0.28mm. This study also supports idea that suspension-feeders greatly effected by sediment instability. Data indicates that abundance of both deposit and suspension feeders greatly reduced in areas with pronounced surficial ripple marks (evidence of sediment instability).

Wilkinson, B. H., and Byrne, J. R., 1977, Lavaca Bay transgressive deltaic sedimentation in central Texas USA estuary, American Association of Petroleum Geologists Bulletin, v. 61, no. 4, p. 527-545.

ESTUARY, LAVACA BAY, DELTA, SEDIMENTATION, TEXAS

Following from BIOSIS: Holocene sediments filling the Lavaca River Valley [USA] comprise a transgressive fluvial-deltaic-estuarine sequence which was deposited during the past 10,000 yrs. Coarse fluvial sands which occupy the valley axis grade upward into deltaic sands and muddy sands. These are overlain by 4 lithologically distinct mud facies which reflect estuarine conditions. Several different sources of argillaceous sediment were important during deposition of this valley-filling sequence. Muds delivered into Lavaca Bay by fluvial discharge are volumetrically most important, comprising 75% of the total fine fraction. Wave erosion of Pleistocene sedimentary strata exposed along the bay margin is a less important but significant source of argillaceous sediment, making up 25% of the estuarine muds. Sedimentation rates determined for the past 10,000 yr are comparable to those reported for the past 100 yr, but suggest increased rates of mud deposition in modern times.

Williams, G. P., and Wolman, G. M., 1984, Downstream effects of dams on alluvial rivers: U. S. Geological Survey Professional Paper 1286, 64 p.

DAMS, RIVERS, SEDIMENT LOADS

Describes changes in mean channel bed elevations, channel width, bed-material sizes, vegetation, water discharges, and sediment loads downstream from 21 dams constructed on alluvial rivers (most from

the semiarid western U.S.; Neches River in Texas included). "Sediment concentrations and suspended loads were decreased markedly for hundreds of kilometers downstream from dams; post-dam annual sediment loads on some rivers did not equal pre-dam loads anywhere downstream from a dam". Riparian vegetation commonly increased downstream from the dam probably as a result of decreased peak flows.

Williams, R. B., and Thomas, L. K., 1967, The standing crop of benthic animals in a North Carolina estuarine area: *Journal of the Elisha Mitchell Scientific Society*, vol. 83, p. 135-139.

STANDING CROP, BENTHOS, BIOMASS, SEAGRASS, NORTH CAROLINA, SEDIMENT, EELGRASS, SCREEN SIZE

This study gives standing crop of benthos for different sediments and for muddy sand with marine grass. Standing crop highest in grass, with muddy sand next. Screen size used was 6mm. Article also talks about larger benthos forming bulk of biomass.

Windom, H. L., Neal, W. J., and Beck, K. C., 1971, Mineralogy of sediments in three Georgia estuaries: *Journal of Sedimentary Petrology*, v. 41, p. 497-504.

ESTUARIES, SEDIMENTS, MINERALOGY, GEORGIA, SOURCES

Mineralogical analyses of suspended sediments indicate that the clastics are mixtures of fluvially derived detritus and fines derived from the continental shelf. Decreasing-flow velocity and focculation of the clays results in a general change in bottom sediments from sand in the rivers to clay in the estuaries. Sands and silts are high in quartz with significant amounts of feldspar. Clay mineralogy is also discussed. Supports conclusions of previous workers that shoreward transport, especially for the fine fraction, is an important process in determining the composition of the estuarine sediments.

Winston, J. E., and Anderson, F. E., 1971, Bioturbation of sediments in a northern temperate estuary: *Marine Geology*, v.10, p. 39-49.

BIOTURBATION, SEDIMENT, GREAT BAY, NEW HAMPSHIRE, SALINITY, SAND, BRACKISH WATER, SILT, MUDFLAT, BIOMASS, ESTUARY, TIDAL FLATS, STRATIFICATION, POLYCHAETE, NEREIS

"In the Great Bay estuarine system the primary controls on biomass and faunal diversity appeared to be the salinity gradient and textural changes within the estuary. In general, bioturbation decreased up the estuary." The polychaete species of the genus *Nereis* were the most important turbating agents. Four different levels of bioturbation were observed ranging from highly turbate in a sandy bottom, marine environment near the mouth of the estuary to non-turbate in a silty bottom, brackish-water environment near the head of the estuary.

Wright, L. D., 1985, River deltas, in Davis, R. A., Jr., Coastal Sedimentary Environments, 2nd revised, expanded edition: Springer-verlag, New York, p. 1-76.

RIVERS, DELTAS, SEDIMENTS, CURRENTS, WAVES, ENVIRONMENTS, FACIES

Zetler, B. D., 1953, Some effects of the diversion of the Santee River on the waters of Charleston Harbor: Transactions, American Geophysical Union, vol.34, no.5, p. 729-732.

DIVERSION, SANTEE RIVER, CHARLESTON, SOUTH CAROLINA, SEA LEVEL, SALINITY, DENSITY, TIDE GAGE, TIDAL RANGE

Copied article from journal in reading room. "Surface temperature and density observations in Charleston Harbor indicated a sharp decrease in salinity beginning in 1942. It was at this time that a hydroelectric plant began diverting some of the waters from the Santee River into the Cooper River which flows into Charleston Harbor. Long-period tide observations at Charleston and at other places along the Atlantic Coast were examined to detect related changes in sea level and range or tide. They indicate that about 0.09 ft of the rise in sea level in Charleston Harbor can be attributed to the diversion of the Santee River and that the tidal range may have been increased about 0.05 ft."

Zimmerman, R. J., Minello, T. J., and Zamora, G., Jr., 1984, Selection of vegetated habitat by brown shrimp, *Penaeus aztecus*, in a Galveston Bay salt marsh: Fishery Bulletin, vol. 82, no. 2, p.325-336.

SHRIMP, BROWN SHRIMP, SPARTINA ALTERNIFLORA, VEGETATION, SALT MARSH, PENAEUS AZTECUS, WEST BAY, GALVESTON, NONVEGETATED

From abstract-"Densities of the brown shrimp, *Penaeus aztecus*, in vegetated and nonvegetated habitats of a Galveston West Bay salt marsh were compared. Each of 81 sample pairs taken between 29 March and 23 July 1982 consisted of one sample from *Spartina alterniflora* habitat and another from nonvegetated habitat. Overall a mean density for shrimp of 11.7/m² in vegetation was significantly greater than the mean density of 1.4/m² in nonvegetated habitat...Mechanisms that may have enhanced utilization of vegetated habitat for *P. aztecus* were reticulation in salt marsh macrostructure, relatively low tidal range, and seasonal periods of high water. The nursery function of the salt marsh was confirmed by dominance of small shrimp, with 95% of all individuals being smaller than 50mm in rostrum through telson length."

Zimmerman, R. J., and Minello, T. J., 1984, Densities of *Penaeus aztecus*, *Penaeus setiferus*, and other natant macrofauna in a Texas salt marsh: Estuaries, vol. 7, no. 4A, p. 421-433.

TEXAS, GALVESTON ISLAND, SALT MARSH, MACROFAUNA, DENSITIES, NATANT, SHRIMP, SPARTINA ALTERNIFLORA, PENAEUS, WATER-LEVEL, VEGETATED, UNVEGETATED, CRUSTACEANS, FISH, CRABS, CALLINECTES

"Densities of the crustaceans *Palaemonetes pugio*, *Penaeus aztecus* and *Callinectes sapidus* were significantly higher in vegetated compared to non vegetated habitat, whereas *Penaeus setiferus* densities were not different. Some natant estuarine macrofauna may have become more or less dependent upon exploitation of vegetated habitat within marshes." Juvenile brown shrimp may spend most of their early residence in or near marsh habitat. "Historically, water-levels in Gulf of Mexico estuaries are highest in the spring and fall (Hicks et al. 1983). As a consequence, intertidal vegetation is more available for longer periods to juvenile shrimp and other organisms during these intervals. Water-level was the single most important factor in regression models for *P. aztecus* densities in the marsh." "Fish species were marginally more numerous within vegetation. White shrimp densities did not differ significantly between vegetated and nonvegetated habitat. Blue crabs were significantly more abundant in vegetation than on nonvegetated bottom."