

**WETLAND PLANT COMMUNITIES,  
GALVESTON BAY SYSTEM**

by  
William A. White and Jeffrey G. Paine

Assisted by  
Damon Waitt

Principal Investigator  
E. G. Wermund

Prepared for  
Galveston Bay National Estuary Program  
under contract no. IAC(90-91)1445

Bureau of Economic Geology  
W. L. Fisher, Director  
The University of Texas at Austin  
Austin, Texas 78713-7508

1992

## TABLE OF CONTENTS

EXECUTIVE SUMMARY .....	1
INTRODUCTION.....	2
General Objectives of Field Investigations.....	2
General Project Objective and Wetland Definition.....	2
Field Surveys.....	3
Methods .....	3
WETLAND COMMUNITIES IN THE GALVESTON BAY AREA.....	5
General Setting of the Galveston Bay System.....	5
Classification of Wetland Communities: Background and Previous Studies .....	5
Species Composition of Wetland Plant Communities, Galveston Bay System.....	13
Wetland Indicator Status of Prevalent Plants at Survey Sites.....	23
Wetland Plant Communities .....	23
Salt-Marsh Community.....	27
Brackish-Marsh Community.....	27
Fresh-Marsh Community.....	38
Forested Wetland Communities (Swamps) .....	42
Submerged Vegetation Community .....	42
Soils and Wetland Community Relationships.....	42
Examples of Wetland Profiles Developed from Topographic Survey Transects .....	49
Smith Point Transect.....	49
Brazoria National Wildlife Refuge Transect.....	51
ACKNOWLEDGMENTS .....	52
REFERENCES.....	53
APPENDICES	
A. Field Site Surveys.....	57
B. List of Plant Species by Site Number.....	63
C. Elevation Transects.....	85

## Figures

1.	Index map of the Galveston Bay area.....	4
2.	Natural systems in the Galveston Bay area.....	6
3.	Location map of field survey sites.....	15
4.	Index map of quadrangles covering the Galveston Bay area.....	19
5.	Low salt-marsh community of <i>Spartina alterniflora</i> and open water on the inland margins of Jones Bay (east end of West Bay).....	28
6.	Salt-marsh community on Follets Island.....	28
7.	Low salt-marsh community inland from West Bay .....	29
8.	Salt-marsh community on the bayward margin of Bolivar Peninsula.....	29
9.	Salt-marsh community at Houston Point .....	30
10.	Salt marsh/sand flat community on Follets Island.....	30
11.	Profile of salt marsh at Smith Point showing relative elevations of plant communities ....	31
12.	Profile of brackish marsh in the Brazoria National Wildlife Refuge showing relative elevations of plant communities.....	32
13.	Brackish-marsh community in the Brazoria National Wildlife Refuge southwest of Hoskins Mound.....	34
14.	Brackish-marsh community in the Brazoria National Wildlife Refuge east of Hoskins Mound.....	34
15.	High and low brackish-marsh communities in the Anahuac National Wildlife Refuge .....	35
16.	Brackish-marsh community dominated by <i>Spartina patens</i> , west of High Island.....	35
17.	Brackish-marsh community in a swale on Galveston Island.....	36
18.	Brackish- to fresh-marsh community in a depression on Follets Island, gulfward of highway.....	36
19.	Brackish-marsh community on the Trinity River delta.....	37
20.	Brackish-marsh community on the Trinity River delta near the delta/bay margin.....	37
21.	Fresh-marsh community in the Trinity River valley north of Interstate Highway 10 .....	39
22.	Fresh- to brackish-marsh community on the Trinity River delta near Old River Lake.....	39
23.	Fresh-marsh and forested-wetland communities in the San Jacinto River valley .....	40
24.	Fresh-marsh community of <i>Scirpus californicus</i> in an ox-bow lake in the Brazoria National Wildlife Refuge .....	41
25.	Fresh-marsh community dominated by <i>Eleocharis quadrangulata</i> (squarestem spikesedge) .....	41

26. Swamp community dominated by <i>Taxodium distichum</i> along the Trinity River inland from Interstate Highway 10 .....	43
27. Generalized map showing the locations of submerged vegetation along the margins of the Trinity River delta .....	44
28. Generalized map showing the locations of submerged vegetation in 1956 and 1987 in the Galveston Bay system.....	45
29. Disturbed-area community on a small spoil mound along the Intracoastal Waterway on the landward margin of Follets Island.....	50

### Tables

1. Generalized characteristics of active coastal processes and conditions in the Galveston Bay area .....	7
2. Dominant and common plants in brackish and fresh marshes in Chambers County.....	9
3. Common species in salt and fresh marshes in the Brazoria National Wildlife Refuge, Brazoria County .....	10
4. Typical plants found in grassflats, marshes, and transitional areas in the Galveston Bay area.....	11
5. Typical plants identified at saline sites at Armand Bayou and vicinity .....	12
6. Water regime descriptions for wetlands used in the Cowardin and others (1979) classification system.....	14
7. List of USGS 7.5-minute topographic maps that encompass the Galveston Bay project area.....	20
8. List of common plant species for various marshes based on field surveys.....	21
9. Species most frequently observed at survey sites in the study area.....	22
10. Wetland indicator status and common names of plants identified in field surveys.....	24
11. Characteristics of typical soils at field survey sites.....	46

## EXECUTIVE SUMMARY

by William A. White and Jeffrey G. Paine

Wetlands and aquatic habitats are critical components of the biologically productive Galveston Bay estuarine system. This report is the culmination of a field investigation of wetland plant communities, and is one phase of the project, "Trends and Status of Wetland and Aquatic Habitats of the Galveston Bay System, Texas," sponsored by the Galveston Bay National Estuary Program (GBNEP). For purposes of this topical report, wetlands are defined and classified in terms of more classical definitions, for example, salt, brackish, and fresh marshes, in accordance with project requirements. The relationship of these wetland classes to the Cowardin and others (1979) classification system used to map wetlands is presented through various examples. Wetlands in this study were not defined in accordance with the "Federal Manual for Identifying and Delineating Jurisdictional Wetlands" (currently being revised) and, thus, should not be regarded as jurisdictional wetlands.

More than 150 sites were examined in the Galveston Bay system. Wetland plants were identified at selected field survey sites, principally along transects aligned perpendicular to the hydrologic gradient so that plant assemblages from the water's edge to upland areas were intercepted. Topography surveys were conducted along several transects. Measurements of elevation, distance, and plant community composition were made along the survey lines, which crossed salt marshes and brackish to fresh marshes. Elevations were measured to the nearest 0.5 cm and distances to the nearest meter. County soil surveys were used to define and characterize soils at the various field check sites. The locations of field survey sites were plotted on aerial photographs, and later accurately transferred to USGS 7.5-minute quadrangle topographic maps. Universal Transverse Mercator (UTM) coordinates were determined for each site and these data were entered into computer data management systems, including the geographic information system, ARC-INFO.

The most widely distributed wetland environments in the Galveston Bay system are marshes, the most extensive of which are brackish. Brackish marshes compose roughly 65 to 70 percent of the marsh system in the Galveston Bay project area. Salt marshes are a distant second, composing roughly 25 to 30 percent. Fresh marshes make up the remaining 5 to 10 percent of the marsh system. Many species can tolerate varying salinity regimes as well as water regimes, and there is, therefore, considerable overlap in the species composition of these marsh systems. Because of the predominance of brackish and salt marshes in the project area, more than 60 percent of the field surveys were located in these marshes. With reference to all sites visited, the 15 most frequently encountered species were headed by *Spartina patens* and *Distichlis spicata*. Other major species include *Spartina alterniflora*, *Batis maritima*, *Salicornia* spp., *Iva frutescens*, *Spartina spartinae*, *Borrichia frutescens*, *Juncus roemerianus*, *Aster* spp., *Typha* spp., *Scirpus maritimus*, and *Monanthochloe littoralis*. Of the species identified at the survey sites, about 34 percent are classified as obligate wetland plants, which means that under natural conditions these plants occur in wetlands with an estimated probability of 99 percent. Among these species are those typically found in wetter conditions (for example, those characterizing topographically low salt, brackish, and fresh marshes). Approximately 37 percent of the species listed are classified as facultative wetland plants. These species usually occur in wetlands or have an estimated probability of 67-99 percent of occurring in wetlands but occasionally they occur in nonwetland areas. These species typically define topographically higher marshes. About 19 percent of the listed species are classified as facultative. These species are equally likely to occur in wetlands or nonwetlands (estimated probability 34-66 percent). At the more than 135 sites surveyed for vegetation around the Galveston Bay system, approximately 40 soil types were identified from county soil surveys. Several soils were encountered more frequently than others, and can be considered the dominant soils corresponding to wetland communities. For example, the soil most frequently occurring at wetland survey sites was the Harris clay. This typically saline, poorly-drained soil is flooded by abnormally high tides, and supports a vegetation assemblage composed predominantly of *Spartina patens* and *Distichlis spicata*. These species were the most frequently encountered during field surveys.

## **WETLAND PLANT COMMUNITIES, GALVESTON BAY SYSTEM**

### **INTRODUCTION**

Wetlands and aquatic habitats are critical components of the biologically productive Galveston Bay estuarine system. Mapping and describing the composition of these important habitats are essential steps in defining their status and in measuring and anticipating the effects of the numerous coastal activities that directly and indirectly influence them. Understanding cause-and-effect relationships can be promoted only through such detailed scientific investigations.

This report is the culmination of a field investigation of wetland plant communities, and is one phase of the project to determine the "Trends and Status of Wetland and Aquatic Habitats of the Galveston Bay System, Texas," sponsored by the Galveston Bay National Estuary Program (GBNEP).

### **General Objectives of Field Investigations**

The purpose of these field investigations was to characterize wetland plant communities through representative field surveys, fundamental to the comparison of various wetland plant communities in the field with corresponding "signatures" on aerial photographs used to define wetland classes, including water regimes, for mapping purposes. In fact, all field work was done with reference to aerial photographs. This topical report presents results of representative field surveys and focuses principally on characterizing prevalent plant associations in the Galveston Bay System. For the grander objectives of the GBNEP contract, these characterizations also provided vital plant community information for defining the appropriate wetland classes and water regimes during the extensive "ground truthing" surveys in which wetland signatures delineated on aerial photographs were correlated with plant communities in the field. Characterization of plant communities in the field surveys allowed mapped wetland classes to be better defined in terms of typical vegetation associations.

### **General Project Objective and Wetland Definition**

The fundamental objective of the GBNEP project, for which this reported study is one phase, is to determine the trends and status of wetlands in the Galveston Bay System using aerial photographs. The definition and identification of wetlands, therefore, is integrally connected to the photographs. Wetlands were delineated on mid-1950's, 1979, and 1989 photographs as part of the U.S. Fish and Wildlife Service (USFWS) National Wetlands Inventory program using the Cowardin and others (1979) wetland classification system. Even though wetlands delineated on aerial photographs are supported by field surveys (especially for the 1989 delineations), field-identified wetlands represent only a small percentage of all the wetlands delineated. During ground-truth surveys, prevalent plant species associations <sup>were</sup> characterized "within the constraints imposed by the resolution of the photos" (as stated in the Project Scope of Work, 1990). Wetlands <sup>were</sup> not identified in accordance with the *Federal Manual for Identifying and Delineating Jurisdictional Wetlands* (this manual is currently being revised). Thus, the wetlands mapped and defined in this study are not jurisdictional wetlands. The following is printed on all wetland maps that are used in this project to determine the status and trends of wetlands in the Galveston Bay system:

This document (map) was prepared primarily by stereoscopic analysis of high altitude aerial photographs. Wetlands were identified on the photographs based on vegetation, visible hydrology, and geography in accordance with "Classification of Wetlands and Deepwater Habitats of the United States" (FWS/OBS - 79/31 December 1979). The aerial photographs typically reflect conditions during the specific year and season when they were taken. In addition, there is a margin of error inherent in the use of the aerial photographs:

Federal, State, and local regulatory agencies with jurisdiction over wetlands may define and describe wetlands in a different manner than that used in this inventory. There is no attempt, in either the design or products of this inventory, to define the limits of proprietary jurisdiction of any Federal, State or local government or to establish the geographical scope of the regulatory programs of government agencies.

For purposes of this report, and in accordance with project requirements, wetlands are defined and classified in terms of more classical definitions, for example, salt, brackish, and fresh marshes. The relationship of these wetland classes to the Cowardin and others (1979) classification system is presented through various examples.

### Field Surveys

More than 150 sites were examined in the Galveston Bay system (fig. 1) at locations that included the Brazoria and Anahuac National Wildlife Refuges, Armand Bayou Nature Center, Follets and Galveston Islands, Bolivar Peninsula, Smith Point, High Island area, Trinity River delta, and other areas. Plant communities were surveyed during the months of June, July, and November 1990, and May and September 1991. The surveys were conducted principally by the authors; other personnel involved in one or more surveys included Larry Handley (USFWS, National Wetlands Research Center), Warren Hagenbuck and Curtis Carley (USFWS National Wetlands Inventory), Todd Mecklenborg (Geonex Martel, Inc.), and Warren Pulich (Texas Parks and Wildlife Department). In addition, Ron Bisbee (Refuge Manager), Richard Antonette and Mike Lange of the Brazoria National Wildlife Refuge, and Jim Neaville and Ed Jackson of the Anahuac National Wildlife Refuge accompanied field parties to their respective areas.

### Methods

During the initial field investigations, methods were developed to characterize prevalent species associations. The primary method was one in which wetland plants were identified at selected field survey sites, principally along transects aligned perpendicular to the hydrologic gradient so that plant assemblages from the water's edge to upland areas were intercepted. A second approach was to conduct a topographic survey along selected transects that crossed representative plant communities to identify relative elevations at which various plant species occur. This is helpful in defining water regimes and in differentiating between high- and low-marsh communities. The boundaries between some plant assemblages are controlled in part by elevation, so elevation measurements focus on such boundaries. Plant species that were difficult to identify in the field were collected for identification in the laboratory or with reference to the plant collection at The University of Texas Herbarium.

Topography surveys were conducted along several transects. Measurements of elevation, distance, and plant community composition were made along the survey lines, which crossed salt marshes (Smith Point, Follets Island, and mainland margin of West Bay) and brackish to




Figure 1. Index map of the Galveston Bay area. (From White and others, 1985)

fresh marshes (Anahuac National Wildlife Refuge, Brazoria National Wildlife Refuge, and Trinity River Delta). Elevations were measured to the nearest 0.5 cm (2 inches) and distances were measured to the nearest meter. Compass bearings of the transects were also recorded.

County soil surveys (Brazoria, Chambers, Galveston, and Harris Counties) were used to define and characterize soils at the various field check sites. Information obtained from the soil surveys included soil type, salinity, drainage, frequency of flooding, position of water table, and prevalent vegetation.

The locations of field survey sites were plotted on aerial photographs, and later accurately transferred to USGS 7.5-minute quadrangle topographic maps using a Zoom Transfer Scope where necessary. Universal Transverse Mercator (UTM) coordinates were determined for each site and these data were entered into computer data management systems, including the geographic information system, ARC-INFO.

## WETLAND COMMUNITIES IN THE GALVESTON BAY AREA

### General Setting of the Galveston Bay System


The geologic framework of the Galveston Bay area consists of Modern-Holocene and Pleistocene systems including the modern wetland, or marsh and marsh-swamp systems (fig. 2). The geomorphic features on which the various types of coastal wetlands have developed are the result of numerous interacting processes. Physical processes that influence wetlands include rainfall, runoff, fluctuations in the water table, streamflow, evapotranspiration, waves and longshore currents, astronomical and wind tides, storms and hurricanes, deposition and erosion, subsidence, faulting, and sea-level rise (table 1). These processes have contributed to the development of a gradational array of permanently inundated to infrequently inundated environments ranging in elevation from the submerged lands of the estuarine system through the topographically higher wetland system, which grades upward from the astronomical-tidal zone through the wind-tidal zone to the storm-tidal zone.

Exchange of marine waters with bay-estuary-lagoon waters in the Galveston Bay system occurs primarily through two major tidal inlets: Bolivar Roads at the north end of Galveston Island and San Luis Pass at its south end (fig. 1). Additional exchange occurs at Rollover Pass, a narrow dredged channel at the east end of Bolivar Peninsula. The predominant sources of fresh-water inflow are the Trinity and San Jacinto Rivers (fig. 1). Salinities in the Galveston Bay system are generally highest in West and Christmas Bays where mean salinities are typically above 20 parts per thousand (ppt) and may range into the 30's. These salinities are in marked contrast to Trinity Bay, where Trinity River fresh-water inflows have a moderating influence; mean monthly salinities in Trinity Bay are usually less than 15 ppt and occasionally are below 5 ppt (Pulich and White, 1991).

These numerous interacting processes in the Galveston Bay system have a major bearing on the location and composition of wetland plant communities.

### Classification of Wetland Communities: Background and Previous Studies

Classification of wetland communities ranges from broad, general systems in which the entire coastal wetland system is encompassed within a single unit (usually as part of a statewide vegetation classification), to the more detailed classifications that focus specifically on coastal



#### EXPLANATION

[Modern marsh system icon]	Modern marsh system	[Pleistocene barrier-strandplain system icon]	Pleistocene barrier-strandplain system
[Modern delta/marsh system icon]	Modern delta/marsh system	[Modern-Holocene fluvial system icon]	Modern-Holocene fluvial system
[Modern marsh-swamp system icon]	Modern marsh-swamp system	[Pleistocene fluvial system icon]	Pleistocene fluvial system
[Modern barrier strandplain system icon]	Modern barrier strandplain system	[Pleistocene fluvial-deltaic system icon]	Pleistocene fluvial-deltaic system

QA 7981

Figure 2. Natural systems in the Galveston Bay area. (From Fisher and others, 1972, 1973)

Table 1. Generalized characteristics of active coastal processes and conditions in the Galveston Bay area. (From White and others, 1985)

Climatic zone:	Humid ( <i>Thornthwaite, 1948</i> )
Average annual precipitation:	41.8 to 51.5 inches/yr (106.2 to 130.8 cm/yr) ( <i>Fisher and others, 1972</i> )
Dominant wind directions:	Southeasterly, northerly ( <i>Fisher and others, 1972</i> )
Average wind speed (in 1978 at Texas City):	6.8 knots (12.6 km/hr) ( <i>Shew and others, 1981</i> )
Astronomical tidal range: Gulf shoreline (Galveston Pleasure Pier) Mean diurnal: Bay shoreline (mean):	2.1 ft (0.6 m) ( <i>U.S. Department of Commerce, 1978</i> ) 0.5 to 1.4 ft (0.2 to 0.4 m) ( <i>Diener, 1975</i> )
Tidal current velocities: Bolivar Roads Average maximum flood: Average maximum ebb:	3.3 knots (1.7 m/sec) ( <i>Bernard and others, 1959</i> ) 4.3 knots (2.2 m/sec) ( <i>Bernard and others, 1959</i> )
Wave height (Gulf): (Caplan, Texas) Onshore wave height:	Between 2.5 and 3.5 ft (0.8 and 1.1 m) about 65% of the time, ( <i>U.S. Army Corps of Engineers, 1956</i> )
Direction of net longshore sediment transport:	Southwesterly ( <i>Fisher and others, 1972</i> )
Maximum hurricane surge height on open coast:	12.7 ft (3.9 m) above MSL ( <i>Bodine, 1969</i> )
Hurricane frequency:	12% in any one year ( <i>Simpson and Lawrence, 1971</i> )
Gulf shoreline change, Bolivar Roads to San Luis Pass from 1850-52 to 1973-74:	Total gain from accretion of 1,074 acres and loss from erosion of 1,183 acres; net loss of 109 acres ( <i>Morton, 1977</i> )
Subsidence: Pasadena - Houston Ship Channel area:	8.5 to 9 ft (2.6 to 2.7 m) during 1906-1973 ( <i>Ratzlaff, 1980</i> )
Faulting: Houston metropolitan area:	Offset by at least 160 faults ( <i>Verbeek and Clanton, 1981</i> )

wetlands and subdivide them into several units. Among the broad descriptive systems are: Bray's (1906) Salt Marsh Meadows; Tharp (1926) and Godfrey and others' (1973) Coastal Marsh; Kuchler's (1966) Southern Cordgrass Prairie; Thomas (1975) and Gould's (1975) Gulf Prairies and Marshes; and Frye and others' (1984) Marsh/Barrier Island.

Among the more specific descriptions of wetland communities, which include a classification scheme and/or which focus on a significant part of the upper (north and central) Texas Coast including the Galveston Bay area, are those by Shaw and Fredine (1956), Fisher and others (1972, 1973), Diener (1975), Lazarine (n.d.), Fleetwood (n.d.), Harcombe and Neaville (1977), Adams and Tingley (1977), Benton and others (1979), Cowardin and others (1979), Gosselink and others (1979), Ward and Armstrong (1980), Shew and others (1981), Thayer and Ustach (1981), and White and others (1985 and 1987).

Most classifications have subdivisions based on salinities because the community composition of coastal wetlands is influenced by the proximity of saline and brackish waters of the marine and estuarine systems. Bray (1906) listed his Salt Marsh Meadow under a more general heading of Salt Water Vegetation. Although Tharp's (1926) coastal marsh unit was not subdivided according to salinities, he did note that giant reed (*Arundo donax*), common reed (*Phragmites australis*), and marshmillet, or southern wildrice (*Zizaniopsis miliacea*) are abundant along streams and other "semi-fresh water bodies." Shaw and Fredine (1956) used two major subdivisions in coastal areas: coastal saline areas and coastal fresh areas. Fisher and others (1972, 1973) subdivided marshes on the basis of salinities into salt-water, brackish (closed), brackish- to fresh-water, and inland fresh-water marshes. Lazarine (n.d.) in a field reference guide to common wetland plants subdivided wetland types into saline, brackish, and fresh. Gosselink and others (1979) followed Chabreck (1972) by subdividing marshes into four categories in order of decreasing salinities: saline, brackish, intermediate, and fresh (for mapping and discussion purposes, intermediate was combined with brackish). Harcombe and Neaville (1977), in describing and mapping wetlands in Chambers County, used brackish and fresh subdivisions (table 2) (salt marshes were not included because of their absence or limited areal extent). Fleetwood (n.d.) in a study of vegetation in the Brazoria Wildlife Refuge, recognized (in addition to fresh marsh) saline, brackish, and intermediate components of the marsh system, but because of "dynamic wet and dry cycle conditions" combined them into a single unit designated as salt marsh (table 3). Among the major subdivisions (systems) used by Cowardin and others (1979) are estuarine and palustrine, which in simplified terms correspond with saline-brackish and fresh marsh areas, respectively, when classifying emergent wetlands. (In coastal tidal areas, palustrine wetlands begin where salinity, due to ocean-derived salts, is below 0.5 ppt; it should be noted that salinity modifiers can be used in both the estuarine and palustrine systems so the palustrine system can have salt marshes in areas where the salts are not ocean derived.) White and others (1985) used three basic categories: salt-, brackish-, and fresh-water marshes (table 4). Saline flats and marshes were among major vegetational areas defined by Brown (1985) for southeastern Harris County (table 5). Harcombe and Neaville (1977), Fleetwood (n.d.), and Brown (1985) have compiled detailed checklists of plants, including wetland species, occurring in Chambers, Brazoria, and Harris Counties.

In addition to subdivision based on relative salinities, some classifications subdivide marsh communities on the basis of inundation frequency determined in large part by elevation with respect to mean sea level. In coastal areas where the range in astronomical (lunar) tides is high, such as along the Atlantic Coast, the salt-marsh community is commonly subdivided into distinct low and high marshes. Broad areas are flooded on a regular (daily) basis, and plants like smooth cordgrass (*Spartina alterniflora*) that live in the intertidal zone represent extensive areas of low marshlands that are readily distinguished from high, irregularly flooded marshes. However, along the Texas coast astronomical tidal ranges are low and, thus, areas flooded on a daily basis, although dominated by smooth cordgrass, are much more restricted in areal extent. Wind-driven tides have a dominant influence along the Texas coast because they flood more extensive areas.

Table 2. Dominant and common plants in brackish and fresh marshes in Chambers County.  
(From Harcombe and Neville, 1977)

**Brackish Marsh**

Dominated by:

*Spartina patens* (marsh-hay or saltmeadow cordgrass)

*Distichlis spicata* (seashore saltgrass)

Isolated clumps of:

*Scirpus maritimus* (saltmarsh bulrush)

*Scirpus olneyi* (Olney bulrush)

Near tidal drains:

*Juncus roemerianus* (needlegrass rush)

Common on levees:

*Phragmites australis* (common reed)

*Spartina cynosuroides* (big cordgrass)

Common in fresher areas:

*Paspalum vaginatum* (seashore paspalum)

*Paspalum lividum* (longtom)

Common locally:

*Spartina alterniflora* (smooth cordgrass)

**Fresh Marsh**

*Phragmites australis* (common reed)

*Cladium jamaicense* (sawgrass)

*Zizaniopsis miliacea* (cutgrass)

*Panicum repens* (torpedograss)

*Paspalum lividum* (longtom)

*Typha latifolia* and *T. domingensis* (cattail)

*Spartina cynosuroides* (big cordgrass)

*Alternanthera philoxeroides* (alligator weed)

Table 3. Common species in salt and fresh marshes in the Brazoria National Wildlife Refuge.  
(From Fleetwood, n.d.)

**Salt Marsh**

Dominants:

*Spartina patens* (marshhay cordgrass)

*Distichlis spicata* (seashore saltgrass)

In fresher areas:

*Paspalum vaginatum* (seashore paspalum)

*Scirpus olneyi* (Olney bulrush)

*Scirpus americanus* (American bulrush)

Other common species:

*Phragmites australis* (common reed)

*Paspalum lividum* (longtom)

*Aster subulatus* (saltmarsh aster)

*Agalinis maritima* (seaside gerardia)

**Salt Flats and Salt Barrens**

*Monanthochloe littoralis* (shoregrass)

*Batis maritima* (saltwort)

*Lycium carolinianum* (Carolina wolfberry)

*Borrichia frutescens* (sea-oxeye)

*Salicornia virginica* (perennial glasswort)

**Fresh Marsh**

*Scirpus californicus* (California bulrush)

*Paspalum lividum* (longtom)

*Leptochloa uninervia* (Mexican spangletop)

*Echinochloa crusgalli* (barnyard grass)

*Pulchea purpurascens* (purple pluchea)

*Pistia stratiotes* (water-lettuce)

*Echinodorus cordifolius* (burhead)

*Sagittaria graminea* (grassy arrowhead)

Table 5. Typical plants identified at saline sites at Armand Bayou and Vicinity. (From Brown, 1985)

Plant Name	Common Name
<i>Panicum repens</i>	torpedograss
<i>Phragmites australis</i>	common reed
<i>Spartina</i> , all species	cordgrasses
<i>Sporobolus virginicus</i>	seashore dropseed
<i>Scirpus americanus</i>	American bulrush
<i>Scirpus maritimus</i>	saltmarsh bulrush
<i>Juncus roemerianus</i>	needlegrass rush
<i>Atriplex arenaria</i>	saltbush
<i>Salicornia bigelovii</i>	annual glasswort
<i>Suaeda linearis</i>	annual seepweed
<i>Sesuvium portulacastrum</i>	sea-purslane
<i>Opuntia lindheimeri</i>	Texas pricklypear
<i>Limonium nashii</i>	sea-lavender
<i>Sabatia arenicola</i>	sand rosegentian
<i>Cuscuta indecora</i>	showy dodder
<i>Ipomoea sagittata</i>	saltmarsh morning glory
<i>Heliotropium curassavicum</i>	seaside heliotrope
<i>Lycium carolinianum</i>	Carolina wolfberry
<i>Bacopa monnieri</i>	coastal waterhyssop
<i>Aster tenuifolius</i>	perennial saltmarsh aster
<i>Borrichia frutescens</i>	sea oxeye
<i>Iva frutescens</i>	big-leaf sumpweed
<i>Machaeranthera phyllocephala</i>	camphor daisy

Table 4. Typical plants found in grassflats, marshes, and transitional areas in the Galveston Bay area. (From White and others, 1985)

Unit	Scientific Name	Common Name	Unit	Scientific Name	Common Name
GRASS-FLAT (subaqueous marine grasses)	<i>Halodule beaudettei</i> <i>Ruppia maritima</i>	shoalgrass widgeongrass	BRACKISH-WATER MARSH (cont.)	<i>Juncus roemerianus</i> <i>Lycium carolinianum</i> <i>Sporobolus</i> spp. <i>Fimbristylis castanea</i> <i>Hydrocotyle</i> spp.	needle rush Carolina wolfberry dropseed fimbrystyle pennywort
SALT-WATER MARSH	<i>Spartina alterniflora</i> <i>Batis maritima</i> <i>Salicornia virginica</i> <i>Salicornia bigelovii</i> <i>Distichlis spicata</i> <i>Borrichia frutescens</i> <i>Monanthochloe littoralis</i> <i>Juncus roemerianus</i> <i>Suaeda</i> spp. <i>Lycium carolinianum</i> <i>Spartina patens</i> <i>Iva frutescens</i> <i>Iva angustifolia</i> <i>Limonium nashii</i> <i>Scirpus maritimus</i> <i>Sporobolus</i> spp. <i>Sesuvium portulacastrum</i> <i>Heliotropium curassavicum</i>	smooth cordgrass saltwort glasswort glasswort seashore saltgrass sea-oxeye shoregrass needle rush seablite or seepweed Carolina wolfberry gulf cordgrass marshhay cordgrass bigleaf sumpweed narrowleaf sumpweed sea-lavender salt-marsh bulrush dropseed sea purslane salt heliotrope	FRESH-WATER MARSH	<i>Spartina spartinae</i> <i>Typha latifolia</i> <i>Typha domingensis</i> <i>Scirpus americanus</i> <i>Scirpus californicus</i> <i>Paspalum lividum</i> <i>Eleocharis</i> spp. <i>Cyperus</i> spp. <i>Alternanthera philoxeroides</i> <i>Juncus</i> spp. <i>Ludwigia</i> spp. <i>Sagittaria</i> spp. <i>Pontederia</i> sp. <i>Polygonum</i> spp. <i>Phragmites australis</i> <i>Bacopa monnieri</i> <i>Echinodorus</i> spp. <i>Eichhornia crassipes</i> <i>Rhynchospora</i> spp. <i>Fimbristylis</i> spp. <i>Echinochloa crusgalli</i> <i>Leptochloa</i> spp. <i>Spartina patens</i> <i>Lemna</i> spp. <i>Hydrocotyle</i> spp. <i>Zizaniopsis miliacea</i> <i>Sesbania drummondii</i> <i>Baccharis halimifolia</i> <i>Cephalanthus occidentalis</i> <i>Salix nigra</i>	gulf cordgrass common cattail narrowleaf cattail three-square bulrush California bulrush longtom spikesedge flatsedge alligatorweed rush seedbox arrowhead pickerelweed knotweed common reed waterhyssop burhead water hyacinth beakrush fimbrystyle barnyard grass sprangletop marshhay cordgrass duckweed marsh pennywort southern wildrice rattlebush groundsel bush buttonbush black willow
BRACKISH-WATER MARSH	<i>Spartina spartinae</i> <i>Spartina patens</i> <i>Borrichia frutescens</i> <i>Distichlis spicata</i> <i>Monanthochloe littoralis</i> <i>Scirpus maritimus</i> <i>Scirpus americanus</i> <i>Scirpus californicus</i> <i>Scirpus olneyi</i> <i>Alternanthera philoxeroides</i> <i>Typha domingensis</i> <i>Typha latifolia</i> <i>Spartina cynosuroides</i> <i>Phragmites australis</i> <i>Eleocharis parvula</i> <i>Eleocharis</i> spp. <i>Cyperus</i> spp. <i>Echinochloa crusgalli</i> <i>Leptochloa</i> spp. <i>Bacopa monnieri</i> <i>Aster tenuifolius</i> <i>Aster subulatus</i> <i>Aster spinosus</i> <i>Paspalum lividum</i> <i>Paspalum vaginatum</i> <i>Setaria geniculata</i> <i>Zizaniopsis miliacea</i> <i>Solidago sempervirens</i> <i>Baccharis halimifolia</i> <i>Iva frutescens</i> <i>Iva angustifolia</i> <i>Iva annua</i> <i>Sesuvium portulacastrum</i> <i>Salicornia</i> spp. <i>Limonium nashii</i>	gulf cordgrass marshhay cordgrass sea-oxeye seashore saltgrass shoregrass salt marsh bulrush three-square bulrush California bulrush Olney bulrush alligatorweed narrowleaf cattail common cattail big cordgrass common reed dwarf spikerush spikerush flatsedge barnyard grass sprangletop coastal waterhyssop saline aster saltmarsh aster spiny aster longtom seashore paspalum knotroot bristlegrass giant cutgrass seaside goldenrod groundsel bush bigleaf sumpweed narrowleaf sumpweed seacoast sumpweed sea purslane glasswort sea-lavender	TRANSITIONAL AREAS	<i>Spartina spartinae</i> <i>Cynodon dactylon</i> <i>Borrichia frutescens</i> <i>Aster spinosus</i> <i>Paspalum monostachyum</i> <i>Paspalum lividum</i> <i>Panicum</i> spp. <i>Rhynchospora</i> spp. <i>Andropogon virginicus</i> <i>Andropogon glomeratus</i> <i>Iva annua</i> <i>Aristida</i> spp. <i>Setaria</i> spp. <i>Helianthus</i> spp. <i>Sorghum halepense</i> <i>Cassia fasciculata</i> <i>Cyperus</i> spp. <i>Eleocharis</i> <i>Scirpus</i> spp. <i>Croton</i> spp. <i>Spartina patens</i> <i>Baccharis halimifolia</i> <i>Sesbania drummondii</i>	gulf cordgrass bermudagrass sea-oxeye spiny aster guifdune paspalum longtom panicum beakrush broomsedge bluestem bushy bluestem seacoast sumpweed threeawn bristlegrass sunflower johnsongrass partridge pea flatsedge spikerush bulrush doveweed marshhay cordgrass groundsel bush rattlebush

Although the periodicity of inundations is irregular, wind tides have developed a relatively broad low marsh that includes species other than regularly flooded *Spartina alterniflora*. Above this level are higher marshes that are flooded less frequently.

Shaw and Fredine (1956) define a regularly flooded salt marsh and an irregularly flooded salt marsh. Cowardin and others (1979) used water-regime modifiers to denote the regularity of flooding (table 6). White and others (1985) used the terms proximal and distal (for salt-water marshes) to differentiate areas that are more frequently flooded because of lower elevations and proximity to estuarine water from those areas less frequently flooded because of higher elevations and distal locations with respect to estuarine water.

### Species Composition of Wetland Plant Communities, Galveston Bay System

To collect information on plant composition, wetland communities were surveyed at more than 150 sites around the Galveston Bay system; more than 135 sites are shown in figure 3, and are listed in appendices A and B. The Galveston Bay project area is defined by 30 USGS 7.5-minute quadrangle maps, although one additional map (Freeport) was included for the field surveys. The maps were assigned numbers from 1 to 31 to simplify numerical designations of the surveyed sites (fig. 4, table 7). Species composition at the various sites along with very brief descriptive notes on the relationship of the identified plant communities to topography (for example, high versus low zones) and local geographic features (such as roads or streams) are presented in appendix B.

Wetland plant communities in the Galveston Bay system include high and low categories of salt, brackish, and fresh marshes, and forested wetlands. Other environments include mud and sand flats, beaches and bars, submerged vascular vegetation, disturbed areas, and open water.

The most widely distributed wetland environments in the Galveston Bay system are marshes, the most extensive of which are brackish. Brackish marshes, as mapped by White and others (1985), compose roughly 65 to 70 percent of the marsh system in the Galveston Bay project area. Salt marshes are a distant second, composing roughly 25 to 30 percent. Fresh marshes make up the remaining 5 to 10 percent of the marsh system. Because many species can tolerate varying salinity regimes as well as water regimes, there is considerable overlap in the species composition of these marsh systems (table 8). The divergent plant communities in the project area are exemplified by the fresh marshes and swamps along the Trinity River which contrast sharply with the salt marshes that fringe Christmas Bay.

Because of the predominance of brackish and salt marshes in the project area, more than 60 percent of the field surveys were located in these marshes. Surveys of other environments ranged from approximately 8 percent in forested wetlands to about 5 percent in transitional areas (appendix A). With reference to all sites visited, the 15 most frequently encountered species, were headed by *Spartina patens* (marshhay or saltmeadow cordgrass) and *Distichlis spicata* (saltgrass) (table 9).

Each of the species in table 9 was observed at more than 20 sites, *Spartina patens* and *Distichlis spicata* occurred at more than 60 sites, and *Spartina alterniflora* (smooth cordgrass) at more than 40 sites. Other species listed as among the top 25 reported include *Solidago* spp., *Limonium nashii*, *Phragmites australis*, *Lycium carolinianum*, *Paspalum vaginatum*, and *Suaeda linearis*. These species plus those listed in table 9 are typical of the brackish and salt marsh systems.

Table 6. Water regime descriptions for wetlands used in the Cowardin and others (1979) classification system.

**Nontidal**

- (A) Temporarily flooded—Surface water present for brief periods during growing season, but water table usually lies well below soil surface. Plants that grow both in uplands and wetlands are characteristic of this water regime.
- (C) Seasonally flooded—Surface water is present for extended periods, especially early in the growing season, but is absent by the end of the growing season in most years. The water table is extremely variable after flooding ceases, extending from saturated to well below the ground surface.
- (F) Semipermanently flooded—Surface water persists throughout the growing season in most years. When surface water is absent, the water table is usually at or very near the land's surface.
- (H) Permanently flooded—Water covers land surface throughout the year in all years.

**Tidal**

- (L) Subtidal—The substrate is permanently flooded with tidal water.
- (M) Irregularly exposed—The land surface is exposed by tides less often than daily.
- (N) Regularly flooded—Tidal water alternately floods and exposes the land surface at least once daily.
- (P) Irregularly flooded—Tidal water floods the land surface less often than daily.
- (S)\* Temporarily flooded—Tidal
- (R)\* Seasonally flooded—Tidal
- (T)\* Semipermanently flooded—Tidal
- (V)\* Permanently flooded—Tidal

\*These water regimes are only used in tidally influenced, freshwater systems.







Figure 3. Location map of field survey sites. Inset maps are shown on figures 3a, b and c. See figure 4 and table 7 for identification of quadrangle maps on which sites are located.



(a) Inset 1 from figure 3.



(b) Inset 2 from figure 3.



(c) Inset 3 from figure 3.




Figure 4. Index map of quadrangles covering the Galveston Bay area (table 7).

Table 7. List of USGS 7.5-minute topographic maps that encompass the Galveston Bay project area. Quadrangle locations shown on figure 4.

Quadrangle Number	Southeast Corner Latitude (N)	Southeast Corner Longitude (W)	USGS Quadrangle Name
1	28° 52.5'	95° 15.0'	Freeport (south of project area)
2	29° 00.0'	95° 00.0'	San Luis Pass
3	29° 00.0'	95° 07.5'	Christmas Point
4	29° 00.0'	95° 15.0'	Oyster Creek
5	29° 07.5'	94° 52.5'	Lake Como
6	29° 07.5'	95° 00.0'	Sea Isle
7	29° 07.5'	95° 07.5'	Hoskins Mound
8	29° 15.0'	94° 37.5'	The Jetties
9	29° 15.0'	94° 45.0'	Galveston
10	29° 15.0'	94° 52.5'	Virginia Point
11	29° 15.0'	95° 00.0'	Hitchcock
12	29° 22.5'	94° 30.0'	Caplen
13	29° 22.5'	94° 37.5'	Flake
14	29° 22.5'	94° 45.0'	Port Bolivar
15	29° 22.5'	94° 52.5'	Texas City
16	29° 22.5'	95° 00.0'	Dickinson
17	29° 30.0'	94° 22.5'	High Island
18	29° 30.0'	94° 30.0'	Frozen Point
19	29° 30.0'	94° 37.5'	Lake Stephenson
20	29° 30.0'	94° 45.0'	Smith Point
21	29° 30.0'	94° 52.5'	Bacliff
22	29° 30.0'	95° 00.0'	League City
23	29° 37.5'	94° 30.0'	Oyster Bayou
24	29° 37.5'	94° 37.5'	Oak Island
25	29° 37.5'	94° 45.0'	Umbrella Point
26	29° 37.5'	94° 52.5'	Morgans Point
27	29° 37.5'	95° 00.0'	La Porte
28	29° 45.0'	94° 37.5'	Anahuac
29	29° 45.0'	94° 45.0'	Cove
30	29° 45.0'	94° 52.5'	Mont Belvieu
31	29° 45.0'	95° 00.0'	Highlands

Table 8. List of common plant species for various marshes based on field surveys. This list characterizes common wetland plants according to general frequencies of occurrence. Many species grow over a range of elevations and salinities and may occur in more than one class.

SALT MARSH	BRACKISH MARSH	FRESH MARSH
LOW	LOW	LOW
<i>Spartina alterniflora</i> <i>Juncus roemerianus</i> <i>Scirpus maritimus</i> <i>Scirpus olneyi</i> <i>Distichlis spicata</i> <i>Batis maritima</i> <i>Salicornia spp.</i>	<i>Scirpus olneyi</i> <i>Scirpus californicus</i> <i>Scirpus maritimus</i> <i>Scirpus americanus</i> <i>Typha spp.</i> <i>Alternanthera philoxeroides</i> <i>Crinum americanum</i> <i>Eleocharis spp.</i> <i>Paspalum vaginatum</i> <i>Bacopa monnieri</i> <i>Zizaniopsis miliacea</i> <i>Panicum dichotomiflorum</i>	<i>Typha spp.</i> <i>Sagittaria spp.</i> <i>Scirpus californicus</i> <i>Juncus spp.</i> <i>Scirpus americanus</i> <i>Zizaniopsis miliacea</i> <i>Alternanthera philoxeroides</i> <i>Eichhornia crassipes</i> <i>Eleocharis spp.</i> <i>Cyperus articulatus</i> <i>Ludwigia spp.</i> <i>Pontederia spp.</i>
HIGH	HIGH	HIGH
<i>Spatina patens</i> <i>Distichlis spicata</i> <i>Spartina spartinae</i> <i>Borrichia frutescens</i> <i>Iva frutescens</i> <i>Batis maritima</i> <i>Salicornia virginica</i> <i>Salicornia bigelovii</i> <i>Monanthochloe littoralis</i> <i>Limonium nashii</i> <i>Lycium carolinianum</i> <i>Aster tenuifolius</i> <i>Suaeda linearis</i> <i>Heliotropium curassavicum</i>	<i>Spartina patens</i> <i>Distichlis spicata</i> <i>Spartina spartinae</i> <i>Spartina cynosuroides</i> <i>Borrichia frutescens</i> <i>Paspalum lividum</i> <i>Paspalum vaginatum</i> <i>Phragmites australis</i> <i>Panicum virgatum</i> <i>Echinochloa crusgalli</i> <i>Leptochloa sp.</i> <i>Scirpus americanus</i> <i>Aster subulatus</i> <i>Aster tenuifolius</i> <i>Hydrocotyle spp.</i> <i>Fimbristylis spp.</i> <i>Setaria geniculata</i>	<i>Polygonum sp.</i> <i>Phragmites australis</i> <i>Echinochloa crusgalli</i> <i>Cyperus articulatus</i> <i>Cyperus spp.</i> <i>Paspalum lividum</i> <i>Scirpus americanus</i> <i>Leptochloa sp.</i> <i>Panicum spp.</i> <i>Spartina spartinae</i> <i>Spartina pectinata</i>

Table 9. Species most frequently observed at survey sites in the study area listed in order by number of sites at which plant was reported.

Plant Name	Common Name
<i>Spartina patens</i>	saltmeadow cordgrass
<i>Distichlis spicata</i>	seashore saltgrass
<i>Spartina alterniflora</i>	smooth cordgrass
<i>Batis maritima</i>	saltwort
<i>Salicornia</i> spp.	glasswort
<i>Iva frutescens</i>	big-leaf sumpweed
<i>Spartina spartinae</i>	gulf cordgrass
<i>Borrichia frutescens</i>	sea oxeye
<i>Juncus roemerianus</i>	needlegrass rush
<i>Aster</i> spp.	aster
<i>Typha</i> spp.	cattail
<i>Scirpus maritimus</i>	saltmarsh bulrush
<i>Monanthochloe littoralis</i>	shoregrass

## Wetland Indicator Status of Prevalent Plants at Survey Sites

The scientific and common names of plant species identified at field survey sites are presented in table 10. Each species is classified in terms of its wetland indicator status for Region 6, which includes Texas, and for the United States. The indicator status is based on the "National List of Plant Species That Occur in Wetlands: 1988, Texas" (Reed, 1988). In addition, the habit for each species as defined in the list (Reed, 1988) is presented in table 10.

Of the species identified at the survey sites (fig. 3), about 34 percent are classified as obligate (OBL) wetland plants, which means that under natural conditions these plants occur in wetlands with an estimated probability of 99 percent. Among the species are those typically found in wetter conditions, for example, those characterizing topographically low salt, brackish, and fresh marshes (table 8). Such species include *Spartina alterniflora*, *Juncus roemerianus*, *Scirpus californicus*, *Scirpus olneyi*, *Eleocharis* spp., *Bacopa monnieri*, *Typha* spp., *Alternanthera philoxeroides*, and *Sagittaria* spp., among others.

Approximately 37 percent of the species listed (table 10) are classified as Facultative Wetland plants (FACW, FACW+, and FACW-). These species usually occur in wetlands or have an estimated probability of 76–99 percent of occurring in wetlands; but occasionally they occur in nonwetland areas. Included, for example, are those species that typically define topographically higher marshes (table 8) such as *Borrichia frutescens*, *Spartina patens*, *Spartina spartinae*, *Phragmites australis*, *Echinochloa crusgalli*, *Hydrocotyle bonariensis*, *Heliotropium curassavicum*, and *Aster spinosus*. Some Facultative Wetland plants (for instance, *Paspalum vaginatum*) may also occur in wetter, typically low marshes.

About 19 percent of the listed species are classified as Facultative (FAC). These species are equally likely to occur in wetlands or nonwetlands (estimated probability 34–66 percent). Such species include grasses like *Setaria geniculata*, *Paspalum urvillei*, and *Panicum repens*. Many trees such as *Carya illinoensis*, *Celtis laevigata*, *Pinus taeda*, and *Ulmus crassifolia* also are listed as FAC plants.

Only 7 percent of the plants listed are classified as Facultative Upland (FACU). These species are usually not found in wetlands; their estimated probability of occurring in wetlands is 1–33 percent. Such species include the grasses *Cynodon dactylon*, *Andropogon virginicus*, and *Eragrostis spectabilis*.

## Wetland Plant Communities

In the following discussion of coastal wetland communities in the Galveston Bay system, marshes are subdivided into salt, brackish, and fresh communities to assist in the discussions of vegetation composition. A lack of long-term field data precludes the establishment of definite salinity values for these units. Because some plant species can tolerate a relatively large range in salinities (Penfound and Hathaway, 1938; Chabreck, 1972), species tend to overlap between the fresh- and the brackish-marsh communities, and the brackish- and the salt-marsh communities. Overlap between communities also occurs between topographically high and low marshes. Some species can tolerate a range in water regimes, or frequencies of inundation, and therefore may occur in wet, low areas as well as in high, dryer areas.

Mapping of wetlands and aquatic habitats by the USFWS follows the classification by Cowardin and others (1979). As mentioned previously, in general terms emergent vegetation in the

Table 10. Plant list compiled from field survey sites; wetland indicator status of plants from Reed (1988).

Emergent spp.	Common Name	Status	Status	Habit
			Reg. 6	
<i>Acacia angustissima</i>	Fern acacia	not listed		
<i>Alternanthera philoxeroides</i>	Alligator weed	OBL	OBL	PIEF
<i>Ambrosia psilostachya</i>	Western ragweed	FAC-	FACU-, FAC	PNF
<i>Ambrosia trifida</i>	Giant ragweed	FAC	FAC,FACW	ANF
<i>Andropogon glomeratus</i>	Bushy bluestem	FACW+	FACW,OBL	PNG
<i>Andropogon virginicus</i>	Broom-sedge	FACU+	FACU,FAC	PNG
<i>Aristida sp.</i>	Three-awn	FACW-to FACU		
<i>Arundo donax</i>	Giant reed	FAC+	FACU-, FACW	PIG
<i>Aster spinosus</i>	Spiny aster	FACW-	FAC, FACW	PNF
<i>Aster subulatus</i>	Annual saltmarsh aster	OBL	FACW,OBL	ANF
<i>Aster tenuifolius</i>	Perennial saltmarsh aster	OBL	OBL	PNF
<i>Baccharis halimifolia</i>	Eastern B., Sea-myrtle	FACW-	FAC,FACW	NS
<i>Bacopa monnieri</i>	Coastal waterhyssop	OBL	OBL	PNF
<i>Batis maritima</i>	Saltwort	OBL	OBL	NS
<i>Borrichia frutescens</i>	Sea oxeye	FACW+	FACW+,OBL	NS
<i>Cardiospermum halicacabum</i>	Balloon vine	FAC	FACU,FAC	AIF
<i>Carya aquatica</i>	Water hickory	OBL	OBL	NT
<i>Carya illinoensis</i>	Pecan hickory	FAC+	FACU,FACW	NT
<i>Celtis laevigata</i>	Sugar-berry	FAC	UPL,FACW	NT
<i>Cephalanthus occidentalis</i>	Common buttonbush	OBL	OBL	NT
<i>Crinum americanum</i>	Swamp lily	OBL	OBL	PNF
<i>Cynodon dactylon</i>	Bermuda grass	FACU+	FACU,FAC	PIG
<i>Cyperus articulatus</i>	Jointed flatsedge	OBL	OBL	PNG
<i>Cyperus elegans</i>	Sticky flatsedge	FACW-	FACW-,FACW	PNG
<i>Cyperus oxylepis</i>	Sharp-scale flatsedge	FACW	FACW	PNEG
<i>Cyperus virens</i>	Green flatsedge	FACW	FACW	PNEG
<i>Dichromena colorata</i>	Starrush whitetop	FACW	FACW	PNG
<i>Distichlis spicata</i>	Seashore saltgrass	FACW+	FAC+,FACW+	PNG
<i>Desmodium canadense</i>	Tickclover	FAC	FACU, FAC	PNF
<i>Echinochloa crusgalli</i>	Barnyard grass, water millet	FACW-	FACU,FACW	AIG
<i>Eichhornia crassipes</i>	Common water-hyacinth	OBL	OBL	PNE/F (I-Ck.Lst.)
<i>Eleocharis parvula</i>	Dwarf spikesedge	OBL	OBL	PNG
<i>Eleocharis cellulosa</i>	Gulf Coast spikesedge	OBL	OBL	PNG
<i>Eleocharis microcarpa</i>	Small-fruit spikerush	OBL	OBL	ANEGL
<i>Eleocharis quadrangulata</i>	Squarestem spikesedge	OBL	OGL	PNEG
<i>Eleocharis lanceolata</i> ?	Lanceleaf spikesedge	OBL	OBL	PNG
<i>Eleocharis sp.</i>	Spikesedge	OBL?	OBL?	PIG?
<i>Eragrostis spectabilis</i>	Purple lovegrass	FACU-	UPL,FACU	PNG
<i>Eustachys petraea</i>	Pinewoods finger grass	FAC-	FACU-, FAC	NG
<i>Fimbristylis castanea</i>	Marsh fimbry	OBL	OBL	PNEG
<i>Fraxinus caroliniana</i>	Carolina ash	OBL	OBL	NETS
<i>Fraxinus pennsylvanica</i>	Green ash	FACW-	FAC,FACW	NT
<i>Gleditsia aquatica</i>	Water locust	OBL	OBL	NET
<i>Heliotropium curassavicum</i>	Seaside heliotrope	FACW	FACW,OBL	APISF
<i>Hydrocotyle bonariensis</i>	Coastal plain penny-wort	FACW	FACW	PNF
<i>Hymenocallis caroliniana</i>	Carolina spider lily	FACW	FACW	PNF
<i>Ilex vomitoria</i>	Yaupon	FAC-	FAC-, FAC	NST
<i>Ipomea sp.</i>	Morning Gloria	FAC?	FAC?	?
<i>Iva annua</i>	Annual sumpweed, marsh-elder	FAC	FAC	AIF
<i>Iva angustifolia</i>	Narrowleaf sumpweed	Not listed		
<i>Iva frutescens</i>	Big-leaf sumpweed	FACW	FACW,FACW+	PNH\$F
<i>Juncus roemerianus</i>	Needlegrass rush	OBL	OBL	PNG
<i>Lemna sp.</i>	Duckweed	OBL	OBL	PN/F
<i>Limonium nashii</i>	Sea-lavender	NA*	OBL	PNF
<i>Liquidambar styraciflua</i>	Sweet gum	FAC	FAC,FACW	NT
<i>Lolium perenne</i>	Perennial ryegrass	FACU	FACU-, FAC	PIG
<i>Lycium carolinianum</i>	Carolina wolf-berry	FACW	VACW	NS
<i>Machaeranthera phyllocephala</i>	Camphor daisy	FACW	FACU,FACW	ANF
<i>Medicago minima</i>	Small medic	Not listed		
<i>Monanthochloe littoralis</i>	Shoregrass	OBL	OBL	PNEG
<i>Nelumbo lutea</i>	American lotus	OBL	OBL	PNZ/F
<i>Panicum dichotomiflorum</i>	Fall panic grass	FACW	FAC,FACW	ANG
<i>Panicum hiatus</i>	Gaping panicum	FACW-	FACW-,OBL	PNG

Table 10 (cont.)

Emergent spp.	Common Name	Status, Reg. 6	Status, Nat.	Habit
<i>Panicum virgatum</i>	Switchgrass	FACW	FAC,FACW	PNG
<i>Panicum repens</i>	Torpedograss	FAC+	FAC+, FACW-	PIG
<i>Parkinsonia aculeata</i>	Retama	FACW-	FAC-, FACW	NT
<i>Paspalum floridanum</i>	Florida paspalum	FACW-	FACW-, FACW	PNG
<i>Paspalum lividum</i>	Longtom	OBL*	OBL	PNEG
<i>Paspalum monostachyum</i>	Gulfdune paspalum	FACW+	FACW,FACW+	PNG
<i>Paspalum urvillei</i>	Vasey grass	FAC	FAC	PIG
<i>Paspalum vaginatum</i>	Seashore paspalum	FACW+	FACW,OBL	PNG
<i>Phragmites australis</i>	Common reed	FACW	FACW,FACW+	PNEG
<i>Phyla lanceolata</i>	Lance leaf frog fruit	FACW	FACW,OBL	PNF
<i>Physostegia intermedia</i>	Intermediate Lionsheart	OBL	FACW-, OBL	PNF
<i>Pinus taeda</i>	Loblolly pine	FAC-	UPL,FAC	NT
<i>Planera aquatica</i>	Water elm	OBL	OBL	NET
<i>Polygonum hydropiperoides</i>	Swamp smartweed	OBL	OBL	PNEF
<i>Polygonum ramosissimum</i>	Bushy knotweed	FACW	FACU-, FACW	ANF
<i>Quercus phellos</i>	Willow oak	FACW	FAC+, FACW	NT
<i>Quercus falcata</i>	Southern red oak	FACU	FACU-, FACU	NT
<i>Quercus nigra</i>	Water oak	FAC+	FAC,FACW	NT
<i>Quercus virginiana</i>	Live oak	FACU+	FACU,FACU+	NT
<i>Sabal minor</i>	Dwarf palmetto	FACW	FACW	NST
<i>Sagittaria latifolia</i>	Coastal arrow-head	OBL	OBL	PNEF
<i>Salicornia bigelovii</i>	Annual glasswort	OBL*	OBL	ANESF
<i>Salicornia virginica</i>	Perennial glasswort	OBL*	OBL	PNESF
<i>Salix nigra</i>	Black willow	FACW+	UPL,OBL	NT
<i>Sapium sebiferum</i>	Chinese tallow	FACU+	FACU+,FAC	IT
<i>Scirpus americanus</i>	Olney's (American) bulrush	OBL	OBL	PNEG
<i>Scirpus californicus</i>	California bulrush	OBL	OBL	PNEG
<i>Scirpus maritimus</i>	Saltmarsh bulrush	NI	OBL	PNEG
<i>Scirpus olneyi (S. americanus)</i>	Olney's bulrush	OBL	OBL	PNEG
<i>Sesbania drummondii</i>	Drummond's rattle-bush	FACW	FACW	NSH
<i>Sesuvium portulacastrum</i>	Sea-purslane	FACW	FACW	PNSF
<i>Setaria geniculata</i>	Knotroot bristlegrass	FAC	FAC	PNG
<i>Setaria magna</i>	Giant bristlegrass	FACW	FACW,FACW+	ANEG
<i>Sisyrinchium exile</i>	Yellow blue-eyed grass	FACW	FAC, FACW-	AIF
<i>Solidago altissima</i>	Tall goldenrod	FACU	FACU-, FACU+	PNF
<i>Solidago sempervirens</i>	Seaside golden-rod	FACW-	FACW-, FACW	PNSF
<i>Spartina spartinae</i>	Gulf cordgrass	FACW+	FACW+,OBL	PNG
<i>Spartina alterniflora</i>	Smooth cordgrass	OBL	OBL	PNEG
<i>Spartina cynosuroides</i>	Big cordgrass	OBL	OBL	PNEG
<i>Spartina patens</i>	Saltmeadow cordgrass	FACW	FACW,OBL	PNG
<i>Spartina pectinata</i>	Prairie cordgrass	FACW+	FACW,OBL	PNG
<i>Spiranthes ovalis</i>	October ladies'-tresses	FAC*	FAC	PNF
<i>Sporobolus virginicus</i>	Seashore dropseed	FACW+	FACW+	PNG
<i>Suaeda linearis</i>	Annual seepweed	OBL	OBL	ANEF
<i>Tamarix gallica</i>	Salt cedar	FACW	FAC, FACW	IT
<i>Taxodium distichum</i>	Bald cypress	OBL	OBL	NET
<i>Teucrium cubense</i>	Small coast germander	FAC+	UPL, FACW	APNF
<i>Typha spp.</i>	Cattail	OBL	OBL	PNEF
<i>Ulmus americana</i>	American elm	FAC	FAC,FACW	NT
<i>Ulmus crassifolia</i>	Cedar elm	FAC	FAC	NT
<i>Zizaniopsis miliacea</i>	Marsh millet, giant cutgrass	OBL	OBL	PNG

Table 10 (cont.)

Habitat symbols	Characteristic or life form			
A	Annual			
E	Emergent			
F	Forb			
/	Floating			
G	grass			
GL	Grass like			
H	Partly woody			
HS	Half shrub			
I	Introduced			
N	Native			
P	Perennial			
S	Shrub			
Z	Submerged			
\$	Succulent			
T	Tree			
V	Herbaceous vine			
WV	Woody vine			
NA	No agreement by regional panel			
*	Tentative assignment based on limited information			
**	More frequently found in wetland			
**	Less frequently found in wetland			
ABBREV.	INDICATOR CATEGORY	DESCRIPTION		
OBL	Obligate wetland	Occur almost always (est. prob. >99%) under natural conditions in wetlands.		
FACW	Facultative wetland	Usually occur in wetlands (est. prob. 76-99%), but occasionally found in nonwetlands.		
FAC	Facultative	Equally likely to occur in wetlands or nonwetlands (est. prob. 34-66%).		
FACU	Facultative upland	Usually occur in nonwetlands (est. prob. 67-99%), but occasionally found in wetlands (e.p. 1-33%).		
UPL	Obligate upland	Occur in wetlands in another region, but occur almost always (e.p. >99%) under natural conditions in nonwetlands		

Estuarine system corresponds to salt and brackish marshes and emergent vegetation in the Palustrine system corresponds to fresh marshes. Water regimes used as modifiers in classifying and mapping wetlands help define high and low wetlands (table 6).

### *Salt-Marsh Community*

Salt marshes were examined principally on Follets and Galveston Islands, and Bolivar Peninsula, along the inland margin of West Bay, near Texas City, and at Houston and Smith Points (figs. 5 through 9). Prevalent species in the salt-marsh community include *Spartina alterniflora* (smooth cordgrass), *Batis maritima* (saltwort), *Distichlis spicata* (saltgrass), *Salicornia virginica* and *Salicornia bigelovii* (glasswort), *Borrichia frutescens* (sea-oxeye), *Monanthochloe littoralis* (shoregrass), *Juncus roemerianus* (needlegrass rush or blackrush), *Suaeda linearis* (seepweed), *Scirpus maritimus* (salt-marsh bulrush), *Limonium nashii* (sea-lavender), *Aster tenuifolius* (perennial saltmarsh aster), and *Lycium carolinianum* (Carolina wolfberry). At higher elevations, *Spartina patens* (marshhay or saltmeadow cordgrass) and *Spartina spartinae* (Gulf cordgrass) occur, although these species are more common in brackish marshes. *Iva frutescens* (big-leaf sumpweed) is locally abundant at higher elevations such as along natural levees.

The low-salt-marsh community is dominated by *Spartina alterniflora*, which lives in the intertidal zone (fig. 5). Species intermixed most frequently with *Spartina alterniflora* along the upper part of the intertidal zone include *Batis maritima* (fig. 6), *Distichlis spicata* (fig. 7), *Scirpus maritimus*, *Juncus roemerianus*, and *Salicornia virginica*.

Wind-tidal sand flats are common features in some areas, especially on the barrier islands (fig. 10). Although algal mats are abundant in these areas, the flats are generally barren of emergent vegetation because of intermittent salt-water flooding and subsequent evaporation—a process that concentrates salts and inhibits the growth of most plants. Soil salinities on the flats can reach concentrations high enough to kill *Spartina alterniflora* and *Spartina patens* (Webb, 1983). The flats may locally have scattered salt-marsh vegetation. Common plant species are *Salicornia virginica*, *Salicornia bigelovii*, *Monanthochloe littoralis*, and *Batis maritima* (fig. 10). Zonation of some salt-marsh species is well defined by elevation transects at Smith Point (fig. 11), in the Brazoria National Wildlife Refuge (fig. 12), and other locations (appendix C).

The salt-marsh community corresponds in general terms to salt marshes (and locally, salt flats) defined by Shaw and Fredine (1956), Fisher and others (1972, 1973), Gosselink and others (1979), and White and others (1985) (table 4), and to saline wetland species identified by Lazarine (n.d.). In accordance with the classification of wetlands by Cowardin and others (1979), this community is designated (down to class) as estuarine, intertidal, emergent wetland (E<sub>2</sub>EM). The water regime modifier, "regularly flooded" (N), is used most frequently to identify low salt marshes; the modifier, "irregularly flooded" (P), is used to define higher marshes (table 6). (The classification by Cowardin and others [1979] has provisions for going beyond the class level and designating species dominance type, water chemistry, and human modifications; examples of the classification given here, however, will be only down to class and water regime.)

### *Brackish-Marsh Community*

The brackish-marsh community is transitional between salt marshes and fresh marshes. These areas are affected both by storm-tidal flooding from bay-estuary-lagoon and Gulf waters and by fresh-water inundation from rivers, precipitation and runoff, or ground water. Because the



Figure 5. Low salt-marsh community of *Spartina alterniflora* and open water on the inland margins of Jones Bay (east end of West Bay). Site No. 10-7, Virginia Point Quad. View is toward Galveston Island.




Figure 6. Salt-marsh community on Follets Island. *Batis maritima*, in foreground, intergrades with *Spartina alterniflora*, in background. Site No. 3-3, Christmas Point Quad. View is landward. See survey line at this site in appendix C.




Figure 7. Low salt-marsh community inland from West Bay. *Distichlis spicata* and scattered *Spartina alterniflora* are in the foreground. *Spartina alterniflora* becomes dominant as elevation decreases in distance. *Scirpus maritimus* is abundant on the margins of the tidal pond on the right; the dark assemblage along the margins of ponds in the upper left is *Juncus roemerianus*. Site No. 10-3, Virginia Point Quad. See survey line at this site in appendix C.




Figure 8. Salt-marsh community on the bayward margin of Bolivar Peninsula. *Spartina patens* and *Distichlis spicata* intergrade with *Scirpus maritimus*. In distance *Spartina alterniflora* is dominant in more regularly flooded areas. Site No. 18-1, Frozen Point Quad. View is toward East Bay.



Figure 9. Salt-marsh community at Houston Point. *Spartina alterniflora* dominates the low marsh in this area and intergrades with *Distichlis spicata* along higher margins. Species in the high marsh include *Spartina patens*, *Aster* sp., *Borrichia frutescens*, *Spartina spartinae*, *Iva frutescens*, and *Lycium carolinianum*. Site No. 26-7, Morgans Point Quad. View is inland (NW).



Figure 10. Salt marsh/sand flat community on Follets Island. Species include *Batis maritima*, *Monanthochloe littoralis*, and *Salicornia* spp. Site No. 3-3, Christmas Point Quad. View is southwestward, roughly parallel to island. See survey line for this site in appendix C.





Figure 11. Profile of salt marsh at Smith Point showing relative elevations of plant communities.  
Site No. 20-1.



#### PLANT COMMUNITIES AND GEOMORPHIC FEATURES ALONG PROFILE

BM to 2	<i>Spartina spartinae</i> (80%), <i>Setaria geniculata</i> , <i>Aster sp.</i> , <i>Iva annua</i> , others (20%)
2 to 4	<i>Spartina spartinae</i> (90%)
4 to 5	Fault Scarp
5 to 6	Mixed flat and emergent vegetation <i>Monanthochloe-Salsicornia-Batis</i>
6 to 7	<i>Distichlis spicata</i> (90%), <i>Salsicornia sp.</i> (10%)

Figure 12. Profile of brackish marsh in the Brazoria National Wildlife Refuge showing relative elevations of plant communities. Site No. 3-1.

brackish-marsh community encompasses a range in salinities from near fresh to near saline, the vegetation types cover a broad spectrum. Species range from those typical of saline marshes to those that occur in fresh marshes.

Areas in which brackish-marsh surveys were conducted included the Brazoria National Wildlife Refuge (figs. 13 and 14), Anahuac National Wildlife Refuge and near High Island (figs. 15 and 16), Galveston and Follets Islands (figs. 17 and 18), and Trinity River delta (figs. 19 and 20). Among the dominant species in topographically higher areas of this community are *Spartina patens*, *Spartina spartinae*, *Borrichia frutescens*, *Phragmites australis* (common reed), *Solidago sempervirens* (seaside goldenrod), *Panicum virgatum* (switchgrass) and *Spartina cynosuroides* (big cordgrass). Other prevalent species, most of which occur in lower, wetter areas (relative to the cordgrasses) include *Scirpus maritimus*, *Scirpus olneyi* (Olney bulrush) (fig. 15), *Juncus roemerianus*, *Typha* spp. (cattail), *Paspalum vaginatum* (seashore paspalum), *Scirpus americanus* (three-square bulrush), *Alternanthera philoxeroides* (alligatorweed), *Eleocharis* spp. (spikesedges), *Bacopa monnieri* (coastal waterhyssop), *Echinochloa crusgalli* (barnyard grass or water millet), and *Aster tenuifolius* and *Aster subulatus* (saline and saltmarsh aster), among others. *Spartina alterniflora* also occurs locally in the brackish-marsh community (fig. 13). Zonation of various species with respect to elevation are illustrated by marsh profiles on the Trinity River delta, and in the Brazoria, and Anahuac National Wildlife Refuges (appendix C). There are considerable differences in brackish marsh composition in the Brazoria and Anahuac National Wildlife Refuges (figs. 13 and 15) compared to brackish marshes in the Trinity River delta (figs. 19 and 20). In general, the Trinity River delta, which has extensive areas of *Alternanthera philoxeroides* (fig. 19) and other species occurring in fresher areas (fig. 20), is toward the fresh end of the brackish salinity spectrum.

The brackish-marsh community corresponds, generally, with the coastal salt meadows (grading into fresh marshes) defined by Shaw and Fredine (1956), the brackish (closed) and brackish- to fresh-water marsh by Fisher and others (1972, 1973), the brackish and intermediate marsh by Gosselink and others (1979), and the brackish marsh by Harcombe and Neaville (1977) (table 2) and White and others (1985) (table 4). In the classification system of Cowardin and others (1979), this community is generally designated (down to class) as estuarine, intertidal, emergent wetland (E<sub>2</sub>EM). Water regimes are generally the same as for the salt marshes—regularly flooded (N) (low marshes) and irregularly flooded (P) (high marshes).

*Spartina spartinae* is a common species in brackish marshes (fig. 14). Because of its tendency to occur mostly in topographically higher areas, it has been placed in the marsh, transitional (occurring between wetlands and uplands), and prairie communities by various researchers. It occurs in many areas in conjunction with *Spartina patens*, becoming more predominant and extensive (relative to *Spartina patens*) south of the Galveston Bay area along the Texas coast. Tharp (1926) listed *Spartina spartinae* as a dominant species in the coastal marsh community, but also included it as part of a coastal prairie-marsh-transition community. McAtee (1976) noted that *Spartina spartinae* flourishes at an elevation between lowland marshes and higher uplands, and apparently requires periodic inundation. The U.S. Army Corps of Engineers, which has jurisdictional responsibilities for wetlands, considers it to be a transitional species (Lazarine, n.d.). Many classifications place it in wetlands, transitional areas, and prairie grasslands (Fisher and others, 1972, 1973; Correll and Correll, 1975; White and others, 1985), presumably depending on associated plants and soil-moisture conditions reflecting inundation frequency. In the list of wetland plants of Texas (Reed, 1988), *Spartina spartinae* is categorized as usually found in wetlands, but occasionally found in nonwetlands. Harcombe and Neaville (1977) place it in their cordgrass prairie unit (table 2), but also list it in a checklist of marsh species and note that it probably once was more extensive (in Chambers County) as an intermediate type between upland prairie and brackish marsh. Fleetwood (n.d.) reported that *Spartina spartinae* was the predominant species in his salty prairie community.



Figure 13. Brackish-marsh community in the Brazoria National Wildlife Refuge southwest of Hoskins Mound. Although dominant species are *Spartina patens* and *Distichlis spicata*, *Spartina alterniflora* occurs along the tidal channel. *Ruppia maritima* (widgeongrass) occurs in the channel. Site No. 3-2, Christmas Point and Oyster Creek Quads. View is landward. This site is on the Oyster Creek Quad, at the west end of the survey line at this site. See survey line in appendix C.



Figure 14. Brackish-marsh community in the Brazoria National Wildlife Refuge east of Hoskins Mound. *Spartina spartinae* is dominant in the foreground and *Juncus roemerianus* in the background. Site No. 7-1, Hoskins Mound Quad. Several elevation surveys were conducted in this area (appendix C).




Figure 15. High and low brackish-marsh communities in the Anahuac National Wildlife Refuge. The high-marsh community is dominated by *Spartina patens* and *Distichlis spicata* in foreground, and the low marsh by *Scirpus olneyi* in the center of the photograph. Site No. 18-9, Frozen Point Quad.; view is landward (NW). See survey line in appendix C.



Figure 16. Brackish-marsh community dominated by *Spartina patens*, west of High Island. Site No. 17-1, High Island Quad.; view is landward (NW).



Figure 17. Brackish-marsh community in a swale on Galveston Island. A dike in this area separates a fresher from a more saline assemblage. Species in the fresher area include *Bacopa monnieri*, *Spartina patens*, *Borrichia frutescens*, and probably *Paspalum vaginatum* and *Scirpus californicus*. The more saline community (not shown in the photograph) includes *Distichlis spicata*, *Batis maritima*, and *Salicornia* spp. Site No. 10-16, Virginia Point Quad.; view is southwest.



Figure 18. Brackish- to fresh-marsh community in a depression on Follets Island, gulfward of highway. Species include *Typha* sp., *Paspalum vaginatum*, *Scirpus americanus*, and *Phragmites australis*. Bayward, across the highway, a salt-marsh community occurs. Site No. 3-12, Christmas Point Quad.; view is gulfward.



Figure 19. Brackish-marsh community on the Trinity River delta. This area is dominated by *Alternanthera philoxeroides* with local patches of *Crinum americanum* (swamp lily). *Phragmites australis* and scattered trees and shrubs line the natural levee along the Trinity River to the left of the photograph. Site No. 28-11, Anahuac Quad.; view is down river toward Trinity Bay.




Figure 20. Brackish-marsh community on the Trinity River delta near the delta/bay margin. Species include *Scirpus olneyi*, *Panicum dichotomiflorum*, *Echinochloa crusgalli*, *Bacopa monnieri*, and *Eleocharis parvula*. This dynamic area of the delta has a dramatic seasonal change in vegetation as described by White and Calnan (1990). Site No. 28-1, Anahuac Quad.; view is westward.

Brackish marshes dominate the coastal marsh community between High Island and Trinity Bay (fig. 2). They are also widely distributed along the lower reaches of the Trinity bay-head delta below Interstate Highway 10, inland from parts of West Bay, and inland of the Intracoastal Waterway in the Christmas Bay area. They occur in swales and intergrade with salt marshes and sand flats on Galveston Island (fig. 17) and Bolivar Peninsula.

### *Fresh-Marsh Community*

Surveys of fresh to intermediate marshes were conducted along the Trinity (figs. 21 and 22) and San Jacinto Rivers (fig. 23), and at other inland sites (figs. 24 and 25). Environments in which fresh marshes occur are generally beyond the limits of salt-water flooding except perhaps locally during hurricanes. The fresh-water influence from rivers, precipitation, runoff, and ground water is sufficient to maintain a fresher water vegetation community (although many species also occur in brackish marshes) consisting of species such as *Typha* spp., *Phragmites australis*, *Zizaniopsis miliacea* (marsh millet or giant cutgrass), *Sagittaria falcata* (coastal arrowhead), *Scirpus californicus*, *Eleocharis quadrangulata* (squarestem spikesedge) and other species of *Eleocharis*, *Cyperus* spp. (flatsedges), *Bacopa monnieri*, *Alternanthera philoxeroides*, *Paspalum lividum* (longtom), and *Eichhornia crassipes* (water hyacinth) in lower, wetter areas. Topographically higher areas generally include such species as *Phragmites australis*, *Paspalum* spp., *Polygonum* spp. (smartweeds), *Panicum* spp. (panic grasses), *Rhynchospora* spp. (beakrushes), and *Aster spinosus* (spiny aster). Shrubs such as *Sesbania drummondii* (rattlebush) are scattered around the margins of some fresh marshes and are locally abundant. Some species that are more common in brackish marshes such as *Spartina spartinae* may also occur in fresh marshes. Harcombe and Neaville (1977) used *Spartina patens* as an indicator of brackish conditions in differentiating brackish from fresh marshes.

The fresh-marsh community corresponds to the deep fresh and shallow fresh marshes of Shaw and Fredine (1956), inland fresh-water marsh and, locally, brackish- to fresh-water marsh of Fisher and others (1972, 1973), and fresh marsh of Fleetwood (n.d.), Harcombe and Neaville (1977) (table 2), Gosselink and others (1979), and White and others (1985) (table 4). Following the classification by Cowardin and others (1979) this community would be designated (down to class) as palustrine, emergent wetland (PEM) in areas where persistent emergent vegetation such as *Typha* spp. is present, and palustrine, aquatic bed (PAB) where floating vascular plants such as *Eichhornia crassipes* occur. A variety of water regimes can be applied under the Cowardin system (table 6). Low fresh marshes are usually characterized by the "semipermanently flooded" (F) or "seasonally flooded" (C) water regimes, and higher marshes by the "temporarily flooded" (A) regime, and occasionally the seasonally flooded regime. Fresh-water marshes in tidally influenced areas, have a different set of modifiers ranging from "semipermanently flooded—tidal" (T) to "temporarily flooded—tidal" (S) (table 6). These regimes are applicable along river systems, for example, and have been applied to some fresh marshes in the Trinity River delta.

Fresh marshes occur inland along river or fluvial systems and in upland basins and depressions on the mainland and perhaps locally on the barrier islands (fig. 18). Upstream along the river valleys of the Trinity and San Jacinto Rivers, salinities decrease and fresh marshes intergrade with and replace brackish marshes (figs. 21 through 23). Fresh marshes also occur locally in swales on the modern barrier islands and on the Pleistocene barrier strandplain, and in abandoned channels and courses of the Pleistocene fluvial-deltaic systems (fig. 2).



Figure 21. Fresh-marsh community in the Trinity River valley north of Interstate Highway 10. Species include *Cyperus articulatus* (jointed flatsedge), *Sagittaria falcata*, *Scirpus californicus*, *Zizaniopsis miliacea*, and *Alternanthera philoxeroides*. Site No. 29-3, Cove Quad.; view is westward.



Figure 22. Fresh- to brackish-marsh community on the Trinity River delta near Old River Lake. Species include *Zizaniopsis miliacea*, *Sagittaria falcata*, and *Alternanthera philoxeroides*. Site No. 29-2, Cove Quad.; view is northwest.



Figure 23. Fresh-marsh and forested-wetland communities in the San Jacinto River valley. Marsh species include *Typha* sp., *Scirpus californicus*, and *Eleocharis* sp. Site No. 31-S, Highlands Quad.; view is southeastward.




Figure 24. Fresh-marsh community of *Scirpus californicus* in an ox-bow lake in the Brazoria National Wildlife Refuge. Site No. 4-5, Oyster Creek Quad.; view is westward.



Figure 25. Fresh-marsh community dominated by *Eleocharis quadrangulata* (squarestem spikesedge). *Hymenocallis caroliniana* (Carolina spider lily) is the flowering plant. Site No. 15-6, Texas City Quad.

### **Forested Wetland Communities (Swamps)**

Forested wetlands as defined by Cowardin and others (1979) include swamps as well as forested areas less frequently inundated. Swamps, as defined most commonly, are woodlands or forested areas that contain saturated soils or are inundated by water during much of the year. This community is located almost entirely in the alluvial valley of the Trinity River. The swamp community is composed principally of *Taxodium distichum* (bald cypress) (fig. 26). Associated species may include *Cephalanthus occidentalis* (button bush), *Planera aquatica* (water-elm), and *Carya aquatica* (water hickory) (Harcombe and Neaville, 1977).

Areas along the floodplains of streams (excluding swamps) support assemblages of water-tolerant trees and shrubs (fig. 23) that are inundated less frequently than swamps. Trees and shrubs occurring in these areas include *Planera aquatica*, *Quercus phellos* (willow oak), *Quercus nigra* (water oak), *Fraxinus pennsylvanica* (Green ash), *Fraxinus caroliniana* (Carolina ash), *Salix nigra* (black willow), *Ulmus* spp. (elm), *Celtis laevigata* (sugar-berry), *Carya illinoensis* (pecan hickory), *Carya aquatica* (water hickory), *Cephalanthus occidentalis*, *Ilex vomitoria* (yaupon), *Liquidambar styraciflua* (sweet gum), *Sepium sebiferum* (Chinese tallow), *Parkinsonia aculeata* (retama), *Gleditsia aquatica* (water locust), and *Sabal minor* (dwarf palmetto). Occurring with hardwoods in some topographically higher areas is *Pinus taeda* (loblolly pine).

### **Submerged Vegetation Community**

Submerged vegetation has a limited distribution in the Galveston Bay system. It occurs principally in patches along the margins of the Trinity River delta, upper Trinity Bay, and Christmas Bay (figs. 27 and 28). Plant species occurring in the comparatively fresh area of the Trinity River delta include *Ruppia maritima* (widgeongrass), *Vallisneria americana* (wild celery), *Potamogeton pusillus* (pondweed), and *Najas quadrangularis* (water nymph) (Pulich and others, 1991). The dominant submerged vegetation along the north and eastern shores of upper Trinity Bay is *Ruppia maritima* (Pulich and White, 1991). In the Christmas Bay area, near Follets island, several true seagrasses occur including *Halodule wrightii* (shoalgrass), the dominant species, *Halophila engelmannii* (clovergrass), and *Thalassia testudinum* (turtlegrass) (Pulich and White, 1991). *Ruppia maritima* is abundant in many inland water bodies and tidal creeks (fig. 13).

The submerged-vegetation community is classified under sounds and bays by Shaw and Fredine (1956); as grassflats by Fisher and others (1972, 1973), and White and others (1985); and as submerged vegetation by Diener (1975). Submerged-vegetation communities are designated as Estuarine, subtidal, aquatic bed (E<sub>1</sub>AB) in the classification by Cowardin and others (1979); the water-regime modifier is "subtidal" (L) (table 6).

### **Soils and Wetland Community Relationships**

At the more than 135 sites surveyed around the Galveston Bay system, approximately 40 soil types were identified from county soil surveys (table 11). Several soils were encountered more frequently than others, and can be considered the dominant soils corresponding to wetland communities. For example, the soil most frequently occurring at wetland survey sites was the Harris clay. This typically saline, poorly drained soil is flooded by abnormally high tides, and




Figure 26. Swamp community dominated by *Taxodium distichum* along the Trinity River inland from Interstate Highway 10. Site No. 29-3, Cove Quad.; view is northward.




Figure 27. Generalized map showing the locations of submerged vegetation along the margins of the Trinity River delta. (From Pulich and others, 1991)




Figure 28. Generalized map showing the locations of submerged vegetation in 1956 and 1987 in the Galveston Bay system excluding areas along the Trinity Delta. The 1956 distribution of submerged vegetation in Trinity and Christmas Bays is not shown. (Modified from Pulich and White, 1991)

Table 11. Characteristics of soils at survey sites. (From USDA Soil Conservation Service County Soil Surveys—Crout, 1976; Wheeler and others, 1976; Crenwelge and others, 1981; and Crenwelge and others, 1988).

SOIL	SALINE OR NONSALINE	DRAINAGE	FREQUENCY OF FLOODING	WATER TABLE
Adine-urban complex	nonsaline	well drained	rarely flooded	
Aaa silty clay loam	nonsaline	moderately well drained	rarely flooded	lower soil saturated 2-4 months in wet season <1 ft below surface during winter
Atascos fine sandy loam, 1 to 4% slopes	nonsaline	poorly drained	rarely flooded	<2 ft below surface during winter
Bacilli clay	nonsaline	somewhat poorly drained	rarely flooded	saturated in and above lower soil in wet season
Bernard clay loam	nonsaline	somewhat poorly drained	rarely flooded	1 to 3 ft below surface in winter
Boy loamy fine sand	nonsaline	somewhat poorly drained	rarely flooded	
Brazoria clay, 0-1% slopes	nonsaline			
Capion muddy silty clay loam	saline	soil saturated		
Capion-Tracosa complex	saline	flooded daily by 2-12 inches of tide water		
Clemmle silty clay loam	nonsaline	flooded daily by 2-12 inches of tide water		
Follett clay loam	saline	rarely flooded		
Galveston-Nass complex	nonsaline-slightly saline	very poorly drained	flooded daily during high tide	at or near surface most of year
Galveston-Nass complex	saline	excessively to very poorly drained	occasionally to frequently flooded	36-50 inches below to 24 inches above surface
Harrit-Tracosa complex	typically saline	very poorly drained	<20 inches below surface throughout year	
Harris clay	saline	poorly drained	<20 inches below surface	
Iam clay, 0-2% slopes	saline	poorly drained	<1.5 ft below surface during winter	
Iam clay, 2-6% slopes	moderately saline	poorly drained	<1.5 ft below surface during winter	
Iam soils	nonsaline	very poorly drained	at surface to 30 inches below surface	
Kerman clay	saline	poorly drained	saturated within 30 inches of surface most of year	
Karenkawa muddy loam	nonsaline	occasionally flooded	depressions 0.3 ft deep, soil saturated	
Kaufman clay	nonsaline	flooded daily with 2 to 12 inches of tide water	surface to 50 inches below during wet season	
Kemah-urban land complex	nonsaline	frequently flooded	<1.5 ft below surface during winter	
Lake Charles clay, 1-3% slopes	nonsaline	rarely flooded		
Miocare-Telon complex	nonsaline	occasionally flooded to poorly drained	<3 ft below surface to 1 ft above	
Norey silt loam	nonsaline to slightly saline			
Mustang fine sand	nonsaline	poorly drained	frequent flooding	
Mustang fine sand, saline	slightly to strongly saline	poorly drained	frequency flooding by abnormally high tides	
Mustang-Nass complex	nonsaline to moderately saline	poorly to very poorly drained	occasionally to frequently flooded	
Nerta fine sandy loam	saline	somewhat poorly drained	occasionally flooded	
Nerta fine sandy loam	moderately saline	somewhat poorly drained	rarely flooded	
Nassa very fine sand loam	slightly to strongly saline	somewhat poorly drained	occasionally flooded by storm tides and rains	
Nass-Galveston complex, shell substratum	nonsaline to moderately saline	poorly to somewhat excessively drained	occasionally to frequently flooded	
Piedoco clay	saline	very poorly drained	frequently flooded	
Sierra loam, 0 to 3% slopes	moderately saline	somewhat poorly drained	rarely flooded	
Surpui clay	nonsaline	poorly drained	ponded for several months during year	
Surside clay	saline	poorly drained	rarely flooded	
Tatum clay loam	saline	very poorly drained	flooded daily during high tide	
Tracosa muddy clay	saline	poorly drained to ponded	flooded daily by 2 to 12 inches of tide water	
Tracosa muddy clay-day, low complex	saline	very poorly drained to ponded	frequently flooded	
Vanton clay	nonsaline	somewhat poorly drained	somewhat poorly drained	
Vanton clay, 1 to 4% slopes	saline	somewhat poorly drained	somewhat poorly drained	
Velasco clay	nonsaline	poorly drained	rarely flooded	
Verland silty clay loam	saline	soil saturated	frequently flooded	
Veston loam	slightly to strongly saline	poorly drained	flooded by unusually high tide	
Veston loam (Galveston Co.)	strongly saline	poorly drained		
Veston silty clay loam			<1 ft below surface	

Table 11 (cont.)

SOIL	DOMINANT PLANTS
Adina-urban complex Aaa silty clay loam	native pine and hardwoods; grasses include little bluestem, beaked panicum, longleaf uniota, brownseed paspalum hardwood trees; under-story—longleaf uniota (10%), kudz sedge (15%), Virginia wildrye (10%), switchgrass (5%), low panicum (5%), nimblewill muhly (5%); forbs such as elephantfoot and drummond waxmallow, vines and shrubs—greenbrier, poison-ivy, yaupon, possumhaw
Aleco fine sandy loam, 1 to 4% slopes Badlift clay	pine (dominant), hardwoods, sedges, beaked panicum, little bluestem (dominant), Indiangrass, switchgrass, eastern gamagrass, Florida paspalum, big bluestem, brownseed paspalum, panicum, sedges little bluestem (dominant), Indiangrass, switchgrass, eastern gamagrass, Florida paspalum, big bluestem, brownseed paspalum, panicum, sedges pine woodlands; bermudagrass, coastal bermuda grass, and bahiagrass
Bernard clay loam Bay loamy fine sand Brazoria clay, 0 to 1% slopes	hardwood trees; under-story—lindt sedge (35%), Virginia wildrye (10%), nimblewill muhly (10%), longleaf uniota (5%), nutgrass paspalum (5%); vines and shrubs—greenbrier, Alabama supple-leaf, yaupon, American elder, dwarf palmetto marshhay cordgrass, common reed, seashore saltgrass, sargentia, bulrushes, big cordgrass, smooth cordgrass Trescas—Smooth cordgrass (dominant), seashore saltgrass, glasswort, maritime salicorn, saltmarsh burrush, widgeongrass; (for Caplen see above)
Caplen muddy silty clay loam Caplen-Tracosa complex Chamblee silty clay loam	hardwood trees; under-story—lindt sedge (15%), Virginia wildrye (10%), longleaf uniota (10%), switchgrass (5%), low panicum (5%), nimblewill muhly; forbs (10%) such as elephantfoot and drummond waxmallow; vines and shrubs—greenbrier, poison-ivy, yaupon, possumhaw smooth cordgrass (40%)
Follett clay Loam Galveston-Nass complex	swales—marshhay cordgrass, seashore saltgrass, seashore paspalum, gulfine paspalum, shorgrass, gulf cordgrass, red lovegrass, needleglass rush, sea-oxeye, glasswort ridge—gulfine paspalum, marshay cordgrass, bushy bluestem, red lovegrass, knotroot bristlegrass, bushy bluestem, bermudagrass, baucharis, southern wax myrtle marshay cordgrass (50%), seashore saltgrass (10%), seashore paspalum (10%), shiny burrush (10%); 5% forbs—saltmarsh aster, sea-oxeye, becerra
Harris-Tracosa complex Harris clay Ham clay, 0-2% slopes Ham clay, 2-5% slopes Ham soils	marshay cordgrass (50%), seashore saltgrass (25%) gulf cordgrass (dominant), little bluestem, switchgrass, Indiangrass, marshay cordgrass, knotroot bristlegrass, longpike tridens gulf cordgrass (dominant), little bluestem, switchgrass, Indiangrass, marshay cordgrass, knotroot bristlegrass, longpike tridens gulf cordgrass (75%), marshay cordgrass (2%), common reedgrass (5%), little bluestem (2%), knotroot bristlegrass (2%), forbs (5%) common bermudagrass and dallisgrass; woodlands—elm, water oak, beach, willow oak, cypress, palmetto, sedges, longleaf uniota, and switch cane smooth cordgrass (20%)
Kermanawa muddy Loam Kuimanian clay Kuiman-Urban land complex Lake Charles clay Mocary-Laton complex	swamps—switchgrass, dallisgrass, tail fescue, johnsongrass, bluebeans, clovers, water-tolerant hardwoods—cypress, water oak, sweetgum not described little bluestem (50%), indiangrass (10%), eastern gamagrass (5%), big bluestem (5%), brownseed paspalum (5%), Florida paspalum (3%) switchgrass, maidencane, eastern gamagrass (dominants); Indiangrass, Florida paspalum, longlom, squarestem spikegrass, brownseed paspalum, knotroot bristlegrass, and low panicum; needlerush grass, rushes, sedge campiglossa, baucharis, sebania prairie grasses such as bermudagrass, bahiagrass, dallisgrass, bluegrass, bulrush, Indiangrass, beaked panicum, paspalum, sedges, and others;
Morey silty loam	woodland species may include lobolly pine, slash pine, white oak, red oak, and sweetgum gulfine paspalum (30%), marshay cordgrass (20%), herbaceous nimble, beach ground cherry, warmyrte, eastern baucharis marshay cordgrass (25%), sedges and rushes (25%), more saline areas—maritime seahort, shoregrass, glasswort, sea-oxeye, seashore saltgrass swales—marshay cordgrass, seashore paspalum, seashore saltgrass, gulf cordgrass, red lovegrass, needleglass, redlovegrass rush, sea-oxeye, glasswort ridge—gulfine paspalum, marshay cordgrass, bushy bluestem, red lovegrass, knotroot bristlegrass, bushy bluestem, bermudagrass, saltmarsh burrush,
Mustang fine sand Mustang fine sand, saline Mustang-Nass complex	gulf cordgrass (60%), marshay cordgrass (5%), switchgrass (5%), little bluestem (5%), seashore saltgrass (5%), forbs—sea-oxeye gulf cordgrass (dominant), little bluestem, switchgrass, Indiangrass, marshay cordgrass, knotroot bristlegrass, longpike tridens marshay cordgrass, seashore saltgrass, seashore paspalum swales—marshay cordgrass, seashore saltgrass, seashore paspalum, gulf cordgrass, bushy bluestem, red lovegrass, knotroot bristlegrass, bushy bluestem, saltmarsh burrush,
Narta fine sandy loam	marshay cordgrass (dominant), little bluestem, switchgrass, Indiangrass, marshay cordgrass, knotroot bristlegrass, longpike tridens giant cordgrass (20%), maidencane (25%), cattail (10%) gulf cordgrass (80%), sea-oxeye and other forbs
Nass very fine sand Nass-Galveston complex, shell substratum	smooth cordgrass (80%) smooth cordgrass (80%)
Placido clay	smooth cordgrass (dominant); seashore saltgrass, glasswort, maritime salicorn, saltmarsh burrush; widgeongrass
Seavers loam, 0 to 3% slopes Sumpf clay Tracosa clay Tracosa muddy clay-day, low complex Varnont clay	little bluestem (dominant), Indiangrass, switchgrass, eastern gamagrass, Florida paspalum, big bluestem, brownseed paspalum, panicum, sedges mixed pine and hardwoods, sedges, switchgrasses, and bluestem; bermudagrass, dallisgrass marshay cordgrass (60%), seashore saltgrass (15%), seashore paspalum (15%) little bluestem (dominant), Indiangrass, switchgrass, eastern gamagrass, Florida paspalum, big bluestem, brownseed paspalum, panicum, sedges gulf cordgrass (50%), marshay cordgrass (20%), Indiangrass, little bluestem, switchgrass, knotroot bristlegrass in higher areas maritime seahort, shoregrass, glasswort, and sea-oxeye interspersed with barren flat, seashore saltgrass, sea lavender, sepweed, Carolina wolfberry, and eastern baucharis shoregrass (35%), bushy sea-oxeye (10%)
Vanson loam (Galveston Co.) Vanson silty clay loam	

supports a vegetation assemblage composed predominantly of *Spartina patens* and *Distichlis spicata* (table 11). These species were the most frequently encountered during field surveys.

To simplify the discussion of soil types and their relationships to wetland communities, Marsh Rangeland Sites defined by Crenwelge and others (1988) in the soil survey of Galveston County will be used for comparing soils with wetland communities described in this report.

Marsh Rangeland Sites (Crenwelge and others, 1988) include the following sites, or complexes: (1) Salt Marsh Range Site, (2) Tidal Flat Range Site, (3) Salt Flat Range Site, (4) Low Coastal Range Site, (5) Coastal Swale Range Site, (6) Deep Marsh Range Site, (7) Salty Prairie Range Site, and (8) Coastal Sand Range Site.

The Salt Marsh Range Site, with elevations of 1 to 4 ft above sea level, occurs in relatively level coastal marsh areas and in flood plains. It is composed of the Harris clay (Ha and 19), Placedo clay (Pd), and Veston loam, strongly saline (Vx) (table 11). Almost 40 sites, or about 30 percent of all the sites surveyed, corresponded to the Salt Marsh Range Site complex as defined by Crenwelge and others (1988). Based on field survey locations, the wetland communities that were typically found on these soils are brackish-water and salt-water marshes (as mapped by White and others, 1985) (appendix A). These communities make up 70 percent of the survey sites within the Salt Marsh Range. High brackish-water marshes represented 30 percent of the sites. Among the dominant species in high brackish- and high salt-water marshes are *Spartina patens* and *Distichlis Spicata* (table 8).

The Tidal Flat Range Site corresponds to broad coastal tidal marshes at elevations slightly below sea level to about 1 ft above sea level. It consists of the Follet clay loam (Fo), Tatlum clay loam (Ta) and the Tracosa soil in the Caplen-Tracosa complex (Ct), the Tracosa mucky clay (Tm), and the Tracosa mucky clay-clay, low complex (Tx) (table 11). Approximately 15 percent of the field survey sites are located within the the Tidal Flat Range Site. The predominant wetland communities (as defined and mapped by White and others, 1985) are proximal salt-water marshes, which represent about 70 percent of the field survey sites located in the Tidal Flat Range Site. The predominant vegetation is *Spartina alterniflora*; other species may include *Batis maritima*, *Distichlis spicata*, *Salicornia* spp., *Scirpus maritimus*, and *Juncus roemerianus*.

The Salt Flat Range Site occurs in nearly level coastal marshes with elevations slightly above mean sea level to about 3 ft above sea level. Soils of this range site are strongly saline Mustang fine sand (Ms) and very strongly saline Veston loam (Vx) (table 11). Sixteen survey sites were located within these soils, or slightly more than 10 percent of all sites surveyed. Wetland communities represented on the Salt Flat Range Site are predominantly salt-water marshes, but some include transitional areas and mixtures of marshes and barren sand flats (White and others, 1985) (appendix A). Vegetation includes *Batis maritima*, *Monanthochloe littoralis*, *Salicornia* spp., *Borrichia frutescens*, *Distichlis spicata*, *Limonium nashii*, *Lycium carolinianum*, and others.

The Coastal Swale Range Site occurs in swales between beach ridges and in shallow depressions on nearly level coastal flats. Soils in this range site are principally in the Nass soil of the Galveston-Nass complex (Gc), the Mustang-Nass complex (Mt), and the Nass-Galveston complex shell substratum (Nx). Vegetation communities were surveyed at nine sites corresponding to soils in the Coastal Swale Range Site. The areas surveyed were mostly located on Galveston Island, much of which is characterized by relict beach ridge and swale topography. Vegetation communities are predominantly defined by brackish- and salt-water marshes, both low and high marshes (White and others, 1985). Vegetation includes *Spartina patens*, *Distichlis spicata*, *Paspalum vaginatum*, *Paspalum monostachyum*, *Monanthochloe littoralis*, *Spartina spartinae*, *Juncus roemerianus*, *Salicornia* spp., and *Borrichia frutescens*.

The Deep Marsh Range Site commonly corresponds with marshes near bays and bayous where tidal-water salinities are lower because of saltwater and freshwater mixing. Elevations range from sea level to 1 ft above. Soils include the Caplen mucky silty clay loam (Ca), and the Caplen soil in the Caplen-Tracosa complex. Dominant vegetation is *Spartina patens* and *Distichlis spicata*. *Spartina cynosuroides* has been a dominant species on this range site in the past, but has been replaced principally by *Spartina patens* (Crenwelge and others, 1988). Depending on water depth and salinities, *Sagittaria* and bulrushes may also occur in this marsh range site. Only a couple of survey sites (high, or distal, salt-water marshes) occur within this range site.

The Salty Prairie Range Site occurs on broad, relatively level coastal flats and marshes, where elevations range from 2 to 8 ft above sea level. Among the soils characterizing this range site is the Ijam soil in the Ijam clay, 0-2 percent slopes (ImA), and 2-8 percent slopes (ImB), Narta fine sandy loam (Na), Sievers loam (SeB), and slightly saline Veston loam (Vx). Most of the survey sites in this range site correspond to the Ijam soils, which might be considered a disturbed soil complex (fig. 29). Ijam soils are formed in saline, clayey, marine and alluvial sediment deposits that were dredged to construct and maintain canals or waterways. Plant communities on these soils vary widely because of the variations in salinities and elevations that characterize this range site. Plant communities may include brackish and salt marshes, barren flats, transitional areas, and uplands. The dominant vegetation in many topographically higher areas is *Spartina spartinae*. Other species may include *Borrichia frutescens*, *Panicum virgatum*, *Spartina patens*, *Phragmites australis*, and *Setaria geniculata*.

The Low Coastal Range Site consists of level to gently sloping coastal sands that roughly parallel the Gulf shoreline; elevations are less than 3.3 m (10 ft) above sea level. Soils in this range site are the Galveston soil in the Galveston-Nass complex (Gc) and Nass-Galveston complex (Nx), and Mustang soils in Mustang fine sand (Mn), Mustang-Nass complex (Mt), and Mustang fine sand, slightly saline (Ms). The Galveston and Mustang soils are at elevations generally too high for marsh development, and therefore, correspond most frequently to uplands (U) and possibly transitional areas as mapped by White and others (1985). Wetlands occur in the Nass soils of the Gc and Nx complexes (see Coastal Swale Range Site).

The Coastal Sand Range Site is composed of nearly level to undulating coastal ridges that parallel the Gulf shoreline. Elevations, which are up to 4 m (12 ft) above mean sea level, preclude marsh development on this range site.

#### **Examples of Wetland Profiles Developed From Topographic Survey Transects**

Topographic surveys of marsh communities were conducted at selected sites around the Galveston Bay system. These data are presented in appendix C. Descriptions of the zonation of plant species along two transects are presented here.

#### **Smith Point Transect**

The elevation survey of the Smith Point marsh is shown in figure 11. The transect has a bearing of south 45 degrees west (S45°W) and is approximately 85 m (279 ft) long. The southwest end of the transect intersects the shoreline of East Galveston Bay. The total range in elevation of the transect is approximately 1.5 m (5 ft), which is the vertical distance from station 1 (just below the water line) to station 6, the crest of the shell berm. Marsh plants, which are absent on the shell berm, have a much lower range in elevation, about 45 cm (1.5 ft) (fig. 11). This salt marsh community, which is classified as an estuarine intertidal emergent community (E<sub>2</sub>EM) as



Figure 29. Disturbed-area community on a small spoil mound along the Intracoastal Waterway on the landward margin of Follets Island. A mixed assemblage of approximately 10 salt-marsh species occurs on the mound. Species range from *Iva frutescens* and *Spartina spartinae* at the top, to *Batis* and *Salicornia* at the base. Site No. 1-2, Freeport Quad. (This Quad is not officially part of the project area).

defined by Cowardin and others (1979), is made up of about 8 different species. *Spartina alterniflora* (smooth cordgrass), as expected, occurs at the lowest elevation (water line), and a community composed of *Spartina spartinae* (gulf cordgrass or sacahuista), *Spartina patens* (marshhay cordgrass), *Iva frutescens* (bigleaf sumpweed or marshelder), and *Borrichia frutescens* (sea-oxeye) occurs at the highest elevation (stations 18 and 19, fig. 11). The profile exemplifies how small changes in elevation along the microtidal Texas coast can affect plant distribution. Occurring at elevations between the water line and the highest marsh plants on the profile are several species (fig. 11) including, at lower elevations, *Scirpus maritimus* (salt-marsh bulrush) and *Juncus roemerianus* (needle rush); at slightly higher elevations *Distichlis spicata* (seashore saltgrass) occurs. *Spartina patens* and *Borrichia* also occur at intermediate elevations, but are still higher than *Spartina alterniflora*, *Scirpus*, *Juncus*, and *Distichlis*. The range in elevation for *Spartina alterniflora* is about 25 cm (0.8 ft) along this transect, so it occurs mixed with other species locally.

A close look at the profile (fig. 11) shows that very small changes in elevation can apparently increase the regularity of flooding and enable species like *Spartina alterniflora* to become established. Stations 10 and 14 have *Spartina alterniflora* mixed with *Distichlis*. At slightly higher elevations toward station 12, only *Distichlis* is present.

This particular survey shows that, in general, the species occurring at lowest (and therefore most frequently flooded) elevations are *Spartina alterniflora*, *Scirpus maritimus*, and *Juncus roemerianus*, with *Distichlis* mixing with these species locally. Occurring at higher elevations are *Spartina patens*, *Borrichia*, *Spartina spartinae*, and *Iva frutescens*.

Wetland indicator plant species designations in the *National List of Plant Species that Occur in Wetlands: 1988 Texas*, by P. B. Reed, USFWS, were used as a guide to help delineate species associations in some areas. Species identified along the Smith Point profile are all wetland species, but *Spartina alterniflora*, *Scirpus maritimus*, and *Juncus roemerianus* are classified as obligate (OBL) wetland plants, which means that under natural conditions they have an estimated probability of occurring in wetlands greater than 99 percent of the time. The other species listed above (i.e., those occurring at slightly higher and drier elevations) are facultative wetland (FACW) plants, which means that they usually occur in wetlands (estimated probability of 67 to 99 percent), but occasionally are found in nonwetlands. As expected, the elevation measurements properly defined the species that can tolerate wetter conditions and are therefore more frequently found in wetlands.

### Brazoria National Wildlife Refuge Transect

The second salt marsh transect along which elevations, distances, and bearings were measured was located in the Brazoria National Wildlife Refuge (fig. 12). The transect, which is approximately 375 m (1,230 ft) long, is oriented roughly perpendicular to the hydrologic gradient and was tied to a USGS bench mark with an elevation of 2.2 m (6.6 ft) at the northwest end of the transect. Lower elevations occur on the downthrown side of a fault located at stations 4 and 5 on the profile (fig. 12). The difference in elevations on each side of the fault line produces a dramatic effect in the vegetation communities. Between the bench mark and station 4 at the edge of the fault (this segment of the transect marks the upthrown side of the fault) the plant community is dominated by *Spartina spartinae*, with scattered species including *Setaria geniculata* (knotroot bristlegrass), *Iva annua* (seacoast sumpweed), and *Aster* spp. Additional species reported in this area in the Brazoria County Soil Survey include *Nothoscordum bivalve* (false garlic) and *Sabatia campestris* (prairie rose-gentian). The dominant species *Spartina spartinae* is classified as a facultative wetland (FACW), but other species, except for *Aster* (OBL), are found much less frequently in wetlands. *Iva annua* and *Setaria* are classified as facultative

(FAC), and are, therefore, equally likely to occur in nonwetlands as wetlands. *Sabatia* and *Nothoscordum* are classified as facultative upland species (FACU), which means the probability of their occurring in a wetland is only 1 to 33 percent.

On the downthrown side of the fault, a definite wetland community occurs. The drop in elevation from the top of the fault scarp to the wetland community is more than 30 cm (1 ft). Plant species between stations 5 and 6 (fig. 12) on the profile are composed of patches of *Monanthochloe*, *Salicornia*, and *Batis* occurring within a sand/mud flat that is capped by an algal mat. At lower elevations, between stations 6 and 7, *Distichlis* composes about 90 percent of the community, with scattered *Salicornia* making up the remaining 10 percent. All the species on the downthrown side of the fault, where wetter conditions characterize the lower elevations, are obligate wetlands.

#### ACKNOWLEDGMENTS

This study was sponsored by the Galveston Bay National Estuary Program, funded by the United States Environmental Protection Agency under assistance agreement with the Texas Water Commission. Special thanks go to Diane Spinney and Thomas Tremblay of the Bureau of Economic Geology for assistance in the geographic information system, ARC-INFO, and to Margaret Evans and Jamie Coggin for assistance in figure and table paste-up. Editing was by Kitty Challstrom and final word processing was by Susan Lloyd, under the direction of Susann Doenges, Editor in chief.

## REFERENCES

- Adams, D. B., and Tingley, C. A., 1977, A study of the hydrological and nutrient exchange processes in the Trinity River delta, Texas, part II, marsh habitats and primary production in the Trinity River delta: report prepared for Texas Department of Water Resources by Espey, Huston and Associates, Inc., document no.77-103, job no. 0216, 20 pp.
- Anderson, J. R., Hardy, E. E., Roach, J. T., and Witmer, R. E., 1976, A land use and land cover classification system for use with remote sensor data: U.S. Geological Survey Professional Paper 964, 27 pp.
- Benton, A. R., Jr., Snell, W. W., and Clark, C. A., 1979, Monitoring and mapping of Texas coastal wetlands, Galveston Bay and Sabine Lake areas, 1978 growing season: Texas A&M University, College Station, Remote Sensing Center, technical report RSC-102, 123 pp.
- Bray, W. L., 1906, Vegetation of Texas: University of Texas Bulletin, v. 82, no. 10, p. 1-106.
- Brown, L. E., 1985, An annotated checklist of the vascular plants of the Armand Bayou Nature Park and vicinity, southeastern Harris County, Texas.
- Chabreck, R. H., 1972, Vegetation, water, and soil characteristics of the Louisiana coastal region: Louisiana State University and Agricultural and Mechanical College Bulletin 664, 72 pp.
- Correll, D. S., and Correll, H. B., 1975, Aquatic and wetland plants of southwestern United States: Stanford University Press, Stanford, California, 2 volumes, 1777 pp.
- Cowardin, L. M., Carter, V., Golet, F. C., and LaRoe, E. T., 1979, Classification of wetlands and deep-water habitats of the United States: U.S. Department of the Interior, Fish and Wildlife Service, Office of Biological Services, Washington, D.C., FWS/OBS-79/31, 103 pp.
- Crenwelge, G. W., Crout, J. D., Griffin, E. L., Golden, M. L., and Baker, J. K., 1981, Soil survey of Brazoria County, Texas: U.S. Department of Agriculture, Soil Conservation Service, 182 pp., 36 maps.
- Crenwelge, G. W., Griffin, E. L., and Baker, J. K., 1988, Soil survey of Galveston County, Texas: U.S. Department of Agriculture, Soil Conservation Service, 182 pp., 36 maps.
- Crout, J. D., 1976, Soil survey of Chambers County, Texas: U.S. Department of Agriculture, Soil Conservation Service, 53 pp., 54 maps.

Diener, R. A., 1975, Cooperative Gulf of Mexico estuarine inventory and study—Texas; area description: National Oceanic and Atmospheric Administration, technical report, National Marine Fisheries Service Circular 393, 129 pp.

Fisher, W. L., Brown, L. F., Jr., McGowen, J. H., and Groat, C. G., 1973, Environmental Geologic Atlas of the Texas Coastal Zone—Beaumont–Port Arthur Area: The University of Texas at Austin, Bureau of Economic Geology, 9 maps, 93 pp.

Fisher, W. L., McGowen, J. H., Brown, L. F., Jr., and Groat, C. G., 1972, Environmental Geologic Atlas of the Texas Coastal Zone—Galveston–Houston Area: The University of Texas at Austin, Bureau of Economic Geology, 9 maps, 91 pp.

Fleetwood, R. J., no date, Plants of Brazoria/San Bernard National Wildlife Refuges, Brazoria County, Texas: U.S. Department of the Interior, Fish and Wildlife Service, 61 pp.

Frye, R. G., Brown, K. L., and McMahan, C. A., 1984, The vegetation types of Texas: Texas Parks and Wildlife Department, color map, scale 1:500,000.

Gabrysich, R. K., and Bonnet, C. W., 1975, Land-surface subsidence in the Houston–Galveston region, Texas, Texas Water Development Board, Report 188, 19 pp.

Godfrey, C., McKee, G. S., and Oakes, H., 1973, General soils map of Texas: Texas Agricultural Experiment Station Miscellaneous Publication, MP-1304, color map, scale 1:1,500,000.

Gosselink, J. G., Cordes, C. L., and Parsons, J. W., 1979, An ecological characterization study of the Chenier Plain coastal ecosystem of Louisiana and Texas: U.S. Fish and Wildlife Service, Office of Biological Services, FWS/OBS-78/9 through 78/11, 3 volumes.

Gould, F. W., 1975, Texas plants—a checklist and ecological summary: Texas Agricultural Experiment Station, MP-585/Revised, College Station, Texas, 121 pp.

Harcombe, P. A., and Neaville, J. E., 1977, Vegetation types of Chambers County, Texas: Texas Journal of Science, v. 29, p. 209–234.

Kuchler, A. W., 1966, Potential natural vegetation: U.S. Geological Survey, Washington, D.C., color map, scale 1:7,500,000.

Lazarine, P., no date, Common wetland plants of southeast Texas: U.S. Army Corps of Engineers, Galveston District, various pagination.

McAtee, J. W., 1976, Autecological aspects of Gulf cordgrass [*Spartina spartinae* (Trin.) Hitchc.] communities of the Texas coastal prairie, p. 2–12, in Range ecological and management

research on the coastal prairie, progress report of cooperative studies: Welder Wildlife Foundation and Texas A&M University, Texas Agricultural Experiment Station, Sinton, Texas.

Penfound, W. T., and Hathaway, E. S., 1938, Plant communities in the marshlands of southeastern Louisiana: Ecological Monographs, v. 8, no. 1, 56 pp.

Pulich, W. M., Jr., and White, W. A., 1991, Decline of submerged vegetation in the Galveston Bay System: chronology and relationship to physical processes: Journal of Coastal Research, v. 7, no. 4, p. 1125-1138.

Pulich, W. M., Jr., White, W. A., Castiglione, M., and Zimmerman, R. J., 1991, Status of submerged vegetation in the Galveston Bay System, in Shipley, F. S., and Kiesling, R. W., eds., proceedings, Galveston Bay characterization workshop, February 21-23, Galveston Bay National Estuary Program Publication GBNEP-6, p. 127-132.

Reed, P. B., Jr., 1988, National list of plant species that occur in wetlands: 1988—Texas: U.S. Fish and Wildlife Service, St. Petersburg, Florida, various pagination.

Shaw, S. P., and Fredine, C. G., 1956, Wetlands of the United States, their extent and their value to waterfowl and other wildlife: U.S. Fish and Wildlife Service, Circular 39, 47 pp.

Shew, D. M., Baumann, R. H., Fritts, T. H., and Dunn, L. S., 1981, Texas barrier island region ecological characterization: environmental synthesis papers: U.S. Department of the Interior, Fish and Wildlife Service, Office of Biological Services, Washington, D.C., FWS/OBS-81/32, 413 pp.

Tharp, B. C., 1926, Structure of Texas vegetation east of the 98th meridian: University of Texas, Austin, Bulletin 2606, 97 pp.

Thayer, G. W., and Ustach, J. F., 1981, Gulf of Mexico wetlands: value, state of knowledge and research needs, in Proceedings of a symposium on environmental research needs in the Gulf of Mexico, Atwood, D. K., convener, v. 2B, Key Biscayne, Florida, September 30-October 5, 1979 p. 1-30.

Thomas, G. W., 1975, Texas plants—an ecological summary, in Texas plants—a checklist and ecological summary, Gould, F. W., ed., Texas Agricultural Experiment Station, MP-585/Revised, College Station, Texas, p. 7-14.

Ward, G. H., Jr., and Armstrong, N. E., 1980, Matagorda Bay, Texas: its hydrography, ecology and fishery resources: U.S. Fish and Wildlife Service, FWS/OBS-81/52, 230 pp.

Webb, J. W., 1983, Soil water salinity variations and their effects on *Spartina alterniflora*: Contributions in Marine Science, v. 26, p. 1-13.

Wheeler, F. F., and others, 1976, Soil survey of Harris County, Texas: U.S. Department of Agriculture, Soil Conservation Service, 140 pp., 140 maps.

White, W. A., and Calnan, T. R., 1990, Sedimentation and historical changes in fluvial-deltaic wetlands along the Texas Gulf Coast with emphasis on the Colorado and Trinity River deltas: The University of Texas at Austin, Bureau of Economic Geology, final report prepared for the Texas Parks and Wildlife Department, 124 p., 7 Appendices.

White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., Littleton, T. G., McGowen, J. H., Nance, H. S., and Schmedes, K. E., 1985, Submerged lands of Texas, Galveston-Houston area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands: The University of Texas at Austin, Bureau of Economic Geology, Special Publication, 145 pp.

White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., Littleton, T. G., McGowen, J. H., and Nance, H. S., 1987, Submerged lands of Texas, Beaumont-Port Arthur area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands: The University of Texas at Austin, Bureau Economic Geology, Special Publication, 110 pp.

## **Appendix A**

**APPENDIX A. Field Site Surveys.**

Site Survey numbers, locations including UTM, soils and general wetland units.

Quadrangle	Site Number			Photograph	Site	UTM Coordinates		Soil*	Sub Land **
	Quad	Site	County			Easting	Northing		
Freeport	1 — 1	Brazoria	GHGH 8011	5	274150	3209100	10	TA/FH	
Freeport	1 — 2	Brazoria	GHKL 8098	1	278955	3207845	44	SD	
Freeport	1 — 3	Brazoria	GHKL 8098	2	280480	3207620	30,31	SP	
Christmas Point	3 — 1	Brazoria	GHGH 8012	3	281040	3216595	20,39	U	
Christmas Point	3 — 2	Brazoria	GHCD 7931	3	281300	3221205	32	BH/BL	
Christmas Point	3 — 3	Brazoria	GHGH 8012	1a	281620	3208600	30,31,16	SP/U	
Christmas Point	3 — 4	Brazoria	GHGH 8012	2b	283000	3215800	19	BH	
Christmas Point	3 — 5	Brazoria	GHCD 7932	3b	283350	3221240	19	BH/BL	
Christmas Point	3 — 6	Brazoria	GHCD 7932	3a	283905	3220980	19	BH/BL	
Christmas Point	3 — 7	Brazoria	GHGH 8013	2	283995	3211210	31,16	SP	
Christmas Point	3 — 8	Brazoria	GHCD 7932	3	284210	3220750	19	BH	
Christmas Point	3 — 9	Brazoria	GHCD 7932	2	284775	3220460	19	WU	
Christmas Point	3 — 10	Brazoria	GHCD 7932	2a	284860	3220745	19	BL	
Christmas Point	3 — 11	Brazoria	GHGH 8013	1a	286485	3213510	16	SP	
Christmas Point	3 — 12	Brazoria	GHGH 8014	1	292720	3217345	30,31	BL/SD/SP	
Oyster Creek	4 — 1	Brazoria	GHGH 8011	3	273610	3212160	36	U	
Oyster Creek	4 — 2	Brazoria	GHGH 8011	1	272260	3213300	water	FL	
Oyster Creek	4 — 3	Brazoria	GHGH 8011	2	273860	3212000	38	FH	
Oyster Creek	4 — 4	Brazoria	GHGH 8011	4	274070	3210700	12	FH/U	
Oyster Creek	4 — 5	Brazoria	GHGH 8012	5	279750	3216610	water	FL	
Oyster Creek	4 — 6	Brazoria	GHGH 8012	4	279850	3216575	39	TA	
Oyster Creek	4 — 7	Brazoria	GHGH 8012	3a	280035	3216575	39	BL	
Lake Como	5 — 1	Galveston	GHEF 7947	2	312785	3234040	Mt/Mn	BH/W	
Lake Como	5 — 2	Galveston	GHEF 7946	2a	313770	3234075	Gc	W	
Lake Como	5 — 3	Galveston	GHEF 7945	9	316380	3236780	Gc	BL	
Lake Como	5 — 4	Galveston	GHEF 7945	10	316480	3236400	Gc	BL	
Lake Como	5 — 5	Galveston	GHEF 7947	4b	309310	3231800	Ka	Sp	
Lake Como	5 — 6	Galveston	GHEF 7948	7	306510	3231330	Ka	Sp	
Sea Isle	6 — 1	Galveston	GHEF 7948	3a	304440	3236280	ImA	upland	
Sea Isle	6 — 2	Galveston	GHCD 7934	1	301170	3225900	Ka	Sp	
Hoskins Mound	7 — 1	Brazoria	GHCD 7932	1	283870	3226410	32	BH/TA	
Hoskins Mound	7 — 2	Brazoria	GHEF 7951	1	285660	3233485	16	BH	
Galveston	9 — 1	Galveston	GHEF 7945	1	318655	3237310	Mn	SP/BH	
Galveston	9 — 2	Galveston	GHAB 7868	2	323540	3245640	SeB	MU/SD	
Galveston	9 — 3	Galveston	GHAB 7868	1	326000	3245410	ImA,ImB	WU	
Virginia Point	10 — 1	Galveston	GHEF 7948	2c	306000	3238420	Tm/Pd	SP	
Virginia Point	10 — 2	Galveston	GHEF 7948	2b	306065	3238350	Tm/Pd	SP	
Virginia Point	10 — 3	Galveston	GHEF 7948	2a	306125	3238280	Tm/Pd	SP	
Virginia Point	10 — 4	Galveston	GHEF 7948	1c	307880	3240450	Vx	SP	
Virginia Point	10 — 5	Galveston	GHEF 7948	1b	307925	3240390	Vx	SP	
Virginia Point	10 — 6	Galveston	GHEF 7948	1a	307950	3240340	Vx	upland	
Virginia Point	10 — 7	Galveston	GHAB 7870	2a	311740	3244810	Tx	SO	
Virginia Point	10 — 8	Galveston	GHAB 7870	2b	312500	3245410	Tx	SO	
Virginia Point	10 — 9	Galveston	GHEF 7945	6	314535	3237970	Ka	SP/SO	
Virginia Point	10 — 10	Galveston	GHAB 7870	3	314725	3243510	Tx	SP	
Virginia Point	10 — 11	Galveston	GHAB 7870	1	313310	3248440	Na	SD	
Virginia Point	10 — 12	Galveston	GHEF 7945	7	315660	3238040	Mt	BH/MU	
Virginia Point	10 — 13	Galveston	GHEF 7945	8	315980	3237480	Nx	BL/BH	
Virginia Point	10 — 14	Galveston	GHEF 7945	3	317130	3239220	Mt	SP	
Virginia Point	10 — 15	Galveston	GHEF 7945	4	317490	3238810	Mu/Gc	BL/BH	
Virginia Point	10 — 16	Galveston	GHEF 7945	5	317675	3238510	Mu	BL/BH	
Virginia Point	10 — 17	Galveston	GHEF 7945	2	317780	3239330	Mn	upland	
Hitchcock	11 — 1	Galveston	GHAB 7873	1	296540	3244420	Be	W	
Hitchcock	11 — 2	Galveston	GHAB 7872	3,3a	302820	3249020	Ba	WL	

APPENDIX A (cont.)

Quadrangle	Site Number			Photograph	Site	UTM Coordinates		Soil*	Sub Land **
	Quad	Site	County			Eastng	Northing		
Caplen	1 2 — 1	Galveston	GHGH 7513	1a	347305	3262445	Vx	SP/SD	
Flake	1 3 — 1	Galveston	GHIJ 7568	2a	332245	3251545	Ms	SD/U	
Flake	1 3 — 2	Galveston	GHIJ 7568	1	332250	3251870	Mt	SD/SP	
Flake	1 3 — 3	Galveston	GHIJ 7568	2	332435	3251680	Mn	U/SD	
Flake	1 3 — 4	Galveston	GHIJ 7569	1a	334900	3257000	Vx, Ka	SP	
Flake	1 3 — 5	Galveston	GHIJ 7569	1	336520	3256410	Mt	SP/SD/LF	
Flake	1 3 — 6	Galveston	GHIJ 7569	2	337625	3258655	Vx	MU/SP	
Texas City	1 5 — 1	Galveston	GHIJ 7565	4	309340	3261240	Be	MU/BL	
Texas City	1 5 — 2	Galveston	GHIJ 7565	1	309800	3256070	Fo	SP	
Texas City	1 5 — 3	Galveston	GHIJ 7565	3	312050	3255500	Fo	SP	
Texas City	1 5 — 4	Galveston	GHIJ 7565	5	312740	3263430	Fo	BL	
	1 5 — 5	Site No. not used							
Texas City	1 5 — 6	Galveston	GHIJ 7565	6	314350	3263740	Md	FL	
Texas City	1 5 — 7	Galveston	GHIJ 7565	2a	314400	3255060	Vx	TA	
Texas City	1 5 — 8	Galveston	GHIJ 7566	1	316260	3256060	Fo	SD/SP	
Texas City	1 5 — 9	Galveston	GHIJ 7566	2	316480	3257140	Vx	SD	
Dickinson	1 6 — 1	Galveston	GHIJ 7563	1	299050	3259710	Va	WL	
Dickinson	1 6 — 2	Galveston	GHIJ 7563	1a	299485	3260040	LaB	WL	
Dickinson	1 6 — 3	Galveston	GHIJ 7564	2	304340	3261070	Ba	BL	
High Island	1 7 — 1	Galveston	BPA-GH 7510	1b	363320	3269905	Pd	BL	
High Island	1 7 — 2	Galveston	BPA-GH 7511	1	363340	3271955	Pd	BL/BH	
High Island	1 7 — 3	Galveston	BPA-GH 7510	4	365095	3273830	Pd	BH	
Frozen Point	1 8 — 1	Galveston	GHGH 7513	4c	346785	3266695	Ca	SD	
Frozen Point	1 8 — 2	Galveston	GHGH 7513	4b	346900	3266900	Ct/Ca	SD	
Frozen Point	1 8 — 3	Galveston	GHGH 7513	4a	346900	3267040	Ct	SD	
Frozen Point	1 8 — 4	Galveston	GHGH 7513	3a	347780	3267550	Ct	SD	
Frozen Point	1 8 — 5	Galveston	GHGH 7513	3b	347830	3267500	Ct	SP	
Frozen Point	1 8 — 6	Galveston	GHGH 7513	2b	349340	3267320	Ct	SP	
Frozen Point	1 8 — 7	Galveston	GHGH 7513	2a	349360	3267420	Ct	SD	
Frozen Point	1 8 — 8	Chambers	GHGH 7513	1	351620	3273900	Ha	BH/BL	
Frozen Point	1 8 — 9	Chambers	BPA-GH 7512	2	352140	3273220	Ha	SD	
Frozen Point	1 8 — 10	Chambers	GHGH 7512	5b	352400	3268760	Ve	TA	
Frozen Point	1 8 — 11	Chambers	GHGH 7512	5a	352690	3268800	Ve	TA	
Frozen Point	1 8 — 12	Galveston	GHGH 7512	1	353765	3265090	Vx	MU	
Lake Stephenson	1 9 — 1	Chambers	GHCD 7464	1	337640	3275970	Ha	BL	
Smith Point	2 0 — 1	Chambers	GHGH 7516	1	326700	3269590	Ve	SD/SP	
Bacliff	2 1 — 1	Chambers	GHCD 7469	2a	310890	3278320	Im	SP	
Bacliff	2 1 — 2	Chambers	GHCD 7469	2b	311050	3278360	Im	SP	
Bacliff	2 1 — 3	Chambers	GHCD 7469	3	311960	3277930	water	LF	
League City	2 2 — 1	Galveston	GHGH 7522	1	297100	3268610	Vx	TA	
League City	2 2 — 2	Harris	GHCD 7471	1	298020	3275400	Na	FL	
League City	2 2 — 3	Harris	GHCD 7471	2	299380	3274600	AtB	BL/BH	
League City	2 2 — 4	Harris	GHCD 7470	1	302510	3276460	VaB	WL	
League City	2 2 — 5	Harris	GHGH 7521	1	303160	3271750	AtB	BL	
Morgans Point	2 6 — 1	Harris	GHCD 7469	1	307935	3284355	Na	TA/BH	
Morgan's Point	2 6 — 2	Chambers	GHCD 7469	7	308750	3285730	Im	LF/U	
Morgan's Point	2 6 — 3	Chambers	GHCD 7469	6	309280	3284850	Im	BL	
Morgan's Point	2 6 — 4	Chambers	GHCD 7469	5	309710	3283040	Im	BL	
Morgan's Point	2 6 — 5	Chambers	GHCD 7469	4b	311070	3280480	Im	W	
Morgan's Point	2 6 — 6	Chambers	GHCD 7469	4a	311150	3280380	water	upland	
Morgan's Point	2 6 — 7	Chambers	GHCD 7468	1	314130	3282425	Im	SP	
Morgan's Point	2 6 — 8	Chambers	GHCD 7468	2	314560	3282320	LaB	WL/SD	
Morgan's Point	2 6 — 9	Chambers	GHCD 7468	3	314910	3285395	Ha	BL/LWL	
La Porte	2 7 — 1	Harris	GHEF 7496	1	299420	3292140	LcB	BH	
Anahuac	2 8 — 1	Chambers	GHAB 7451	4a	332330	3295850	Ha	LF/BL	
Anahuac	2 8 — 2	Chambers	GHAB 7451	9e	333420	3297190	Ha	BH	
Anahuac	2 8 — 3	Chambers	GHAB 7451	9d	333460	3297270	Ha	BH	
Anahuac	2 8 — 4	Chambers	GHAB 7451	9c	333490	3297320	Ha	BH	
Anahuac	2 8 — 5	Chambers	GHAB 7451	9b	333520	3297360	Ha	BH	
Anahuac	2 8 — 6	Chambers	GHAB 7451	9a	333560	3297420	Ha	U	
Anahuac	2 8 — 7	Chambers	GHAB 7452	4a	334570	3298880	Ka	BH	
Anahuac	2 8 — 8	Chambers	GHEF 7501	4a	335100	3293100	Ha	LF	
Anahuac	2 8 — 9	Chambers	GHEF 7501	4b	335150	3293170	Ha	BL	
Anahuac	2 8 — 10	Chambers	GHEF 7501	3b	336405	3293860	Ha	BL	

**APPENDIX A (cont.)**

Quadrangle	Site Number			Photograph	Site	UTM Coordinates		Soil*	Sub Land **
	Quad	Site	County			Easting	Northing		
Anahuac	2 8 — 1 1	Chambers	GHEF 7501	3a	336465	3293860	Ha	BH	
Anahuac	2 8 — 1 2	Chambers	GHAB 7452	1a	337000	3300540	Ha	FL	
Cove	2 9 — 1	Chambers	GHAB 7451	6	327000	3297800	Ha	BH	
Cove	2 9 — 2	Chambers	GHAB 7451	5	327220	3298140	Ha	BH	
Cove	2 9 — 3	Chambers	GHAB 7451	7	328370	3301790	Ka/Ha	FH/W	
Cove	2 9 — 4	Chambers	GHAB 7451	11a	328655	3298960	reservoir	BH	
Cove	2 9 — 5	Chambers	GHAB 7451	11b	328680	3299080	reservoir	BL	
Cove	2 9 — 6	Chambers	GHAB 7451	10a	330190	3301000	reservoir	U	
Cove	2 9 — 7	Chambers	GHAB 7451	10b	330340	3300920	reservoir	BL/BH	
Cove	2 9 — 9	Chambers	GHAB 7451	10d	330720	3300550	reservoir	BL	
Cove	2 9 — 1 0	Chambers	GHAB 7451	8	330120	3302400	Mo	SW	
Highlands	3 1 — 1	Harris	GHAB 7446	8	296500	3296740	An	WL	
Highlands	3 1 — 2	Harris	GHAB 7446	3	298460	3302465	Ka	WL	
Highlands	3 1 — 3	Harris	GHEF 7495	1	298720	3293845	Is	BH	
Highlands	3 1 — 4	Harris	GHAB 7446	6	298960	3297900	Ku	BL	
Highlands	3 1 — 5	Harris	GHAB 7446	1d	300000	3302300	Ka	WS	
Highlands	3 1 — 6	Harris	GHAB 7446	1c	300100	3302300	Ka	WL	
Highlands	3 1 — 7	Harris	GHAB 7446	1b	300180	3302400	Ka	FH	
Highlands	3 1 — 8	Harris	GHEF 7496	2	300580	3295820	VaB	WL	
Highlands	3 1 — 9	Harris	GHAB 7446	1a	300265	3302280	Ka	FL/WL	
Highlands	3 1 — 1 0	Harris	GHAB 7446	5	300000	3297805	Bo	BL	

\*Soil abbreviations are shown in table 2 A.

\*\* Sub Land column shows wetland units as mapped by White and others (1985). Abbreviations are defined below.

- BH Brackish-water marsh, high
- BL Brackish-water marsh, low
- FH Fresh-water marsh, high
- FL Fresh-water marsh, low
- LF Low sand flat
- MU Marshes/sand flats undifferentiated
- SD Salt-water marsh, distal (high)
- SP Salt-water marsh, proximal (low)
- SO Salt-water marsh/open water undifferentiated
- SW Swamps
- LWL Fluvial woodlands, low
- WL Fluvial woodlands
- TA Transitional areas
- WU Wetlands/uplands, undifferentiated
- U Uplands

## **Appendix B**

**APPENDIX B. List of Plant Species by Survey Site number.**

General Location	Site Number on Photo (Aerial photo + location No.)	General descriptions	Site number	Prevalent Species
			Quad No.	Site No.
Follets Island	GHGH 8011 5		1 — 1	<i>OUT OF MAP AREA</i>
			1 — 1	<i>Paspalum urvillei</i> -collected
	GHKL 8098 1		1 — 2	<i>Batis-Monanthochloe</i> community
			1 — 2	<i>Distichlis spicata</i>
			1 — 2	<i>Distichlis-Batis</i> community
			1 — 2	<i>Batis, Borrichia, Spartina alterniflora</i> (around water)
			1 — 2	<i>Distichlis-Monanthochloe</i> , mixed with <i>Batis</i>
			1 — 2	<i>Lycium carolinianum</i> (sparse)
			1 — 2	<i>Juncus roemerianus</i> (along some ponds)
	Disturbed areas (soil)		1 — 2	<i>Iva frutescens</i>
			1 — 2	<i>Spartina patens</i>
			1 — 2	<i>Borrichia frutescens</i>
			1 — 2	<i>Spartina spartinae</i>
			1 — 2	<i>Batis maritima</i>
			1 — 2	<i>Suaeda</i> sp.
			1 — 2	<i>Aster</i> sp.
			1 — 2	<i>Limonium nashii</i>
			1 — 2	<i>Monanthochloe littoralis</i>
			1 — 2	<i>Salicornia</i> sp.
	Depression		1 — 2	<i>Spartina patens-Batis</i>
	Margins of depression		1 — 2	<i>Iva frutescens</i>
			1 — 2	<i>Setaria</i> sp., composites
	GHKL 8098 2		1 — 3	
	near hwy and generators		1 — 3	
	Upland belt of scrub/shrub		1 — 3	<i>Baccharis halimifolia</i>
	Along upper margins of marsh		1 — 3	<i>Iva frutescens-Borrichia</i>
	Bayward transect		1 — 3	<i>Spartina patens</i>
			1 — 3	<i>Scirpus americanus</i>
			1 — 3	<i>Spartina patens</i>
			1 — 3	then <i>Monanthochloe</i>
			1 — 3	<i>Salicornia</i>
			1 — 3	<i>Suaeda</i>
			1 — 3	<i>Limonium</i>
			1 — 3	<i>Batis</i>
			1 — 3	<i>Distichlis</i>
			1 — 3	<i>Aster</i>
	Tidal channel/pond		1 — 3	<i>Batis</i> (dead)
			1 — 3	<i>Scirpus maritimus</i>
			1 — 3	<i>Batis</i>
			1 — 3	<i>Juncus roemerianus</i>
	Higher margins		1 — 3	<i>Iva</i>
			1 — 3	<i>Borrichia</i>
			1 — 3	<i>Spartina patens</i>
			1 — 3	<i>Distichlis</i>
	GHGH 8012 3		3 — 1	
	Elevation transect		3 — 1	
	Higher side of fault		3 — 1	<i>Spartina spartinae</i> (80%)
			3 — 1	<i>Setaria geniculata</i>
			3 — 1	<i>Aster</i> sp.
			3 — 1	<i>Cyperus</i> sp.
	Lower side of fault		3 — 1	
	Flat/emergent		3 — 1	<i>Monanthochloe-Salicornia-Batis</i>
			3 — 1	<i>Distichlis</i> (80%)
			3 — 1	<i>Salicornia</i> (20%)

APPENDIX B (cont.)

General Location	Site Number on Photo (Aerial photo + location No.)	Site number Quad Site No. No.	Prevalent Species
	General descriptions		
		3 — 1	<i>Spartina alterniflora</i>
GHCD 7931 3		3 — 2	SEE HOSKINS MOUND TRANSECT 7 (WINDMILL TRANSECT)
GHGH 8012 1a Lower area Landward margin		3 — 3	<i>Batis-Salicornia bigelovii-Spartina alterniflora</i>
		3 — 3	<i>Monanthochloe-Distichlis-Batis</i>
		3 — 3	<i>Borrichia</i>
		3 — 3	<i>Baccharis halimifolia</i>
		3 — 3	<i>Spartina spartinae</i>
		3 — 3	<i>Spartina patens</i>
Highest assemblage toward Hwy		3 — 3	<i>Batis grades into Spartina alterniflora</i>
		3 — 3	<i>Paspalum monostachyum-collected</i>
		3 — 3	<i>Hydrocotyle bonariensis</i>
		3 — 3	<i>Borrichia</i>
		3 — 3	<i>Fimbristylis</i>
		3 — 3	<i>Andropogon glomeratus</i>
		3 — 3	<i>composites, other species</i>
Follets Island	GHGH 8012 1a SEE SURVEY LINE FOR THIS SITE	3 — 3	
		3 — 3	
Brazoria National Wildlife Refuge			
GHGH 8012 2b		3 — 4	<i>Salicornia-Distichlis</i>
		3 — 4	<i>Batis (scattered)</i>
		3 — 4	<i>Spartina patens (patches)</i>
GHCD 7932 3b Wet areas in distance Higher area above fringing flat		3 — 5	<i>Distichlis - Spartina patens - Paspalum vaginatum dominance</i>
		3 — 5	<i>Phragmites australis</i>
		3 — 5	<i>Spartina spartinae</i>
		3 — 5	<i>Setaria sp.</i>
		3 — 5	<i>Aster sp.</i>
		3 — 5	<i>Cyperus articulatus</i>
		3 — 5	<i>Solidago sp.</i>
		3 — 5	<i>composites</i>
Flat/Emergent area		3 — 5	<i>Distichlis, Salicornia, Monanthochloe</i>
GHCD 7932 3a In water		3 — 6	<i>Distichlis spicata</i>
		3 — 6	<i>Paspalum vaginatum</i>
		3 — 6	<i>Spartina patens dominant in distance</i>
		3 — 6	<i>Salicornia virginica and Suaeda sp.</i>
		3 — 6	<i>Juncus roemerianus</i>
		3 — 6	<i>Spartina alterniflora (small patch)</i>
GHGH 8013 2 Adjacent lower areas Slightly higher Landward Depressions Rims		3 — 7	<i>Spartina alterniflora-Batis-Distichlis</i>
		3 — 7	<i>Spartina alterniflora (100%)</i>
		3 — 7	<i>Distichlis &amp; Batis</i>
		3 — 7	<i>Salicornia</i>
		3 — 7	<i>Distichlis, Batis</i>
		3 — 7	<i>some Borrichia</i>
		3 — 7	<i>Salicornia, and others</i>
		3 — 7	<i>Spartina alterniflora, patches</i>
		3 — 7	<i>Iva frutescens, Borrichia, Spartina spartinae-Dominants</i>
		3 — 7	<i>Spartina patens, Opuntia sp.</i>
GHCD 7932 3		3 — 8	<i>Distichlis spicata dominant</i>
In distance		3 — 8	<i>Scattered Salicornia</i>
		3 — 8	<i>Phragmites australis</i>
		3 — 8	<i>Eleocharis microcarpa</i>
Disturbed area adjacent to site 3		3 — 8	<i>Cynodon dactylon</i>
From higher to lower areas		3 — 8	<i>Andropogon glomeratus</i>
		3 — 8	<i>Machaeranthera</i>

APPENDIX B (cont.)

General Location	Site Number on Photo (Aerial photo + location No.)	General descriptions	Site number Quad No.	Site number Site No.	Prevalent Species
			3 — 8		<i>Borrichia</i>
			3 — 8		<i>Spartina spartinae</i>
			3 — 8		<i>Spartina patens</i>
			3 — 8		<i>Distichlis spicata</i>
			3 — 8		<i>Typha sp.</i>
			3 — 8		<i>Bacopa</i>
			3 — 8		<i>Cyperus sp.</i>
			3 — 8		<i>Eleocharis sp.</i>
			3 — 8		<i>Paspalum vaginatum</i>
Hoskins Mound S.			3 — 9		
	GHCD 7932 2		3 — 9		
	Disturbed (diked) area at well site		3 — 9		
	Saline areas around diked pond		3 — 9		<i>Spartina spartinae, Borrichia, Distichlis, Machaeranthera, Iva Monanthochloe, Salicornia</i>
	In fresher diked area		3 — 9		<i>Typha sp., Bacopa monnieri, Cyperus oxylepis-collected</i>
			3 — 9		<i>Iva frutescens, Borrichia, Distichlis, Spartina spartinae</i>
	Flat/Emergent south of diked area		3 — 9		<i>Fimbristylis castanea, Andropogon glomeratus, Monanthochloe</i>
	Adjacent to ICWW		3 — 9		<i>Salicornia-Monanthochloe dominant</i>
	Dark patches in water		3 — 9		<i>Distichlis spicata (dominant)-Spartina alterniflora-Batis</i>
	(No sea grasses in drift line)		3 — 9		<i>Brown algae</i>
	SE corner of diked area		3 — 9		<i>Spartina patens</i>
	GHCD 7932 2a		3 — 10		
	Flat/Emergent assemblage		3 — 10		<i>Monanthochloe, Salicornia, Spartina spartinae, Batis, Limonium, Borrichia</i>
			3 — 10		
	GHGH 8013 1a		3 — 11		Patches of vegetation included
	sand flats/emergents		3 — 11		<i>Monanthochloe</i>
			3 — 11		<i>Batis</i>
			3 — 11		<i>Salicornia</i>
			3 — 11		Algal mats on flats, damp soils near vegetation
	GHGH 8014 1		3 — 12		
	Brackish/Intermediate		3 — 12		<i>Typha - dominant</i>
	Gufward of Rd.		3 — 12		<i>Juncus</i>
			3 — 12		<i>Scirpus americanus</i>
			3 — 12		<i>Spartina patens</i>
			3 — 12		<i>Phragmites</i>
			3 — 12		<i>Paspalum vaginatum</i>
			3 — 12		<i>Baccharis halimifolia</i>
	Salt/brackish		3 — 12		<i>Spartina patens dominant</i>
	Bayward of Rd.		3 — 12		<i>Scirpus americanus</i>
			3 — 12		<i>Juncus roemerianus</i>
			3 — 12		<i>Borrichia</i>
	Adjacent area		3 — 12		
	grading from		3 — 12		<i>Spartina patens - dominant</i>
			3 — 12		<i>Batis</i>
			3 — 12		<i>Juncus roemerianus</i>
			3 — 12		<i>Scirpus maritimus</i>
			3 — 12		<i>Batis</i>
	GHGH 8011 3		4 — 1		Uplands
	Water		4 — 1		Trees appear dead
	Margin of water		4 — 1		<i>Sesbania, Celtis sp., Sapium sebiferum</i>
			4 — 1		<i>Andropogon glomeratus</i>
Lake Jackson Area			4 — 2		
	GHGH 8011 1		4 — 2		<i>Scirpus californicus dominant</i>
	Stubblefield Lake		4 — 2		<i>Nelumbo lutea</i>

APPENDIX B (cont.)

General Location	Site Number on Photo (Aerial photo + location No.)	Site number		Prevalent Species
		Quad General descriptions	Site No.	
	On margin of lake		4 — 2	<i>Phragmites australis</i>
			4 — 2	<i>Polygonum sp.</i>
			4 — 2	<i>Bacopa monnieri</i>
			4 — 2	<i>Salix nigra</i>
			4 — 2	<i>Sesbania sp.</i>
			4 — 2	<i>Cyperus articulatus</i>
			4 — 2	<i>Scirpus americanus</i>
			4 — 2	<i>Andropogon glomeratus</i>
			4 — 2	<i>Spartina spartinae</i>
			4 — 2	<i>Spartina patens</i>
			4 — 2	<i>Aster sp.</i>
			4 — 2	<i>Typha sp.</i>
			4 — 2	<i>Setaria sp.</i>
			4 — 2	<i>Solidago sp.</i>
			4 — 2	<i>Baccharis halimifolia</i>
	GHGH 8011 2		4 — 3	
	Ditch has drained water		4 — 3	<i>Sesbania sp.</i>
			4 — 3	<i>Cyperus sp.</i>
			4 — 3	<i>Cynodon dactylon (probably)</i>
			4 — 3	
	GHGH 8011 4		4 — 4	<i>Spartina spartinae</i>
			4 — 4	<i>Scirpus or Juncus</i>
			4 — 4	<i>Ulmus crassifolia</i>
			4 — 4	<i>Celtis laevigata</i>
			4 — 4	<i>Quercus virginiana</i>
			4 — 4	<i>Sabal minor</i>
			4 — 4	<i>Sapium sebiferum</i>
			4 — 4	<i>Baccharis halimifolia ?</i>
	GHGH 8012 5		4 — 5	<i>Scirpus californicus dominant</i>
	GHGH 8012 4		4 — 6	<i>Spartina spartinae dominant</i>
	GHGH 8012 3a		4 — 7	
	Brackish/Intermediate		4 — 7	<i>Paspalum vaginatum</i>
			4 — 7	<i>Typha sp.</i>
			4 — 7	<i>Scirpus olneyi</i>
			4 — 7	<i>Spartina patens</i>
			4 — 7	<i>Echinochloa crusgalli</i>
			4 — 7	<i>Spartina spartinae</i>
			4 — 7	<i>Aster</i>
			4 — 7	<i>Cyperus sp.</i>
Galveston Island	GHEF 7947 2		5 — 1	<i>Salicornia</i>
			5 — 1	<i>Spartina patens</i>
			5 — 1	<i>Borrichia</i>
			5 — 1	<i>Iva frutescens</i>
			5 — 1	<i>Aster</i>
			5 — 1	<i>Batis (along channel)</i>
	GHEF 7946 2a		5 — 2	<i>Spartina alterniflora</i>
			5 — 2	<i>Juncus roemerianus</i>
	GHEF 7945 9		5 — 3	
	Heavily grazed, grass unidentified		5 — 3	<i>Cynodon dactylon possibly</i>
	Across road (southwest)		5 — 3	<i>Scirpus californicus</i>
	small ponded area		5 — 3	<i>Bacopa monnieri</i>
			5 — 3	<i>Cyperus articulatus</i>
			5 — 3	<i>Sesbania sp.</i>
			5 — 3	<i>Cynodon dactylon</i>

**APPENDIX B (cont.)**

General Location	Site Number on Photo (Aerial photo + location No.)	General descriptions	Site number Quad No.	Site number Site No.	Prevalent Species
	GHEF 7945 10		5 — 4		
	Lower area (southwest)		5 — 4		<i>Distichlis</i> dominant
	surrounding higher flats		5 — 4		<i>Salicornia</i>
	Toward northeast of road		5 — 4		<i>Distichlis</i> dominant
	On flats		5 — 4		<i>Salicornia</i>
			5 — 4		<i>Distichlis</i>
			5 — 4		<i>Machaeranthera</i>
			5 — 4		<i>Limonium</i>
			5 — 4		<i>Borrichia</i>
			5 — 4		<i>Monanthochloe</i>
	GHEF 7947 4b		5 — 5		
	along channel		5 — 5		<i>Spartina alterniflora</i> dominant
	grading upward above channel		5 — 5		<i>Distichlis</i> , <i>Spartina patens</i> , <i>Juncus roemarianus</i>
	higher zones		5 — 5		<i>Iva frutescens</i> , <i>Spartina spartinae</i> , <i>Spartina patens</i>
			5 — 5		<i>Andropogon</i> , <i>Setaria</i> , <i>Hydrocotyle</i>
	GHEF 7948 7		5 — 6		<i>Spartina alterniflora</i> dominant
	above smooth cordgrass		5 — 6		<i>Batis</i> , <i>Salicornia</i> , Scattered <i>Distichlis</i>
	near road		5 — 6		<i>Juncus roemarianus</i> , <i>Spartina patens</i>
	along road		5 — 6		clumps of <i>Baccharis</i> , <i>Iva</i> , and <i>Spartina spartinae</i>
	GHEF 7948 3a		6 — 1		<i>Suaeda linearis</i>
	Spoil island-local algal flat and patches of emergent vegetation		6 — 1		<i>Batis maritima</i>
			6 — 1		<i>Spartina patens</i>
			6 — 1		<i>Spartina spartinae</i>
			6 — 1		<i>Borrichia frutescens</i>
			6 — 1		<i>Iva frutescens</i>
			6 — 1		<i>Limonium nashii</i>
			6 — 1		<i>Opuntia</i> in higher areas
			6 — 1		<i>Setaria</i>
			6 — 1		<i>Cynodon dactylon</i>
			6 — 1		<i>Distichlis spicata</i>
			6 — 1		<i>Acacia angustissima</i>
			6 — 1		<i>Salicornia bigelovii</i>
			6 — 1		<i>Iva annua</i>
			6 — 1		other composites
	GHEF 7934 1		6 — 2		<i>Spartina alterniflora</i> dominant (100%)
Hoskins Mound area					
	GHCD 7932 1		7 — 1		<i>Cyperus oxylepis</i> -collected
	SEE SURVEY LINE FOR THIS SITE		7 — 1		<i>Iva augustifolia</i> -collected
			7 — 1		<i>Cyperus virens</i> -collected
			7 — 1		<i>Paspalum floridanum</i> -collected
			7 — 1		<i>Andropogon glomeratus</i> -collected
			7 — 1		<i>Eragrostis spectabilis</i> -collected
			7 — 1		<i>Eleocharis cellulosa</i> -collected
Chocolate Bayou area					
	GHEF 7951 1		7 — 2		<i>Juncus roemerianus</i> dominant
			7 — 2		<i>Spartina alterniflora</i>
			7 — 2		<i>Scirpus maritimus</i>
	Away from water		7 — 2		<i>Distichlis spicata</i> dominant
			7 — 2		<i>Spartina patens</i>
			7 — 2		Scattered <i>Aster</i> sp.
	vegetation/flat mix		7 — 2		<i>Distichlis spicata/dry flats</i>
Galveston Island					
	GHEF 7945 1		9 — 1		
	low marsh		9 — 1		<i>Distichlis spicata-Spartina alterniflora</i> assemblage

APPENDIX B (cont.)

General Location	Site Number on Photo (Aerial photo + location No.) General descriptions	Site number Quad Site No. No.	Prevalent Species
		9 — 1	<i>Spartina alterniflora</i> abundance increases toward bayou
		9 — 1	<i>Salicornia bigelovii</i>
high marsh		9 — 1	<i>Borrichia frutescens</i> (dominant)
		9 — 1	<i>Spartina spartinae</i> (scattered)
		9 — 1	<i>Machaeranthera phyllocephala</i>
		9 — 1	<i>Fimbristylis castanea</i>
		9 — 1	<i>Solidago</i> sp.
sand flat/emergent mix		9 — 1	<i>Salicornia</i>
		9 — 1	<i>Batis</i>
		9 — 1	<i>Limonium nashii</i>
		9 — 1	<i>Suaeda</i> sp.
		9 — 1	<i>Monanthochloe littoralis</i>
		9 — 1	<i>Lycium carolinianum</i>
fresher small marsh near road		9 — 1	<i>Typha</i> sp.
GHAB 7868 2		9 — 2	<i>Borrichia frutescens</i> - <i>Distichlis spicata</i> dominance
		9 — 2	<i>Limonium nashii</i>
		9 — 2	<i>Suaeda</i> sp.
		9 — 2	<i>Salicornia bigelovii</i>
Pelican Island			
	GHAB 7868 1 and area	9 — 3	<i>Borrichia frutescens</i>
		9 — 3	<i>Distichlis spicata</i>
		9 — 3	<i>Machaeranthera</i>
	Depressions	9 — 3	<i>Typha</i> sp.
		9 — 3	<i>Scirpus maritimus</i>
	Trees and shrubs on Island include	9 — 3	<i>Sapium sebiferum</i>
		9 — 3	<i>Tamarix gallica</i>
		9 — 3	<i>Celtis</i> sp.
		9 — 3	<i>Sesbania</i> spp.
		9 — 3	<i>Baccharis halimifolia</i>
		9 — 3	<i>Iva frutescens</i>
Virginia Point Quad	GHEF 7948 2c	10 — 1	<i>Distichlis spicata</i> 35% water depth 6-7cm
		10 — 1	<i>Spartina alterniflora</i> 60%
		10 — 1	<i>Batis maritima</i> 5%
	GHEF 7948 2b	10 — 2	<i>Distichlis spicata</i> 60% water depth 3cm
		10 — 2	<i>Spartina alterniflora</i> 40%
	GHEF 7948 2a transect	10 — 3	<i>Distichlis spicata</i> 75% water depth 1cm
		10 — 3	<i>Spartina alterniflora</i> 15%
		10 — 3	<i>Batis maritima</i> 5%
	GHEF 7948 1c	10 — 4	<i>Spartina alterniflora</i> 100%
	GHEF 7948 1b	10 — 5	<i>Distichlis spicata</i> 60%
		10 — 5	<i>Spartina alterniflora</i> 40%
		10 — 5	<i>Salicornia</i> 1%
Mainland shore			
West Bay	GHEF 7948 1a transect	10 — 6	<i>Batis maritima</i>
		10 — 6	<i>Borrichia frutescens</i>
		10 — 6	<i>Limonium nashii</i>
		10 — 6	<i>Spartina spartinae</i>
		10 — 6	<i>Lycium carolinianum</i>
West of Jones Bay			
	GHAB 7870 2a	10 — 7	<i>Spartina alterniflora</i> (dominant, 100%)
	GHAB 7870 2b	10 — 8	<i>Spartina alterniflora</i> (dominant, > 90%)
		10 — 8	<i>Salicornia</i> sp.

APPENDIX B (cont.)

General Location	Site Number on Photo (Aerial photo + location No.) General descriptions	Site number Quad Site No. No.	Prevalent Species
		10 — 8	<i>Distichlis spicata</i>
		10 — 8	<i>Batis maritima</i>
GHEF 7945 6 (GHEF 7946 3b)		10 — 9	<i>Spartina alterniflora</i> (dominant, > 90%)
		10 — 9	scattered <i>Batis</i> , <i>Distichlis</i> , <i>Salicornia</i> locally
GHAB 7870 3		10 — 10	<i>Juncus roemerianus</i>
		10 — 10	<i>Batis maritima</i>
		10 — 10	<i>Distichlis spicata</i>
		10 — 10	<i>Scirpus maritimus</i>
		10 — 10	Moist algal flats
Texas City area			
GHAB 7870 1 (east side of highway)		10 — 11	<i>Spartina alterniflora</i> (dominant)
		10 — 11	<i>Scirpus maritimus</i>
		10 — 11	<i>Distichlis spicata</i>
		10 — 11	<i>Juncus roemerianus</i>
		10 — 11	<i>Iva frutescens</i>
		10 — 11	<i>Aster sp.</i>
	(Ponds on west side of highway)	10 — 11	<i>Typha sp.</i>
GHEF 7945 7 flat/emergent mix (Southwest) In distance		10 — 12	
		10 — 12	<i>Monanthochloe</i> , <i>Salicornia</i> spp., <i>Suaeda</i> , <i>Limonium</i> , <i>Lycium</i>
		10 — 12	<i>Spartina patens</i> dominant
		10 — 12	<i>Juncus roemerianus</i> (less dominant)
		10 — 12	<i>Scirpus californicus</i> (patch)
		10 — 12	<i>Eleocharis cellulosa</i> -collected
GHEF 7945 8 (Southwest)		10 — 13	
		10 — 13	<i>Spartina patens-Juncus roemerianus</i> dominant
		10 — 13	<i>Tamarix</i> nearby
	Standing water across road higher areas	10 — 13	<i>Bacopa</i> on margins of water
		10 — 13	<i>Spartina patens-Juncus roemerianus</i> dominant
		10 — 13	<i>Iva frutescens</i>
GHEF 7945 3		10 — 14	<i>Juncus roemerianus</i> (dominant in some areas)
		10 — 14	<i>Distichlis spicata</i> (codominant with <i>S. patens</i> locally)
		10 — 14	<i>Spartina patens</i>
	on flats in distance	10 — 14	<i>Monanthochloe littoralis</i>
		10 — 14	<i>Batis</i>
		10 — 14	<i>Salicornia</i> spp.
GHEF 7945 4		10 — 15	
flats on swale		10 — 15	<i>Monanthochloe littoralis</i> (dominant)
		10 — 15	<i>Salicornia bigelovii</i>
		10 — 15	<i>Distichlis spicata</i> -collected
		10 — 15	<i>Batis</i>
	flanks of swale in distance	10 — 15	<i>Spartina alterniflora</i>
	fringing water	10 — 15	<i>Monanthochloe littoralis</i>
	slightly higher marsh near road lows	10 — 15	<i>Batis-Borrichia-Distichlis</i>
	slightly higher	10 — 15	<i>Iva-Spartina spartinae</i>
	slightly higher	10 — 15	<i>Suaeda</i> sp.
	other less abundant species	10 — 15	<i>Lycium carolinianum</i>
		10 — 15	<i>Limonium nashii</i>
		10 — 15	<i>Juncus roemerianus</i>
	ridge assemblage near flat	10 — 15	<i>Spartina spartinae</i>
		10 — 15	<i>Spartina patens</i>
		10 — 15	<i>Iva frutescens</i>
		10 — 15	<i>Juncus roemerianus</i>
	toward bay	10 — 15	<i>Spartina patens-Juncus-Iva assemblage</i>

APPENDIX B (cont.)

General Location	Site Number on Photo (Aerial photo + location No.)	General descriptions	Site number Quad Site No. No.	Prevalent Species
GHEF 7945 5		saltier assemblage near road and NE	10 — 16	
			10 — 16	<i>Distichlis-Batis-Salicornia</i>
			10 — 16	<i>Juncus, Iva, Spartina patens, Limonium</i>
			10 — 16	
		fresher west of dike	10 — 16	<i>Spartina patens</i>
			10 — 16	<i>Distichlis spicata</i>
			10 — 16	<i>Bacopa monnieri</i>
			10 — 16	<i>Sesbania sp.</i>
			10 — 16	<i>Typha sp.</i>
			10 — 16	<i>Scirpus californicus ? (in distance)</i>
			10 — 16	<i>Paspalum vaginatum (probable)</i>
		swale across road (NE)	10 — 16	
		flat/emergent mix	10 — 16	<i>Distichlis, Salicornia, short S. patens, Suaeda,</i>
			10 — 16	<i>Machærantha and Cynodon dactylon</i>
GHEF 7945 2			10 — 17	
		sand flat/emergent mix	10 — 17	<i>Salicornia bigelovii</i>
			10 — 17	<i>Salicornia virginica</i>
			10 — 17	<i>Distichlis spicata</i>
			10 — 17	<i>Limonium nashii</i>
		slightly higher	10 — 17	<i>Borrichia frutescens</i>
			10 — 17	<i>Spartina spartinae</i>
			10 — 17	<i>Juncus</i>
GHAB 7873 1			11 — 1	
Willow Bayou			11 — 1	
Forested margin			11 — 1	<i>Sapium sebiferum</i>
			11 — 1	<i>Salix nigra</i>
			11 — 1	<i>Celtis laevigata</i>
		Along stream	11 — 1	<i>Alternanthera philoxeroides</i>
			11 — 1	<i>Panicum dichotomiflorum</i>
			11 — 1	<i>Sagittaria sp.</i>
			11 — 1	<i>Sesbania sp.</i>
			11 — 1	<i>Ambrosia sp.</i>
Hitchcock area				
GHAB 7872 3 and 3a			11 — 2	<i>Pinus sp.</i>
Highland Bayou			11 — 2	<i>Ulmus crassifolia</i>
			11 — 2	<i>Quercus virginiana</i>
			11 — 2	<i>Ilex vomitoria</i>
			11 — 2	<i>Carya illinoensis</i>
			11 — 2	<i>Platanus occidentalis</i>
			11 — 2	<i>Salix nigra</i>
			11 — 2	<i>Juniper</i>
GHGH 7513 1a			12 — 1	
Low areas			12 — 1	<i>Spartina alterniflora (dominant in lows)</i>
Slightly higher			12 — 1	<i>Distichlis spicata (dominant overall)</i>
			12 — 1	<i>Spartina patens (abundant)</i>
			12 — 1	<i>Aster, Batis, Borrichia (scattered)</i>
Flat			12 — 1	<i>Monanthochloe littoralis (dominant on flats)</i>
Higher areas			12 — 1	<i>Spartina spartinae, Borrichia, Iva, Lycium, Limonium</i>
			12 — 1	<i>Salicornia, Suaeda, Machærantha, Solidago</i>
GHIJ 7568 2a			13 — 1	
Higher flanks of swale			13 — 1	<i>Spartina patens (60-70%)-Borrichia frutescens (30-40%)</i>
Edge of flats			13 — 1	<i>Juncus roemerianus</i>
Flats			13 — 1	<i>Monanthochloe littoralis, Salicornia spp., Distichlis, Batis</i>
In distance toward Boliv. Rds			13 — 1	<i>Spartina alterniflora</i>
Beach ridge-- prairie assemblage			13 — 1	<i>Andropogon glomeratus, Dichromena colorata, Fimbristylis</i>
			13 — 1	<i>castanea, Iva frutescens, Solidago sp., Aristida sp., Paspalum</i>
			13 — 1	<i>monostachym, other composites</i>

## APPENDIX B (cont.)

General Location	Site Number on Photo (Aerial photo + location No.)	General descriptions	Site number Quad No.	Site number Site No.	Prevalent Species
		Next swale (gulfward but cutoff from marine waters--no flat)	13 — 1	13 — 1	<i>Spartina patens, Distichlis, Batis, Juncus in lows</i>
			13 — 1	13 — 1	<i>Scirpus americanus locally abundant</i>
			13 — 1	13 — 1	<i>Borrichia, Limonium, Lycium</i>
Bolivar Peninsula					
	GHIJ 7568 1		13 — 2	13 — 2	<i>Juncus roemerianus-Batis dominant</i>
	Edge of flat		13 — 2	13 — 2	<i>Salicornia</i>
	In depression		13 — 2	13 — 2	<i>Spartina alterniflora scattered</i>
	Higher areas		13 — 2	13 — 2	<i>Borrichia frutescens</i>
	GHIJ 7569 1a		13 — 4	13 — 4	<i>Spartina alterniflora dominant</i>
	higher prairie		13 — 4	13 — 4	<i>Spartina spartinae</i>
	GHIJ 7569 1		13 — 5	13 — 5	<i>Typha, Cyperus articulatus, Hydrocotyle</i>
			13 — 5	13 — 5	<i>Scirpus americanus, Sesbania</i>
	Higher clumps		13 — 5	13 — 5	<i>Setaria</i>
	Wetter, narrow zone in swale		13 — 5	13 — 5	<i>Scirpus californicus (?) in distance</i>
			13 — 5	13 — 5	<i>Polygonum hydropiperoides-collected</i>
	GHIJ 7569 2		13 — 6	13 — 6	<i>Distichlis dominant, Spartina alterniflora in lows</i>
			13 — 6	13 — 6	<i>Batis, Aster, Borrichia</i>
	GHIJ 7565 4		15 — 1	15 — 1	<i>Spartina alterniflora (codominant with Juncus in low marsh)</i>
	(low marsh to higher areas)		15 — 1	15 — 1	<i>Juncus roemerianus (codominant with S. alterniflora)</i>
			15 — 1	15 — 1	<i>Distichlis spicata</i>
			15 — 1	15 — 1	<i>Salicornia sp.</i>
			15 — 1	15 — 1	<i>Spartina patens</i>
			15 — 1	15 — 1	<i>Borrichia frutescens</i>
			15 — 1	15 — 1	<i>Iva frutescens</i>
			15 — 1	15 — 1	<i>Aster tenuifolius ?</i>
			15 — 1	15 — 1	<i>Lycium carolinium</i>
			15 — 1	15 — 1	<i>Spartina spartinae</i>
			15 — 1	15 — 1	<i>Andropogon glomeratus</i>
			15 — 1	15 — 1	<i>Cynodon dactylon</i>
	GHIJ 7565 1		15 — 2	15 — 2	<i>Spartina alterniflora (dominant)</i>
	GHIJ 7565 3		15 — 3	15 — 3	<i>Spartina alterniflora (dominant)</i>
	(edge of Moses Lake)		15 — 3	15 — 3	<i>Juncus roemerianus (patch)</i>
			15 — 3	15 — 3	<i>Distichlis spicata</i>
			15 — 3	15 — 3	<i>Spartina patens</i>
			15 — 3	15 — 3	<i>Iva frutescens</i>
			15 — 3	15 — 3	<i>Borrichia frutescens</i>
			15 — 3	15 — 3	<i>Aster sp.</i>
			15 — 3	15 — 3	<i>Limonium nashii</i>
			15 — 3	15 — 3	<i>Salicornia sp.</i>
			15 — 3	15 — 3	<i>Spartina spartinae</i>
	GHIJ 7565 5 (Factory Bayou)		15 — 4	15 — 4	<i>Scirpus maritimus</i>
			15 — 4	15 — 4	<i>Juncus roemerianus</i>
			15 — 4	15 — 4	<i>Distichlis spicata</i>
			15 — 4	15 — 4	<i>Spartina alterniflora (margins of channel)</i>
	high marsh		15 — 4	15 — 4	<i>Iva frutescens</i>
			15 — 4	15 — 4	<i>Spartina patens</i>
			15 — 4	15 — 4	<i>Distichlis spicata</i>
			15 — 4	15 — 4	<i>Spartina spartinae</i>
	others		15 — 4	15 — 4	<i>Limonium nashii</i>
			15 — 4	15 — 4	<i>Lycium carolinium</i>
			15 — 4	15 — 4	<i>Phragmites australis</i>
	mud flats (low tide)		15 — 4	15 — 4	

APPENDIX B (cont.)

General Location	Site Number on Photo (Aerial photo + location No.)	Site number		Prevalent Species
		Quad	Site	
	General descriptions	No.	No.	
San Leon		15 — 5		This site number was not used
	GHIJ 7565 6  (forested area mostly willow)	15 — 6		<i>Eleocharis quadrangulata</i> (dominant, 90%)
		15 — 6		<i>Sesbania</i> sp.
		15 — 6		<i>Salix nigra</i>
		15 — 6		<i>Sapium sebiferum</i>
		15 — 6		<i>Hymenocallis caroliniana</i>
	GHIJ 7565 2a  (exact location not confirmed because of new housing develop.) Site in relatively small drainage.	15 — 7		<i>Typha</i> sp.
		15 — 7		<i>Rhynchospora</i> sp.
		15 — 7		<i>Panicum</i> sp.
		15 — 7		<i>Sesbania</i> sp.
		15 — 7		<i>Andropogon glomeratus</i>
		15 — 7		<i>Setaria</i> sp.
		15 — 7		<i>Aristida</i> sp.
		15 — 7		<i>Aster tenuifolius?</i>
	GHIJ 7566 1  (from low to high marsh)	15 — 8		<i>Spartina alterniflora</i> (dominant)
		15 — 8		<i>Distichlis spicata</i> (abundant)
		15 — 8		<i>Scirpus maritimus</i> (abundant)
		15 — 8		<i>Spartina patens</i> (abundant)
		15 — 8		<i>Iva frutescens</i> (higher fringe)
	GHIJ 7566 2  mud/sand flat	15 — 9		<i>Distichlis spicata</i> (dominant)
		15 — 9		<i>Spartina alterniflora</i> (dominant near water)
		15 — 9		<i>Salicornia virginica</i>
		15 — 9		<i>Salicornia bigelovii</i>
		15 — 9		<i>Monanthochloe littoralis</i>
		15 — 9		<i>Limonium nashii</i>
		15 — 9		<i>Suaeda</i> sp.
	higher marsh	15 — 9		<i>Iva frutescens</i>
		15 — 9		<i>Spartina spartinae</i>
		15 — 9		<i>Borrichia frutescens</i>
	margin of pond near road	15 — 9		<i>Scirpus maritimus</i>
Dickinson area				
	GHIJ 7563 1	16 — 1		<i>Ulmus crassifolia</i>
		16 — 1		<i>Ilex vomitoria</i>
		16 — 1		<i>Celtis laevigata</i>
		16 — 1		<i>Sabal minor</i>
		16 — 1		<i>Quercus nigra</i>
		16 — 1		<i>Pinus taeda</i>
		16 — 1		<i>Fraxinus</i> sp.
		16 — 1		<i>Liquidambar styraciflua</i>
	GHIJ 7563 1a  Magnolia Bayou	16 — 2		<i>Quercus phellos</i>
		16 — 2		<i>Quercus nigra</i>
		16 — 2		<i>Ilex vomitoria</i>
		16 — 2		<i>Ulmus crassifolia</i>
		16 — 2		<i>Quercus falcata</i>
		16 — 2		<i>Sabal minor</i>
		16 — 2		<i>Pinus taeda</i>
	GHIJ 7564 2	16 — 3		<i>Iva frutescens</i> dominant
		16 — 3		<i>Spartina spartinae</i>
		16 — 3		<i>Phragmites australis</i>
		16 — 3		<i>Solidago</i> sp.
		16 — 3		<i>Cynodon dactylon</i>
		16 — 3		<i>Ambrosia</i> sp.
		16 — 3		<i>Ilex vomitoria</i>

APPENDIX B (cont.)

General Location	Site Number on Photo (Aerial photo + location No.)	General descriptions	Site number	Prevalent Species
			Quad	Site No.
			16 — 3	<i>Parkinsonia aculeata</i>
			16 — 3	<i>Ulmus crassifolia</i>
			16 — 3	<i>Sapium sebiferum</i>
High Island area	BPA-GH 7510 1b		17 — 1	<i>Spartina patens</i> dominant
			17 — 1	<i>Distichlis</i>
	Along channel		17 — 1	<i>Typha, Bacopa, Paspalum lividum ?</i>
	Back toward hwy		17 — 1	<i>Scirpus olneyi</i> patch
High Island Area	BPA-GH 7511 1		17 — 2	<i>Spartina patens - Scirpus maritimus</i> co-dominant
			17 — 2	<i>Distichlis</i> abundant
			17 — 2	<i>Scattered Aster, Phragmites, Spartina alterniflora</i>
	BPA-GH 7510 4		17 — 3	
	West of ICWW near High Island		17 — 3	<i>Spartina patens - Typha mix</i>
			17 — 3	<i>Scirpus olneyi</i>
			17 — 3	<i>Distichlis</i> abundant
	Near ICWW		17 — 3	<i>Phragmites</i>
	GHGH 7513 4c		18 — 1	<i>Distichlis spicata</i>
			18 — 1	<i>Spartina patens</i>
			18 — 1	patches of <i>Scirpus maritimus</i>
			18 — 1	patches of <i>Juncus roemerianus</i>
	GHGH 7513 4b		18 — 2	<i>Spartina alterniflora</i>
			18 — 2	<i>Scirpus maritimus</i>
	GHGH 7513 4a		18 — 3	<i>Scirpus maritimus</i>
			18 — 3	<i>Spartina patens</i>
			18 — 3	small <i>Borrichia</i>
			18 — 3	<i>Spartina alterniflora</i>
	GHGH 7513 3a		18 — 4	<i>Scirpus maritimus</i>
			18 — 4	<i>Spartina patens</i>
			18 — 4	<i>Spartina alterniflora</i>
			18 — 4	scattered aster
	GHGH 7513 3b		18 — 5	<i>Spartina alterniflora</i>
			18 — 5	<i>Distichlis spicata</i>
	—		—	
	GHGH 7513 2b		18 — 6	<i>Spartina alterniflora</i>
			18 — 6	scattered <i>Scirpus maritimus</i>
			18 — 6	scattered <i>Distichlis spicata</i>
Bolivar Peninsula	GHGH 7513 2a		18 — 7	<i>Scirpus maritimus</i>
			18 — 7	<i>Spartina patens</i>
			18 — 7	<i>Spartina alterniflora</i>
			18 — 7	<i>Salicornia</i> sp.
Anahuac National Wildlife Refuge	GHGH 7513 1		18 — 8	
	Low Brackish/Intermediate		18 — 8	<i>Scirpus olneyi</i>
			18 — 8	<i>Typha</i>
			18 — 8	<i>Spartina patens</i>
			18 — 8	<i>Scirpus maritimus</i>
	Higher marsh near bay		18 — 8	<i>Spartina spartinae</i>
			18 — 8	<i>Setaria geniculata</i>
Anahuac NWR	BPA GH 7512 2		18 — 9	SEE MARSH PROFILE
			18 — 9	<i>Echinochloa crusgalli</i> -collected

APPENDIX B (cont.)

General Location	Site Number on Photo (Aerial photo + location No.)	Site number Quad Site No. No.	Prevalent Species
	General descriptions		
		18 — 9	<i>Panicum virgatum</i> -collected
		18 — 9	<i>Paspalum vaginatum</i> -collected
	GHGH 7512 5b	18 — 10	<i>Spartina spartinae</i>
		18 — 10	<i>Spartina patens</i>
		18 — 10	<i>Iva frutescens</i>
		18 — 10	<i>Borrichia frutescens</i>
		18 — 10	<i>Sporobolus virginicus</i>
		18 — 10	<i>Scirpus olneyi</i>
		18 — 10	<i>Scirpus americanus</i>
		18 — 10	<i>Juncus effusus</i>
		18 — 10	others collected
Frozen Point			
	GHGH 7512 5a	18 — 11	<i>Spartina patens</i>
		18 — 11	<i>Scirpus maritimus</i>
		18 — 12	
	Northeast (flat/emergents)	18 — 12	<i>Monanthochloe</i> , <i>Salicornia</i> spp., <i>Limonium</i>
	Higher mounds	18 — 12	<i>Batis</i> , some <i>Suaeda</i> , <i>Spartina spartinae</i>
	Toward bay	18 — 12	<i>Spartina alterniflora</i>
		18 — 12	patches of <i>Spartina spartinae</i> , <i>S. patens</i>
	Toward bay, wet conditions	18 — 12	<i>Distichlis</i> dominant, <i>Spartina patens</i> , <i>Spartina alterniflora</i>
		18 — 12	<i>Batis</i> in distance
Smith Point Area			
	GHCD 7464 1	19 — 1	
	Brackish/Intermediate	19 — 1	<i>Spartina patens</i>
		19 — 1	<i>Spartina spartinae</i>
		19 — 1	<i>Scirpus maritimus</i>
		19 — 1	<i>Juncus roemerianus</i>
		19 — 1	<i>Phragmites</i>
		19 — 1	<i>Spartina cynosuroides</i>
		19 — 1	<i>Paspalum vaginatum</i>
		19 — 1	<i>Typha</i>
	GHGH 7516 1	20 — 1	<i>Distichlis</i>
		20 — 1	<i>Spartina alterniflora</i>
		20 — 1	<i>Juncus roemerianus</i>
		20 — 1	<i>Scirpus maritimus</i>
		20 — 1	<i>Spartina patens</i>
		20 — 1	<i>Borrichia</i>
		20 — 1	<i>Spartina spartinae</i>
		20 — 1	<i>Iva frutescens</i>
Spoil Islands along Houston Ship Channel			
	GHCD 7469 2	21 — 1	<i>Borrichia frutescens</i>
		21 — 1	<i>Tamarix</i>
		21 — 1	<i>Sesbania drummondii</i>
		21 — 1	<i>Baccharis halimifolia</i>
		21 — 1	<i>Teucrium cubense</i>
		21 — 1	<i>Solidago altissima</i>
		21 — 1	<i>Acacia angustissima</i>
		21 — 1	<i>Ambrosia psilostachya</i>
	GHCD 7469 3	21 — 3	<i>Distichlis spicata</i>
		21 — 3	<i>Spartina alterniflora</i>
		21 — 3	<i>Spartina patens</i>
		21 — 3	<i>Borrichia frutescens</i>
		21 — 3	<i>Salicornia</i> sp.
Clear Creek	GHGH 7522 1 east of highway	22 — 1	<i>Baccharis halimifolia</i> -collected
		22 — 1	<i>Spartina patens</i>
		22 — 1	<i>Distichlis spicata</i>
		22 — 1	<i>Scirpus maritimus</i>

**APPENDIX B (cont.)**

General Location	Site Number on Photo (Aerial photo + location No.)	Site number		Prevalent Species
		Quad No.	Site No.	
		22 — 1		<i>Iva frutescens</i>
		22 — 1		<i>Solidago sp.</i>
		22 — 1		<i>Borrichia frutescens</i>
	west of highway	22 — 1		<i>Spartina patens</i> (dominant)
		22 — 1		<i>Iva frutescens</i>
		22 — 1		<i>Andropogon glomeratus</i>
		22 — 1		<i>Setaria sp.</i>
		22 — 1		<i>Solidago sp.</i>
		22 — 1		<i>Lycium carolinianum</i>
	wetter areas	22 — 1		<i>Typha sp.</i>
		22 — 1		<i>Scirpus maritimus</i>
Armand Bayou Bay Area Park	GHCD 7471 1	22 — 2		<i>Sagittaria sp.</i>
		22 — 2		<i>Polygonum sp.</i>
		22 — 2		<i>Scirpus maritimus</i>
		22 — 2		<i>Spartina patens</i>
		22 — 2		<i>Vigna luteola</i>
		22 — 2		<i>Iva frutescens</i>
		22 — 2		<i>Aster sp.</i>
	forested area	22 — 2		<i>Echinochloa crusgalli</i> -collected
		22 — 2		<i>Sabal minor</i>
		22 — 2		<i>Ulmus crassifolia</i>
		22 — 2		<i>Celtis laevigata</i>
		22 — 2		<i>Ilex vomitoria</i>
		22 — 2		<i>Carya illinoensis</i>
		22 — 2		<i>Pinus sp.</i>
		22 — 2		<i>Quercus aquatica</i>
		22 — 2		<i>Quercus phellos</i>
		22 — 2		<i>Ulmus americana</i>
A. B. Nature Center GHCD 7471 2		22 — 3		<i>Spartina patens</i>
		22 — 3		<i>Spartina spartinae</i>
		22 — 3		<i>Scirpus maritimus</i>
		22 — 3		<i>Iva frutescens</i>
		22 — 3		<i>Spartina alterniflora</i> (near water)
		22 — 3		<i>Cyperus sp.</i>
		22 — 3		<i>Solidago sp.</i>
Taylor Bayou at Port Rd.		22 — 4		<i>Quercus phellos</i>
	GHCD 7470 1	22 — 4		<i>Ulmus crassifolia</i>
		22 — 4		<i>Ilex vomitoria</i>
		22 — 4		<i>Fraxinus sp.</i>
	In water	22 — 4		<i>Scirpus maritimus</i>
		22 — 4		<i>Iva frutescens</i>
		22 — 4		<i>Distichlis spicata</i>
		22 — 4		<i>Solidago sp.</i>
		22 — 4		<i>Typha sp.</i>
Clear Lake		22 — 5		<i>Spartina patens</i> (dominant)
	GHGH 7521 1	22 — 5		<i>Distichlis spicata</i> (co-dominant)
		22 — 5		<i>Scirpus maritimus</i>
		22 — 5		<i>Aster sp.</i>
		22 — 5		<i>Iva frutescens</i>
		22 — 5		<i>Suaeda sp.</i>
		22 — 5		<i>Spartina alterniflora</i> (creek margins)
Morgans Point				

APPENDIX B (cont.)

General Location	Site Number on Photo (Aerial photo + location No.)	Site number		Prevalent Species
		Quad	Site	
		No.	No.	
GHCD 7469 1		26 — 1		
low marsh		26 — 1	<i>Juncus roemerianus</i>	
		26 — 1	<i>Typha sp.</i>	
		26 — 1	<i>Scirpus maritimus</i>	
		26 — 1	<i>Distichlis spicata</i>	
high marsh		26 — 1	<i>Paspalum vaginatum</i>	
		26 — 1	<i>Spartina patens</i>	
		26 — 1	<i>Spartina spartinae</i>	
		26 — 1	<i>Iva frutescens</i>	
		26 — 1	<i>Cynodon dactylon</i>	
		26 — 1	<i>Borrichia frutescens</i>	
		26 — 1	<i>Andropogon glomeratus</i>	
		26 — 1	<i>Solidago sp.</i>	
		26 — 1	<i>Aster sp.</i>	
		26 — 1	<i>Phragmites australis</i>	
		26 — 1	<i>Arundo donax</i>	
		26 — 1	<i>Sesbania sp.</i>	
		26 — 1	<i>Andropogon sp.</i>	
		26 — 1	<i>Baccharis sp.</i>	
forested area in adjacent upland		26 — 1	<i>Cetis laevigata</i>	
		26 — 1	<i>Ulmus crassifolia</i>	
		26 — 1	<i>Ilex vomitoria</i>	
		26 — 1	<i>Carya illinoensis</i>	
		26 — 1	<i>Sapium sebiferum</i>	
		26 — 1	<i>Quercus nigra</i>	
		26 — 1	<i>Quercus phellos</i>	
GHCD 7469 7		26 — 2	lower area- <i>Spartina alterniflora</i>	
		26 — 2	<i>Suaeda</i>	
		26 — 2	<i>Heliotropium</i>	
		26 — 2	<i>Salicornia</i>	
		26 — 2	<i>Batis</i>	
		26 — 2	higher area- <i>Spartina patens</i>	
		26 — 2	<i>Spartina patens</i>	
		26 — 2	<i>Limonium nashii</i>	
		26 — 2	<i>Tamarix</i>	
		26 — 2	<i>Machaeranthera phyllocephala</i>	
		26 — 2	<i>Ambrosia psilostachya</i>	
		26 — 2	<i>Acacia angustissima</i>	
		26 — 2	<i>Phyla lanceolata</i>	
		26 — 2	<i>Eustachys petraea</i>	
		26 — 2	<i>Spiranthes ovalis</i>	
		26 — 2	<i>Juncus roemerianus</i>	
		26 — 2	<i>Desmodium canadense</i>	
		26 — 2	<i>Medicago minima</i>	
GHCD 7469 6		26 — 3	<i>Spartina alterniflora</i>	
		26 — 3	<i>Scirpus maritimus</i>	
		26 — 3	Higher berms- <i>Spartina patens</i>	
		26 — 3	<i>Borrichia frutescens</i>	
		26 — 3	<i>Iva frutescens</i>	
		26 — 3	<i>Lycium carolinianum</i>	
		26 — 3	<i>Alternanthera philoxeroides</i>	
GHCD 7469 5		26 — 4	<i>Spartina alterniflora</i>	
GHCD 7469 4b		26 — 5	<i>Spartina alterniflora</i>	
GHCD 7469 4a		26 — 6	<i>Distichlis spicata</i>	
		26 — 6	<i>Borrichia frutescens</i>	
		26 — 6	<i>Heliotropum curassivicum</i>	

APPENDIX B (cont.)

General Location	Site Number on Photo (Aerial photo + location No.)	General descriptions	Site number Quad Site No. No.	Prevalent Species
Houston Point				
	GHCD 7468 1	low marsh	26 — 7	<i>Spartina alterniflora</i> (dominant over whole area)
			26 — 7	<i>Distichlis spicata</i>
		high marsh	26 — 7	<i>Spartina spartinae</i> (fringes low marsh)
			26 — 7	<i>Spartina patens</i>
			26 — 7	<i>Borrichia frutescens</i>
			26 — 7	<i>Lycium carolinianum</i>
			26 — 7	<i>Iva frutescens</i>
			26 — 7	<i>Aster subulatus</i>
		fresher water drainage zone	26 — 7	<i>Paspalum vaginatum</i>
			26 — 7	<i>Scirpus maritimus</i>
	GHCD 7468 2	high marsh	26 — 8	<i>Iva frutescens</i>
			26 — 8	<i>Spartina spartinae</i>
			26 — 8	<i>Spartina patens</i>
			26 — 8	<i>Phragmites australis</i>
			26 — 8	<i>Arundo donax</i>
			26 — 8	<i>Solidago</i> sp.
			26 — 8	<i>Typha</i> sp.
		lower marsh along channel	26 — 8	<i>Scirpus maritimus</i>
			26 — 8	<i>Spartina alterniflora</i>
		shrubs	26 — 8	<i>Celtis laevigata</i>
			26 — 8	<i>Parkinsonia aculeata</i>
			26 — 8	<i>Baccharis halimifolia</i>
	GHCD 7468 3		26 — 9	
	Transitional assemblage		26 — 9	<i>Iva frutescens</i>
	(east side of highway)		26 — 9	<i>Aster</i> sp.
			26 — 9	<i>Lycium carolinianum</i>
			26 — 9	<i>Baccharis</i> sp.
	(west side of highway)		26 — 9	<i>Iva frutescens</i>
			26 — 9	<i>Baccharis halimifolia</i>
			26 — 9	<i>Setaria</i> sp.
			26 — 9	<i>Andropogon glomeratus</i>
			26 — 9	<i>Solidago</i> sp.
			26 — 9	<i>Aster</i> sp.
	more abundant off levee		26 — 9	<i>Scirpus maritimus</i>
			26 — 9	<i>Distichlis spicata</i>
			26 — 9	<i>Spartina patens</i>
			26 — 9	<i>Spartina spartinae</i>
			26 — 9	<i>Lycium carolinianum</i>
San Jacinto Park	GHEF 7496 1		27 — 1	<i>Iva frutescens</i> dominant
			27 — 1	<i>Spartina patens</i>
			27 — 1	<i>Spartina alterniflora</i>
			27 — 1	<i>Borrichia frutescens</i>
			27 — 1	<i>Sesuvium portulacastrum</i>
			27 — 1	<i>Spartina spartinae</i>
		shrubs	27 — 1	<i>Solidago</i> sp.
			27 — 1	<i>Parkinsonia aculeata</i>
			27 — 1	<i>Celtis laevigata</i>
			27 — 1	<i>Ulmus crassifolia</i>
			27 — 1	<i>Baccharis halimifolia</i>
Trinity River Delta	GHAB 7451 4a		28 — 1	<i>Scirpus olneyi</i>
			28 — 1	<i>Panicum dichotomiflorum</i>
			28 — 1	<i>Echinochloa crusgalli</i>

APPENDIX B (cont.)

General Location	Site Number on Photo (Aerial photo + location No.) General descriptions	Site number Quad Site No. No.	Prevalent Species
		28 — 1	<i>Bacopa monnieri</i>
		28 — 1	<i>Eleocharis parvula</i>
		28 — 1	<i>Eleocharis sp.</i>
GHAB 7451 9e (SEE TRANSECT 28-2, APP. B)		28 — 2	<i>Edge of Eleocharis</i>
		28 — 2	<i>Bacopa monnieri</i> 60%
		28 — 2	<i>Eleocharis sp.</i>
		28 — 2	<i>Polygonum hydropiperoides</i>
		28 — 2	<i>Zizaniopsis miliacea</i>
		28 — 2	<i>Crinum americanum</i>
		28 — 2	<i>Paspalum vaginatum?</i>
GHAB 7451 9d		28 — 3	Tall <i>Eleocharis</i> assemblage 90%
		28 — 3	<i>Polygonum hydropiperoides</i>
		28 — 3	<i>Scirpus olneyi</i>
		28 — 3	<i>Bacopa monnieri</i>
Transition zone between 28-3 and higher assemblage of 28-4 listed below		28 — 4	tall grass <i>Spartina patens?</i>
		28 — 4	<i>Paspalum vaginatum</i>
		28 — 4	<i>Polygonum hydropiperoides</i>
		28 — 4	<i>Cyperus articulatus</i>
		28 — 4	<i>Eleocharis sp.</i>
		28 — 4	<i>Alternanthera philoxeroides</i>
GHAB 7451 9c		28 — 4	Tall grass assemblage <i>Spartina patens?</i>
		28 — 4	<i>Setaria geniculata</i>
		28 — 4	<i>Alternanthera philoxeroides</i>
		28 — 4	<i>Cyperus articulatus</i>
		28 — 4	<i>Lycium carolinianum</i>
GHAB 7451 9b		28 — 5	<i>Panicum repens</i>
		28 — 5	<i>Alternanthera philoxeroides</i>
		28 — 5	<i>Polygonum hydropiperoides</i>
		28 — 5	others collected <i>Physostegia intermedia</i>
		28 — 5	<i>Iva annua</i>
GHAB 7451 9a Transect from edge of into backmarsh		28 — 6	<i>Salix nigra</i>
		28 — 6	<i>Sapium sebiferum</i>
		28 — 6	<i>Phragmites australis</i>
GHAB 7452 4a		28 — 7	<i>Spartina patens</i> (co-dominant)
		28 — 7	<i>Paspalum vaginatum</i> (co-dominant)
		28 — 7	<i>Spartina spartinae</i>
		28 — 7	<i>Cyperus articulatus</i>
		28 — 7	<i>Borrichia frutescens</i>
GHEF 7501 4a		28 — 8	<i>Alternanthera philoxeroides</i>
GHEF 7501 4b		28 — 9	<i>Scirpus olneyi/barren flat</i>
GHEF 7501 3b		28 — 10	<i>Alternanthera philoxeroides</i> 90%
		28 — 10	<i>Crinum americanum</i>
Trinity Delta		28 — 11	<i>Phragmites australis</i>
	GHEF 7501 3a	28 — 11	<i>Salix nigra</i>
		28 — 11	<i>Sapium sebiferum</i>
		28 — 11	<i>Alternanthera philoxeroides</i>
		28 — 11	<i>Celtis laevigata</i>
		28 — 11	<i>Ipomea tricolor</i>
		28 — 11	<i>Panicum repens</i>
		28 — 11	<i>Hymenocallis caroliniana</i>

APPENDIX B (cont.)

General Location	Site Number on Photo (Aerial photo + location No.)	Site number Quad Site No. No.	Prevalent Species
General descriptions			
		28 — 11	<i>Alternanthera philoxeroides</i>
		28 — 11	<i>Iva frutescens</i>
		28 — 11	<i>Polygonum hydropiperoides</i>
North of Lake Anahuac			
	GHAB 7452 1a	28 — 12	<i>Typha sp.</i>
		28 — 12	<i>Eichhornia crassipes</i>
		28 — 12	<i>Lemna sp.</i>
		28 — 12	<i>Juncus roemerianus</i>
		28 — 12	<i>Bacopa monnieri</i>
		28 — 12	<i>Scirpus americanus</i>
		28 — 12	<i>Cyperus articulatus</i>
		28 — 12	<i>Spartina patens</i>
		28 — 12	<i>Sesbania</i>
	GHAB 7451 6	29 — 1	<i>Paspalum vaginatum</i>
		29 — 1	<i>Spartina patens</i>
		29 — 1	<i>Eleocharis sp.</i>
		29 — 1	<i>Spartina patens</i>
		29 — 1	<i>Paspalum lividum</i>
		29 — 1	<i>Cyperus articulatus</i>
		29 — 1	<i>Eleocharis parvula</i>
		29 — 1	<i>Cynodon dactylon</i>
		29 — 1	<i>Polygonum sp.</i>
		29 — 1	<i>Lycium carolinianum</i>
		29 — 1	<i>Aster tenuifolius</i>
	GHAB 7451 5	29 — 2	<i>Alternanthera philoxeroides</i>
		29 — 2	<i>Sagittaria falcata</i>
		29 — 2	<i>Sagittaria lancifolia</i>
		29 — 2	<i>Zizaniopsis miliacea</i>
	GHAB 7451 7	29 — 3	<i>Cyperus articulatus</i>
		29 — 3	<i>Scirpus californicus</i>
		29 — 3	<i>Zizaniopsis miliacea</i>
		29 — 3	<i>Sagittaria falcata</i>
		29 — 3	<i>Phragmites australis</i>
		29 — 3	<i>Alternanthera philoxeroides</i>
		29 — 3	<i>Polygonum sp.</i>
		29 — 3	<i>Aster spinosus (higher margins)</i>
		29 — 3	<i>Lycium carolinianum (scattered)</i>
		—	
	GHAB 7451 11a	29 — 4	<i>Phragmites australis</i>
		29 — 4	<i>Sapium sebiferum</i>
		29 — 4	<i>Crinum americanum</i>
		29 — 4	<i>Alternanthera philoxeroides</i>
		29 — 4	<i>Panicum dichotomiflorum</i>
		29 — 4	<i>Echinochloa crusgalli</i>
	GHAB 7451 11b	29 — 5	<i>Alternanthera philoxeroides</i>
Trinity Delta			
	GHAB 7451 10a levee woodlands	29 — 6	<i>Celtis laevigata</i>
		29 — 6	<i>Aster spinosus</i>
		29 — 6	<i>Sapium sebiferum</i>
		29 — 6	<i>Cynodon dactylon</i>
	GHAB 7451 10b	29 — 7	<i>Cynodon dactylon</i>
		29 — 7	<i>Paspalum vaginatum?</i>
		29 — 7	<i>Cyperus articulatus</i>
		29 — 7	<i>Juncus effusus</i>
		29 — 7	<i>Lycium carolinianum</i>

APPENDIX B (cont.)

General Location	Site Number on Photo (Aerial photo + location No.)	Site number		Prevalent Species
		Quad No.	Site No.	
	General descriptions			
		29 — 7	<i>Eleocharis</i> sp.?	
		29 — 7	<i>Polygonum hydropiperoides</i>	
		29 — 7	<i>Setaria magna</i>	
		29 — 7	<i>Bacopa monnieri</i>	
	GHAB 7451 10c	29 — 8	<i>Spartina patens</i>	
		29 — 8	<i>Polygonum hydropiperoides</i>	
		29 — 8	<i>Cynodon dactylon</i>	
	GHAB 7451 10d	29 — 9	<i>Spartina patens</i>	
		29 — 9	<i>Scirpus californicus</i> (around ponds)	
		29 — 9	others collected-	
		29 — 9	<i>Sisyrinchium exile</i>	
		29 — 9	<i>Hymenocallis caroliniana</i>	
		29 — 9	<i>Physostegia intermedia</i>	
Trinity River	GHAB 7451 8	29 — 10		
		29 — 10	<i>Taxodium distichum</i>	
		29 — 10	<i>Salix nigra</i>	
		29 — 10	<i>Celtis laevigata</i>	
		29 — 10	<i>Cephaelanthus occidentalis</i>	
		29 — 10	<i>Ulmus crassifolia</i>	
		29 — 10	<i>Sapium sebiferum</i>	
		29 — 10	<i>Sabal minor</i>	
		29 — 10	<i>Carya</i> sp.	
		29 — 10	<i>Sesbania</i> sp.	
	In water	29 — 10	<i>Eichhornia crassipes</i>	
		29 — 10	<i>Lemna</i> sp.	
	Edge of forested area	29 — 10	<i>Alternanthera philoxeroides</i>	
		29 — 10	<i>Paspalum lividum</i>	
		29 — 10	<i>Cyperus articulatus</i>	
		29 — 10	<i>Panicum dichotomiflorum</i>	
		29 — 10	<i>Polygonum</i> sp.	
		29 — 10	<i>Rhynchospora</i> sp.	
Tributary near	GHAB 7446 8	31 — 1	<i>Salix nigra</i> along stream	
	River Terrace Park	31 — 1	<i>Pinus</i> sp.	
		31 — 1	<i>Quercus nigra</i>	
		31 — 1	<i>Quercus phellos</i>	
		31 — 1	<i>Ulmus crassifolia</i>	
		31 — 1	<i>Celtis laevigata</i>	
		31 — 1	<i>Liquidambar styraciflua</i>	
		31 — 1	<i>Ilex vomitoria</i>	
	GHAB 7446 3	31 — 2	<i>Typha</i> sp.	
	dead trees include	31 — 2	<i>Taxodium distichum</i>	
	GHEF 7495 1	31 — 3	<i>Phragmites australis</i>	
		31 — 3	<i>Spartina alterniflora</i>	
		31 — 3	<i>Iva frutescens</i>	
		31 — 3	<i>Eleocharis</i> sp.	
		31 — 3	<i>Bacopa monnieri</i>	
		31 — 3	<i>Spartina patens</i>	
		31 — 3	<i>Sesbania</i> sp.	
		31 — 3	<i>Solidago</i> sp.	
		31 — 3	<i>Aster</i> sp.	
		31 — 3	<i>Sesuvium portulacastrum</i>	
		31 — 3	<i>Cynodon dactylon</i>	
		31 — 3	<i>Borrichia frutescens</i>	
		31 — 3	<i>Andropogon glomeratus</i>	
		31 — 3	<i>Ambrosia</i> sp.	
		31 — 3	<i>Baccharis halimifolia</i>	

## APPENDIX B (cont.)

General Location	Site Number on Photo (Aerial photo + location No.) General descriptions	Site number Quad No.	Site number Site No.	Prevalent Species
shrub/forest		31 — 3		<i>Tamarix gallica</i>
		31 — 3		<i>Ulmus crassifolia</i>
		31 — 3		<i>Salix nigra</i>
		31 — 3		<i>Celtis laevigata</i>
GHAB 7446 6		31 — 4		
Park water body		31 — 4		<i>Taxodium distichum</i>
Margin of water		31 — 4		<i>Bacopa monnieri</i>
Sandy Lake		31 — 4		<i>Spartina patens</i>
		31 — 4		<i>Bacopa monnieri</i>
		31 — 4		<i>Sesuvium sp. ?</i>
		31 — 4		<i>Paspalum vaginatum</i>
		31 — 4		<i>Cynodon dactylon</i>
shrub/forest		31 — 4		<i>Salix nigra</i>
No bald cypress		31 — 4		<i>Quercus nigra</i>
		31 — 4		<i>Celtis laevigata</i>
		31 — 4		<i>Sapium sebiferum</i>
GHAB 7446 1d		31 — 5		<i>Scirpus californicus</i>
		31 — 5		<i>Eleocharis sp.</i>
		31 — 5		<i>Typha sp.</i>
		31 — 5		<i>Alternanthera philoxeroides</i>
GHAB 7446 1c		31 — 6		<i>Ulmus crassifolia</i>
forested area toward river		31 — 6		<i>Celtis laevigata</i>
		31 — 6		<i>Sabal minor</i>
		31 — 6		<i>Sapium sebiferum</i>
		31 — 6		<i>Ilex vomitoria</i>
		31 — 6		<i>Liquidambar styraciflua</i>
		31 — 6		<i>Quercus nigra</i>
		31 — 6		<i>Quercus phellos</i>
		31 — 6		<i>Salix nigra</i>
		31 — 6		<i>Carya aquatica</i>
		31 — 6		<i>Pinus sp.</i>
		31 — 6		<i>Taxodium distichum</i>
GHAB 7446 1b		31 — 7		<i>Typha sp.</i>
		31 — 7		<i>Polygonum hydropiperoides-collected</i>
		31 — 7		<i>Alternanthera philoxeroides</i>
		31 — 7		<i>Cyperus articulatus</i>
		31 — 7		<i>Aster subulatus-collected</i>
		31 — 7		<i>Solidago sempervirens-collected</i>
		31 — 7		<i>Panicum dichotomiflorum</i>
scrub/shrubs fringing marsh		31 — 7		<i>Quercus virginiana</i>
		31 — 7		<i>Ulmus crassifolia</i>
		31 — 7		<i>Sapium sebiferum</i>
GHEF 7496 2		31 — 8		<i>Spartina alterniflora</i>
		31 — 8		<i>Phragmites australis</i>
		31 — 8		<i>Iva frutescens</i>
		31 — 8		<i>Colocasia antiquorum</i>
		31 — 8		<i>Typha sp.</i>
		31 — 8		<i>Eleocharis sp.</i>
		31 — 8		<i>Solidago sp.</i>
		31 — 8		<i>Sesbania sp.</i>
		31 — 8		<i>Salix nigra</i>
		31 — 8		<i>Celtis laevigata</i>
		31 — 8		<i>Ilex vomitoria</i>
		31 — 8		<i>Sapium sebiferum</i>

## APPENDIX B (cont.)

General Location	Site Number on Photo (Aerial photo + location No.)	Site number Quad Site No. No.	Prevalent Species
San Jacinto River	GHAB 7446 1a	31 — 9	<i>Typha sp.</i>
	GHAB 7446 5	31 — 10	<i>Iva frutescens dominant</i>
		31 — 10	<i>Spartina spartinae</i>
		31 — 10	<i>Eleocharis parvula ?</i>
		31 — 10	<i>Spartina patens</i>
		31 — 10	<i>Aster sp.</i>
		31 — 10	<i>Sesuvium portulacastrum</i>
		31 — 10	<i>Paspalum vaginatum</i>
In ditch across frontage rd.		31 — 10	<i>Scirpus maritimus</i>
Shrubs/forest		31 — 10	<i>Typha sp.</i>
		31 — 10	<i>Pinus sp.</i>
		31 — 10	<i>Ulmus crassifolia</i>
		31 — 10	<i>Ilex vomitoria</i>
		31 — 10	<i>Liquidambar styraciflua</i>
		31 — 10	<i>Sapium sebiferum</i>

## Appendix C

APPENDIX C. Elevation Transects.


Galveston Bay Elevation Transect: Site No. 7-1

Hoskins Mound Transect 1

11/14/90  
 Instrument height (m): 1.470  
 Ground elevation (m): 0.000

Shot	Bearing °	Bearing "	Height (m)	Top (m)	Bottom (m)	Bearing (°)	Distance (m)	Relative Elevation (m)	X (m)	Y (m)	Line Distance (m)
0	0	0	0	1.470	1.470	1.470	0.000	0	0.000	0.0	0.0
4	200	58	10	1.445	1.540	1.340	200.969	20	0.025	-7.2	20.0
3	198	11	52	1.525	1.690	1.370	198.198	32	-0.055	-10.0	32.0
2	198	44	40	1.350	1.575	#N/A	198.744	45	0.120	-14.5	45.0
1	198	4	40	0.650	0.895	#N/A	198.078	49	0.820	-15.2	-46.6
											49.0


Hoskins Mound Transect 1 Shotpoints



Site No. 7-1

APPENDIX C (cont.)

Hoskins Mound Transect 1



APPENDIX C (cont.)

Hoskins Mound Transect 1: Site No. 7-1

Station No.

1	Edge of gravel road
2	<i>Juncus roemerianus, Spartina patens, Polygonum sp., Cyperus sp., others</i>
2-4	<i>Juncus roemerianus</i>
4-Instru.	<i>Spartina spartinae</i> (90%), <i>Spartina patens</i> (10%)
Instru.	<i>Spartina spartinae</i>

APPENDIX C (cont.)

Galveston Bay Elevation Transect: Site No. 7-1


Hoskins Mound Transect 2

11/14/90                  1.470  
 Instrument height (m):      0.000  
 Ground elevation (m):        0.000

Shot	Bearing °	Height (m)	Top (m)	Bottom (m)	Bearing °	Distance (m)	Relative Elevation (m)	X (m)	Y (m)	Line Distance (m)
0	0 0 0	0	1.470	1.470	1.470	0.00	0	0.000	0.0	0.0
5	127 19 50	50	1.470	1.505	1.445	127.33	6	0.000	4.8	-3.6
6	124 56 20	20	1.460	1.620	1.300	124.94	32	0.010	26.2	-18.3
11	125 19 20	20	1.555	1.750	1.360	125.32	39	-0.085	31.8	-22.5
12	125 10 50	50	1.530	1.755	1.300	125.18	45	-0.060	37.2	-26.2
13	125 32 40	40	1.370	1.640	#N/A	125.54	54	0.100	43.9	-31.4
14	124 28 0	0	1.920	2.220	#N/A	124.47	60	-0.450	49.5	-34.0
15	123 52 40	40	1.375	1.720	#N/A	123.88	69	0.095	57.3	-38.5
										69.0

**APPENDIX C (cont.)**


**Hoskins Mound Transect 2 Shotpoints**



Site No. 7-1

Hoskins Mound Transect 2

APPENDIX C (cont.)



APPENDIX C (cont.)

Hoskins Mound Transect 2: Site No. 7-1

Station No.	
Instru. to 5	<i>Spartina spartinae</i>
5 to 6	<i>Juncus roemerianus</i>
6	<i>Spartina spartinae, Spartina patens, Setaria sp., Juncus roemerianus, Andropogon glomeratus, Solidago sp.</i>
11	<i>Juncus roemerianus</i>
12	<i>Spartina spartinae, Andropogon glomeratus, Fimbristylis castanea, Aster sp., Borrichia, annuals</i>
13	<i>Juncus roemerianus, Andropogon glomeratus, Paspalum laeve, Setaria sp.</i>
14	<i>Typha sp.</i>
15	<i>Spartina spartinae, Spartina patens, Eleocharis sp., Setaria sp., Fimbristylis castanea, Andropogon glomeratus, Solidago sp.</i>

APPENDIX C (cont.)


Galveston Bay Elevation Transect: Site No. 7-1

Hoskins Mound Transect 3

11/14/90                          1.470  
 Instrument height (m):        0.000  
 Ground elevation (m):

Shot	Bearing		Height (m)	Top (m)	Bottom (m)	Decimal Bearing (°)	Distance Elevation (m)	Relative Elevation (m)	X		Y		Line Distance (m)
	°	"							(m)	(m)	(m)	(m)	
0	0	0	0	1.470	1.470	1.470	0.00	0	0.000	0.0	0.0	0.0	
7	135	3	30	1.480	1.625	1.335	135.06	29	-0.010	20.5	-20.5	29.0	
8	134	57	50	1.830	1.980	1.680	134.96	30	-0.360	21.2	-21.2	30.0	
9	135	6	10	2.080	2.260	1.905	135.10	35	-0.610	25.1	-25.1	35.5	
10	134	45	40	1.360	1.555	1.165	134.76	39	0.110	27.7	-27.5	39.0	


**Hoskins Mound Transect 3 Shotpoints**



**APPENDIX C (cont.)**

Site No. 7-1

**Hoskins Mound Transect 3**



APPENDIX C (cont.)

Hoskins Mound Transect 3: Site No. 7-1

Station No.

Instru. to 7

7

*Juncus roemerianus*

*Spartina patens, Spartina spartinae, Setaria sp., Andropogon glomeratus, Juncus roemerianus, Solidago sp.*

8

*Typha sp. (Water)*

9

*Typha sp. (Water)*

10

*Spartina spartinae, Spartina patens, Andropogon glomeratus, Setaria sp., Juncus roemerianus, Polygonum sp.*

APPENDIX C (cont.)

Galveston Bay Elevation Transect: Site No. 7-1


Hoskins Mound Transect 4

11/14/90  
 Instrument height (m): 1.470  
 Ground elevation (m): 0.000

Shot	Bearing °	Bearing "	Height (m)	Top (m)	Bottom (m)	Bearing (°)	Distance Elevation (m)	X (m)	Y (m)	Line Distance (m)
0	0	0	0	1.470	1.470	0.00	0	0.000	0.0	0.0
16	87	21	40	1.500	1.538	1.462	87.36	8	-0.030	7.6
17	92	39	50	1.525	1.640	1.410	92.66	23	-0.055	23.0
18	92	46	20	1.595	1.805	1.390	92.77	42	-0.125	41.5
19	93	34	0	1.620	1.860	1.375	93.57	48	-0.150	47.9
20	94	27	30	1.510	1.850	1.170	94.46	68	-0.040	67.8
21	94	43	30	1.280	1.750	0.810	94.73	94	0.190	93.7
										-7.7
										94.0

**APPENDIX C (cont.)**


**Hoskins Mound Transect 4 Shotpoints**



Site No. 7-1

Hoskins Mound Transect 4

APPENDIX C (cont.)



APPENDIX C (cont.)

Hoskins Mound Transect 4: Site No. 7-1

Station No.

Instru.	<i>Spartina spartinae</i>
16 to 17	<i>Juncus roemerianus</i>
17 to 18	Barren flats, <i>Spartina patens</i> patches, <i>Eleocharis</i> sp., <i>Paspalum vaginatum</i>
18	<i>Paspalum vaginatum</i> , <i>Spartina patens</i> patches
19	Edge of <i>Spartina patens</i> patch
19 to 20	Mixtures of vegetation and barren flat
20	<i>Spartina spartinae</i> (short), scattered <i>Salicornia</i> sp., <i>Limonium nashii</i> , <i>Fimbristylis castanea</i> , <i>Panicum</i> sp., <i>Cyperus articulatus</i> , algae mats
21	Prairie assemblage, <i>Spartina spartinae</i> , <i>Setaria</i> sp., <i>Aristida</i> sp., <i>Solidago</i> sp., <i>Andropogon glomeratus</i> , short <i>Distichlis spicata</i> , <i>Paspalum vaginatum</i>

**APPENDIX C (cont.)**

**Galveston Bay Elevation Transect: Site No. 7-1**

Hoskins Mound Transect 5


11/14/90

Instrument height (m): 1.470

Ground elevation (m): 0.000

Shot	Bearing			Height (m)	Top (m)	Bottom (m)	Decimal Bearing (°)	Distance (m)	Relative Elevation (m)	X (m)	Y (m)	Line Distance (m)
	°	'	"									
0	0	0	0	1.470	1.470	1.470	0.00	0	0.000	0.0	0.0	0.0
22	30	44	10	1.445	1.555	1.340	30.74	21	0.025	11.0	18.5	21.5
23	32	56	20	1.390	1.690	1.090	32.94	60	0.080	32.6	50.4	60.0
24	36	47	20	1.600	2.550	#N/A	36.79	190	-0.130	113.8	152.2	190.0


Hoskins Mound Transect 5 Shotpoints



Site No. 7-1

Hoskins Mound Transect 5

APPENDIX C (cont.)



APPENDIX C (cont.)

Hoskins Mound Transect S: Site No. 7-1

Station No.

Instru.

22

Edge of tall *Spartina spartinae* (dominant)—*Juncus roemerianus* mix, Into short  
*Spartina spartinae*—*Spartina patens* assemblage, scattered *Cyperus articulatus*,  
*Fimbristylis castanea*, *Suaeda* sp., *Borrichia frutescens*, composites

23

Edge of prairie, short *Spartina spartinae*, *Spartina patens*, *Fimbristylis castanea*,  
*Panicum* sp., *Borrichia frutescens*, *Andropogon glomeratus*, *Aristida* sp., *Setaria*  
sp., *Aster* sp., composites, barren spots along trails

24

Edge of Prairie, short *Spartina spartinae*, *Distichlis spicata*, scattered *Fimbristylis*  
*castanea*, *Panicum* sp., *Borrichia frutescens*, (Damp soils in lows)

APPENDIX C (cont.)

Galveston Bay Elevation Transect: Site No. 7-1

Hoskins Mound Transect 6


11/14/90

Instrument height (m): 1.560 (same location as T5, shot 24)

Ground elevation (m): -0.040 (relative to instrument position 1)

Shot	Bearing °	Bearing "	Height (m)	Top (m)	Bottom (m)	Bearing (°)	Distance (m)	Elevation (IP1, m)	Relative Elevation (m)	X (m)	Y (m)	Line Distance (m)
30	226	25	20	0.560	1.685	#N/A	226.42	0.960	-163.0	-155.1	-225	
0	0	0	0	1.560	1.560	1.560	0.00	0	-0.040	0.0	0.0	0
25	43	53	50	1.640	1.760	1.520	43.90	24	-0.120	16.6	17.3	24
26	42	15	30	2.180	2.330	2.030	42.26	30	-0.660	20.2	22.2	30
27	43	6	20	1.620	1.800	1.440	43.11	36	-0.100	24.6	26.3	36
28	42	45	0	1.690	2.030	1.350	42.75	68	-0.170	46.2	49.9	68
29	42	41	20	1.680	2.220	1.130	42.69	109	-0.160	73.9	80.1	109


**Hoskins Mound Transect 6 Shotpoints**



Site No. 7-1

Hoskins Mound Transect 6

APPENDIX C (cont.)



APPENDIX C (cont.)

Hoskins Mound Transect 6: Site No. 7-1

Station No.	Instru.	
24		Edge of Prairie, short <i>Spartina spartinae</i> , <i>Distichlis spicata</i> , scattered <i>Fimbristylis castanea</i> , <i>Panicum</i> sp., <i>Borrichia frutescens</i> , (Damp soils in lows)
25		Short <i>Spartina spartinae</i> , <i>Spartina patens</i> , <i>Setaria</i> sp., <i>Andropogon glomeratus</i> , <i>Solidago</i> sp.
26		<i>Typha</i> sp. (Water 30 cm)
27		Short <i>Spartina spartinae</i> , <i>Spartina patens</i> , <i>Setaria</i> sp., <i>Andropogon glomeratus</i> , <i>Solidago</i> sp.
28		Tall <i>Spartina spartinae</i> — <i>Spartina patens</i> , <i>Fimbristylis castanea</i> , some <i>Juncus roemerianus</i>
29		Channel assemblage, tall <i>Spartina patens</i> (up to 75-90%)— <i>Juncus roemerianus</i> (up to 50-60% locally), <i>Cyperus articulatus</i>

**APPENDIX C (cont.)**

**Galveston Bay Elevation Transect: Site No. 3-2**

Hoskins Mound Transect 7

12/12/90


Instrument height (m): 1.560

Ground elevation (m): 0.000 (relative to instrument position)

Shot	Bearing			Height (m)	Top (m)	Bottom (m)	Decimal		Relative		X (m)	Y (m)	Line Distance (m)
	°	'	"				Bearing (°)	Distance (m)	Elevation (IP1, m)				
1	60	18	50	1.450	2.220	#N/A	60.31	154	0.110	133.8	76.3	154	
2	60	30	40	1.470	2.170	0.780	60.51	139	0.090	121.0	68.4	139	
3	60	48	0	1.520	2.140	0.920	60.80	122	0.040	106.5	59.5	122	
4	60	22	10	1.520	2.090	0.970	60.37	112	0.040	97.4	55.4	112	
5	60	22	0	1.560	1.950	1.180	60.37	77	0.000	66.9	38.1	77	
6	60	32	10	1.570	1.810	1.320	60.54	49	-0.010	42.7	24.1	49	
0				1.560	#N/A	#N/A	0.00	#N/A	0.000	0.0	0.0	0	
7	239	51	40	1.620	1.800	1.440	239.86	36	-0.060	-31.1	-18.1	-36	
8	240	6	30	1.610	1.850	1.360	240.11	49	-0.050	-42.5	-24.4	-49	
9	240	5	20	1.590	1.900	1.270	240.09	63	-0.030	-54.6	-31.4	-63	
10	239	59	0	1.690	2.020	1.360	239.98	66	-0.130	-57.1	-33.0	-66	
11	239	59	50	1.740	2.120	1.360	240.00	76	-0.180	-65.8	-38.0	-76	
12	240	32	20	1.740	2.270	1.220	240.54	105	-0.180	-91.4	-51.6	-105	
13	240	26	10	1.860	2.620	1.100	240.44	152	-0.300	-132.2	-75.0	-152	
14	240	16	50	1.900	2.690	1.120	240.28	157	-0.340	-136.3	-77.8	-157	
15	240	25	50	2.050	2.900	1.210	240.43	169	-0.490	-147.0	-83.4	-169	
16	240	22	30	1.700	2.560	0.850	240.38	171	-0.140	-148.6	-84.5	-171	
17	240	22	30	1.810	2.730	0.890	240.38	184	-0.250	-159.9	-91.0	-184	
18	243	34	10	1.780	3.030	#N/A	243.57	250	-0.220	-223.9	-111.3	-250	

APPENDIX C (cont.)


Hoskins Mound Transect 7 Shotpoints



Site No. 3-2

### Hoskins Mound Transect 7

#### APPENDIX C (cont.)



APPENDIX C (cont.)

Hoskins Mound Transect 7: Site No. 3-2

Station No.	
1	<i>Spartina spartinae, Spartina patens, Distichlis spicata</i> Scattered <i>Borrichia frutescens, Iva frutescens, Cyperus articulatus, Cyperus</i> sp.
2	Barren flat, scattered <i>Salicornia bigelovii</i>
3	<i>Distichlis spicata—Spartina patens—Spartina spartinae</i>
4	<i>Spartina patens</i> dominant, <i>Distichlis spicata</i> abundant, scattered <i>Spartina spartinae, Aster tenuifolius, Borrichia frutescens</i>
5	<i>Distichlis spicata—Spartina spartinae—Spartina patens</i>
6	<i>Spartina spartinae</i> dominant, some <i>Distichlis, Spartina patens</i> , scattered <i>Salicornia</i>
Instru.	<i>Spartina spartinae—Distichlis spicata</i> , scattered <i>Salicornia, Aster tenuifolius, Lymonium</i>
7	Edge of <i>Distichlis spicata—Spartina spartinae</i> zone, beginning of <i>Monanthochloe littoralis</i>
8	Edge of <i>Monanthochloe</i> dominance, beginning of <i>Spartina spartinae - Distichlis</i> zone
9	<i>Spartina spartinae</i> on rim of flat
10	Flat with <i>Monanthochloe</i> , scattered <i>Distichlis, Salicornia</i> spp., algal mat
11	<i>Distichlis spicata</i>
12	<i>Distichlis spicata</i> , scattered <i>Aster tenuifolius</i> and <i>Salicornia</i>
13	Edge of <i>Spartina alterniflora</i> , some <i>Distichlis</i>
14	Tidal channel, standing water, <i>Ruppia maritima</i>
15	Center of tidal channel (0.5 to 1 m wide) water 9 cm deep
16	<i>Spartina patens</i> dominant (margin of channel)
17	<i>Spartina patens</i> (tall and healthy), scattered <i>Juncus roemerianus, Distichlis spicata, Aster tenuifolius</i>
18	<i>Spartina patens—Distichlis spicata</i> zone, scattered <i>Scirpus maritimus, Juncus roemerianus</i> <i>Iva frutescens</i> abundant toward channel to SW (about 25 m)

**APPENDIX C (cont.)**

**Galveston Bay Elevation Transect: Site No. 3-3**

Follets Island Transect 1

11/14/90


Instrument height (m): 1.530

Ground elevation (m): 0.000 (relative to instrument position)

Shot	Bearing			Height (m)	Top (m)	Bottom (m)	Decimal Bearing (°)	Distance (m)	Relative Elevation		X (m)	Y (m)	Line Distance (m)
	°	'	"						X	Y			
31	138	26	30	0.550	1.010	#N/A	138.44	92	0.980	61.0	-68.8	-92.0	
32	138	20	40	1.110	1.360	0.870	138.34	49	0.420	32.6	-36.6	-49.0	
33	140	19	50	1.330	1.460	1.200	140.33	26	0.200	16.6	-20.0	-26.0	
0	0	0	0	1.530	1.530	1.530	0.00	0	0.000	0.0	0.0	0.0	
34	317	42	10	1.690	1.790	1.590	317.70	20	-0.160	-13.5	14.8	20.0	
35	320	52	0	1.870	2.020	1.720	320.87	30	-0.340	-18.9	23.3	30.0	
36	321	35	50	1.930	2.120	1.750	321.60	37	-0.400	-23.0	29.0	37.0	
37	321	42	0	2.010	2.220	1.790	321.70	43	-0.480	-26.7	33.7	43.0	
38	319	50	20	2.070	2.510	1.620	319.84	89	-0.540	-57.4	68.0	89.0	
39	319	41	40	2.100	2.570	1.620	319.69	95	-0.570	-61.5	72.4	95.0	
40	319	46	40	2.170	2.660	1.680	319.78	98	-0.640	-63.3	74.8	98.0	
41	320	15	40	2.330	2.830	1.820	320.26	101	-0.800	-64.6	77.7	101.0	
42	320	16	30	2.180	2.700	1.660	320.28	104	-0.650	-66.5	80.0	104.0	
43	320	47	0	2.110	2.670	1.560	320.78	111	-0.580	-70.2	86.0	111.0	
44	320	51	10	2.200	2.820	1.570	320.85	125	-0.670	-78.9	96.9	125.0	


**APPENDIX C (cont.)**

**Follets Island Transect 1 Shotpoints**



APPENDIX C (cont.)

Site No. 3-3



APPENDIX C (cont.)

Follets Island Transect 1: Site No. 3-3

Station No.	
1	<i>Paspalum monostachyum, Spartina spartinae, Fimbristylis castanea, Andropogon glomeratus, Hydrocotyle bonariensis, Cyperus</i> sp.
2	Edge of <i>Iva frutescens</i> , <i>Paspalum monostachyum</i> , <i>Andropogon glomeratus</i> , <i>Fimbristylis castanea</i> , scattered <i>Hydrocotyle bonariensis</i> , composites
3	Middle of <i>Iva frutescens</i> — <i>Spartina patens</i> dominance, <i>Spartina spartinae</i> abundant, <i>Paspalum monostachyum</i> , <i>Andropogon glomeratus</i> , <i>Solidago</i> sp., <i>Scirpus americanus</i> , <i>Setaria</i> sp., <i>Borrichia frutescens</i>
Instru.	
4	Trailing edge of <i>Iva frutescens</i> , beginning of <i>Spartina patens</i> dominance with <i>Distichlis spicata</i> mix, scattered <i>Borrichia frutescens</i> , <i>Spatina spartinae</i>
5	Edge of <i>Spartina patens</i> , <i>Distichlis spicata</i> dominant (90%), scattered <i>Limonium nashii</i> , <i>Salicornia</i> sp.
6	Leading edge of <i>Monanthochloe littoralis</i> dominance, gradation with <i>Distichlis spicata</i> zone about 1 m
7	Trailing edge of <i>Monanthochloe</i> , leading edge of algal flat
8	<i>Batis maritima</i> , trailing edge of algal flat
9	<i>Spartina alterniflora</i> — <i>Batis maritima</i>
10	Edge of water, <i>Spartina alterniflora</i> dominance, scattered <i>Distichlis spicata</i>
11	<i>Spartina alterniflora</i> (Water 17 cm)
12	<i>Spartina alterniflora</i> — <i>Distichlis spicata</i>
13	<i>Distichlis spicata</i>
14	<i>Spartina alterniflora</i> (90-95%), <i>Batis maritima</i> (5-10%)

**APPENDIX C (cont.)**

**Galveston Bay Elevation Transect: Site No. 18-9**

Anahuac Wildlife Refuge Transect 1

11/16/90


Instrument height (m): 1.620

Ground elevation (m): 0.000 (relative to instrument position)

Shot	Bearing			Height (m)	Top (m)	Bottom (m)	Decimal Bearing		Distance (m)	Relative Elevation (m)	X (m)	Y (m)	Line Distance (m)
	°	'	"				(°)	(m)					
45	305	16	10	1.670	2.020	1.310	305.27		71	-0.050	-58.0	41.0	-71.0
46	304	43	30	1.590	1.780	1.400	304.73		38	0.030	-31.2	21.6	-38.0
0	0	0	0	1.620	1.620	1.620	0.00		0	0.000	0.0	0.0	0.0
47	124	6	0	1.625	1.680	1.570	124.10		11	-0.005	9.1	-6.2	11.0
48	124	0	30	1.675	1.775	1.575	124.01		20	-0.055	16.6	-11.2	20.0
50	117	46	20	1.800	2.020	1.580	117.77		44	-0.180	38.9	-20.5	44.0
49	124	6	0	1.650	1.870	1.420	124.10		45	-0.030	37.3	-25.2	45.0
51	117	15	10	1.640	1.940	1.340	117.25		60	-0.020	53.3	-27.5	60.0
52	116	38	40	1.750	2.100	1.410	116.64		69	-0.130	61.7	-30.9	69.0
53	115	30	40	1.780	2.180	1.380	115.51		80	-0.160	72.2	-34.5	80.0
54	116	28	40	1.720	2.140	1.300	116.48		84	-0.100	75.2	-37.5	84.0
55	116	50	50	1.720	2.180	1.260	116.85		92	-0.100	82.1	-41.5	92.0
56	116	19	40	1.630	2.200	1.050	116.33		115	-0.010	103.1	-51.0	115.0
57	117	58	30	1.660	2.320	0.990	117.98		133	-0.040	117.5	-62.4	133.0

APPENDIX C (cont.)


Anahuac NWR Transect 1 Shotpoints



Site No. 18-9

Anahuac NWR Transect 1

APPENDIX C (cont.)




APPENDIX C (cont.)

Anahuac NWR Transect 1: Site No. 18-9

Station No.


- |         |   |
|---------|---|
| 1       | <i>Spartina spartinae</i> , <i>Spartina patens</i> , <i>Distichlis spicata</i> , scattered <i>Aster</i> sp.,<br><i>Borrichia frutescens</i> , <i>Iva frutescens</i> (Wet)   |
| 2       | <i>Spartina spartinae</i> (Damp Soil)   |
| Instru. | <i>Spartina spartinae</i> , some <i>Iva frutescens</i>  |
| 3       | <i>Spartina patens</i> — <i>Distichlis spicata</i> dominance, <i>Paspalum vaginatum</i> , trailing<br>edge of <i>Spartina spartinae</i> , scattered <i>Aster</i> sp., <i>Setaria</i> , <i>Cyperus</i> sp.                 |
| 4       | <i>Paspalum vaginatum</i> , (Water 1-2 cm)  |
| 5       | Leading edge of <i>Scirpus olneyi</i> , trailing edge of <i>Paspalum vaginatum</i> and<br><i>Spartina patens</i>  |
| 6       | <i>Scirpus olneyi</i> (Water 20 cm)   |
| 7       | <i>Scirpus olneyi</i> (60%), <i>Spartina patens</i> (40%) (Water 4 cm)  |
| 8       | <i>Scirpus olneyi</i> , <i>Spartina patens</i> (Water 6 cm)   |
| 9       | <i>Scirpus olneyi</i> (90%), <i>Spartina patens</i> , <i>Echinochloa crusgalli</i> , <i>Bacopa monnieri</i><br>(Water 7 cm)   |
| 10      | <i>Distichlis spicata</i> (tall), <i>Spartina patens</i> , scattered <i>Scirpus olneyi</i> (Water 2.5 cm)   |
| 11      | Trailing edge of <i>Scirpus olneyi</i> , leading edge of <i>Distichlis spicata</i> — <i>Spartina</i><br><i>patens</i> dominance, scattered <i>Echinochloa crusgalli</i> , <i>Spartina spartinae</i> ,<br><i>Aster</i> sp. |
| 12      | <i>Spartina patens</i> dominance, abundant <i>Distichlis spicata</i> , scattered <i>Borrichia</i> and<br><i>Aster</i> sp. (Soil Damp)   |
| 13      | <i>Spartina spartinae</i> dominance, <i>Spartina patens</i> , <i>Distichlis spicata</i> , <i>Aster</i> sp.,<br><i>Borrichia frutescens</i> , <i>Cyperus articulatus</i> , <i>Echinochloa crusgalli</i> (Soil Damp)        |

APPENDIX C (cont.)



APPENDIX C (cont.)

Trinity River Delta Transect: Site No. 28-2 to 6



Station No.

- 1 Levee (instr.) River's edge; water level  
Upland assemblage: scattered trees and shrubs including *Salix nigra*, *Sapium sebiferum*; grasses include *Panicum repens* and *Phragmites australis*; forbs include *Iva annua*, *Physostegia intermedia*, others (from station 1 to 2)
- 2 Edge of tall grass assemblage including *Spartina patens*, *Setaria geniculata*, *Alternanthera philoxeroides*, *Cyperus articulatus*, *Lycium carolinianum*  
Bayward edge of tall grass assemblage, beginning of assemblage of *Spartina patens*, *Paspalum vaginatum*? (no inflorescence), *Polygonum hydropiperoides*, *Cyperus articulatus*, *Eleocharis* sp., *Alternanthera philoxeroides*; water 0.5 cm deep
- 3 Continuation of assemblage noted above at station 3  
Edge of dominant, tall *Eleocharis* sp.(90%) (0.8 m tall), *Polygonum hydropiperoides*, *Scirpus olneyi*, *Bacopa monnieri*, *Alternanthera philoxeroides*  
Center of tall *Eleocharis* sp. zone (see 5); water 3 to 8 cm deep
- 4 Beginning of less dense and shorter assemblage of *Bacopa monnieri* (60%) *Eleocharis* sp., *Polygonum hydropiperoides*, *Zizaniopsis miliacea*, *Crinum americanum*, *Paspalum vaginatum*?; water 3 to 8 cm deep
- 5
- 6
- 7